



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 04:47 PM GMT

PDB ID : 2VB6
Title : MYOSIN VI (MD-INSERT2-CAM, DELTA INSERT1) POST-RIGOR STATE
(CRYSTAL FORM 2)
Authors : Menetrey, J.; Llinas, P.; Cicolari, J.; Squires, G.; Liu, X.; Li, A.; Sweeney,
H.L.; Houdusse, A.
Deposited on : 2007-09-06
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	788	
2	B	149	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7271 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYOSIN VI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	724	Total	C	N	O	S	65	0	0
			5857	3741	1008	1080	28			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	547	VAL	GLY	CONFLICT	UNP Q29122
A	572	ARG	ALA	CONFLICT	UNP Q29122
A	573	ASP	TYR	CONFLICT	UNP Q29122
A	714	LEU	VAL	CONFLICT	UNP Q29122
A	721	TYR	SER	CONFLICT	UNP Q29122
A	722	MET	LEU	CONFLICT	UNP Q29122

- Molecule 2 is a protein called CALMODULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	145	Total	C	N	O	S	9	0	0
			1141	700	183	249	9			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	99	PHE	TYR	CONFLICT	UNP P62149
B	120	GLU	GLN	CONFLICT	UNP P62149
B	143	THR	GLN	CONFLICT	UNP P62149
B	147	SER	ALA	CONFLICT	UNP P62149

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Be	F	0	0
			4	1	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Mg 1	0	0

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	4	Total 4	Ca 4	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	211	Total 211	O 211	0	0
7	B	26	Total 26	O 26	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: MYOSIN VI

Chain A:



• Molecule 2: CALMODULIN

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.31Å 107.59Å 178.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.18 – 2.30 46.17 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (46.18-2.30) 99.1 (46.17-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.91 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.210 , 0.253 0.206 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	32.7	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 23.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 62868 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7271	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, BEF, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	3/5980 (0.1%)	0.63	2/8052 (0.0%)
2	B	0.56	1/1153 (0.1%)	0.73	3/1548 (0.2%)
All	All	0.56	4/7133 (0.1%)	0.65	5/9600 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
2	B	1	2
All	All	1	5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	494	GLU	CB-CG	7.55	1.66	1.52
1	A	412	LYS	CA-CB	-6.95	1.38	1.53
2	B	78	ASP	CA-CB	6.91	1.69	1.53
1	A	261	GLU	CA-CB	-5.01	1.43	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	773	ASP	N-CA-C	-7.97	89.49	111.00
2	B	24	ASP	N-CA-C	6.20	127.74	111.00
2	B	78	ASP	CB-CA-C	-5.75	98.89	110.40
1	A	177	ASP	N-CA-C	-5.62	95.81	111.00
2	B	23	GLY	N-CA-C	5.02	125.65	113.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	24	ASP	CA

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	176	GLN	Peptide
1	A	729	LEU	Peptide
1	A	772	SER	Peptide
2	B	23	GLY	Peptide
2	B	57	ALA	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5857	0	5813	49	3
2	B	1141	0	1067	19	0
3	A	27	0	12	0	0
4	A	4	0	0	0	0
5	A	1	0	0	0	0
6	B	4	0	0	0	0
7	A	211	0	0	3	0
7	B	26	0	0	2	0
All	All	7271	0	6892	66	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (66) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:116:LEU:HD21	2:B:124:MET:HE1	1.54	0.89
1:A:811:LYS:HB3	1:A:811:LYS:NZ	1.98	0.77
1:A:811:LYS:HB3	1:A:811:LYS:HZ2	1.50	0.74
1:A:811:LYS:CB	1:A:811:LYS:NZ	2.48	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:554:HIS:HD2	1:A:556:ARG:H	1.37	0.70
1:A:532:GLU:OE2	1:A:542:HIS:HD2	1.75	0.70
1:A:548:HIS:CE1	1:A:559:ILE:HD12	2.33	0.63
1:A:667:ARG:NH2	7:A:2180:HOH:O	2.27	0.63
1:A:811:LYS:CB	1:A:811:LYS:HZ2	2.07	0.63
1:A:554:HIS:HE1	7:A:2123:HOH:O	1.83	0.61
1:A:559:ILE:HG22	1:A:562:LYS:HG3	1.81	0.61
2:B:135:GLN:NE2	7:B:2026:HOH:O	2.33	0.60
1:A:748:ASP:HA	1:A:759:ARG:HG3	1.85	0.59
2:B:20:ASP:OD2	2:B:23:GLY:HA2	2.03	0.58
1:A:211:GLU:OE1	1:A:213:HIS:HE1	1.87	0.57
1:A:73:LEU:HD22	1:A:698:LEU:HD11	1.86	0.57
1:A:707:SER:HB2	1:A:758:PHE:HB2	1.87	0.57
1:A:466:ASN:HD22	1:A:470:GLN:HG2	1.69	0.56
1:A:190:GLU:O	1:A:194:ASN:HB2	2.05	0.56
2:B:63:ILE:HD12	2:B:67:GLU:HB3	1.86	0.55
1:A:581:HIS:HE1	1:A:588:TYR:OH	1.90	0.55
1:A:581:HIS:CE1	1:A:588:TYR:OH	2.61	0.54
1:A:178:ILE:HD11	1:A:222:GLY:HA2	1.89	0.54
2:B:52:ILE:HA	2:B:71:MET:CE	2.39	0.53
1:A:807:LYS:HE3	1:A:808:ASN:OD1	2.10	0.52
2:B:64:ASP:OD1	2:B:66:PRO:HD2	2.10	0.51
1:A:726:LEU:HD13	1:A:781:VAL:HG11	1.93	0.50
1:A:213:HIS:HD2	1:A:448:SER:OG	1.95	0.50
2:B:52:ILE:HG12	2:B:71:MET:HE1	1.93	0.50
2:B:27:ILE:HD12	2:B:32:LEU:HD13	1.94	0.49
2:B:28:THR:HG22	2:B:62:THR:HG22	1.94	0.49
1:A:202:ASN:OD1	1:A:246:HIS:HE1	1.95	0.49
1:A:476:CYS:HA	1:A:651:LEU:HD11	1.95	0.48
1:A:513:ASP:OD2	1:A:556:ARG:HD3	2.12	0.48
2:B:52:ILE:HA	2:B:71:MET:HE2	1.95	0.48
1:A:501:LEU:HD23	1:A:750:LYS:HG2	1.96	0.47
1:A:705:PHE:CZ	1:A:750:LYS:HD3	2.50	0.47
1:A:211:GLU:OE1	1:A:213:HIS:CE1	2.67	0.47
1:A:802:SER:O	1:A:806:LEU:HG	2.15	0.46
2:B:24:ASP:OD2	2:B:24:ASP:C	2.54	0.46
1:A:803:VAL:HG13	2:B:36:MET:HG2	1.98	0.46
1:A:674:LYS:O	1:A:676:THR:HG23	2.16	0.45
2:B:116:LEU:CD2	2:B:124:MET:HE1	2.36	0.45
1:A:811:LYS:HB3	1:A:811:LYS:HZ3	1.80	0.45
2:B:90:ARG:HD2	2:B:90:ARG:HA	1.69	0.45
1:A:554:HIS:CD2	1:A:556:ARG:HG2	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:411:LEU:HD22	1:A:415:GLN:HB3	1.99	0.45
2:B:105:LEU:HD21	2:B:124:MET:HE3	1.97	0.45
1:A:479:LYS:HB3	1:A:655:LEU:HD11	1.99	0.44
2:B:37:ARG:HA	2:B:41:GLN:O	2.16	0.44
1:A:497:GLN:HG3	1:A:498:LYS:N	2.28	0.44
2:B:126:ARG:NH2	7:B:2024:HOH:O	2.48	0.44
1:A:736:LYS:HG2	1:A:751:PHE:CZ	2.52	0.44
2:B:115:LYS:HB2	2:B:115:LYS:HE3	1.88	0.44
1:A:509:VAL:O	1:A:509:VAL:HG23	2.18	0.44
1:A:176:GLN:H	1:A:176:GLN:HG2	1.55	0.43
1:A:84:ASP:OD1	1:A:678:HIS:HE1	2.02	0.43
1:A:762:LYS:HA	1:A:762:LYS:HD3	1.85	0.42
1:A:200:ASN:HB3	1:A:203:SER:HB2	2.01	0.42
1:A:321:CYS:SG	1:A:334:LYS:HE3	2.60	0.41
1:A:431:LEU:O	1:A:435:VAL:HG23	2.20	0.41
1:A:134:ALA:O	1:A:145:GLN:HG3	2.21	0.41
1:A:702:GLN:HG2	7:A:2189:HOH:O	2.21	0.41
1:A:591:THR:O	1:A:592:GLN:HB2	2.20	0.41
1:A:728:ARG:HB3	2:B:115:LYS:O	2.21	0.41
1:A:559:ILE:HD11	1:A:573:ASP:HB3	2.04	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:494:GLU:CD	1:A:572:ARG:NH1[4_556]	1.58	0.62
1:A:494:GLU:OE2	1:A:572:ARG:NH1[4_556]	1.66	0.54
1:A:494:GLU:OE1	1:A:572:ARG:NH1[4_556]	1.67	0.53

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	712/788 (90%)	692 (97%)	20 (3%)	0	100	100
2	B	143/149 (96%)	138 (96%)	4 (3%)	1 (1%)	30	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	855/937 (91%)	830 (97%)	24 (3%)	1 (0%)	59	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	58	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	646/697 (93%)	620 (96%)	26 (4%)	42	56
2	B	125/129 (97%)	114 (91%)	11 (9%)	14	16
All	All	771/826 (93%)	734 (95%)	37 (5%)	35	46

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LYS
1	A	66	MET
1	A	162	LYS
1	A	176	GLN
1	A	177	ASP
1	A	229	LEU
1	A	266	SER
1	A	269	ASP
1	A	351	ASP
1	A	442	CYS
1	A	443	PHE
1	A	494	GLU
1	A	497	GLN
1	A	508	TYR
1	A	539	SER
1	A	559	ILE
1	A	614	ASP
1	A	651	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	667	ARG
1	A	670	LYS
1	A	726	LEU
1	A	773	ASP
1	A	777	LEU
1	A	779	GLU
1	A	792	ARG
1	A	811	LYS
2	B	29	THR
2	B	32	LEU
2	B	42	ASN
2	B	52	ILE
2	B	58	ASP
2	B	63	ILE
2	B	90	ARG
2	B	94	LYS
2	B	111	ASN
2	B	116	LEU
2	B	146	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	A	75	HIS
1	A	116	GLN
1	A	194	ASN
1	A	213	HIS
1	A	246	HIS
1	A	383	GLN
1	A	415	GLN
1	A	466	ASN
1	A	481	GLN
1	A	485	ASN
1	A	537	GLN
1	A	542	HIS
1	A	554	HIS
1	A	581	HIS
1	A	678	HIS
2	B	3	GLN
2	B	49	GLN
2	B	111	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ADP	A	998	5,4	29,29,29	1.17	4 (13%)	45,45,45	1.85	8 (17%)
4	BEF	A	999	3,5	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	998	5,4	-	0/16/32/32	0/1/3/3
4	BEF	A	999	3,5	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	998	ADP	C5-C4	3.07	1.47	1.40
3	A	998	ADP	C4-N9	-2.28	1.34	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	998	ADP	PA-O3A	2.16	1.63	1.59
3	A	998	ADP	C2-N3	2.02	1.36	1.32

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	998	ADP	N3-C2-N1	-6.94	122.90	128.71
3	A	998	ADP	N3-C4-N9	5.65	135.64	125.43
3	A	998	ADP	C5-C4-N3	-3.52	118.04	125.70
3	A	998	ADP	O4'-C1'-N9	3.10	111.33	108.44
3	A	998	ADP	C4-C5-N7	-2.99	106.97	109.52
3	A	998	ADP	C2-N3-C4	2.84	122.08	114.01
3	A	998	ADP	O2'-C2'-C1'	-2.16	104.69	111.23
3	A	998	ADP	C4'-O4'-C1'	2.09	112.02	109.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	724/788 (91%)	-0.04	18 (2%) 54 65	16, 31, 61, 75	14 (1%)
2	B	145/149 (97%)	-0.09	1 (0%) 84 91	25, 38, 56, 65	2 (1%)
All	All	869/937 (92%)	-0.05	19 (2%) 59 69	16, 33, 60, 75	16 (1%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	352	PHE	4.5
1	A	413	VAL	4.0
1	A	368	SER	2.9
1	A	391	THR	2.9
1	A	608	LEU	2.8
1	A	411	LEU	2.7
1	A	388	VAL	2.6
1	A	414	GLU	2.5
1	A	216	GLU	2.5
1	A	410	PRO	2.3
1	A	739	PHE	2.3
1	A	749	TYR	2.2
1	A	747	ILE	2.2
1	A	384	ASP	2.2
1	A	447	THR	2.2
1	A	611	GLU	2.1
2	B	75	LYS	2.1
1	A	812	TYR	2.1
1	A	508	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	BEF	A	999	4/4	0.17	0.76	22,22,24,25	0
3	ADP	A	998	27/27	0.15	0.31	17,20,22,26	0
6	CA	B	1150	1/1	0.11	-0.14	31,31,31,31	0
5	MG	A	1000	1/1	0.16	-0.33	24,24,24,24	0
6	CA	B	1148	1/1	0.05	-1.94	64,64,64,64	0
6	CA	B	1149	1/1	0.05	-2.40	72,72,72,72	0
6	CA	B	1151	1/1	0.05	-2.76	41,41,41,41	0

6.5 Other polymers ⓘ

There are no such residues in this entry.