



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 09:09 AM GMT

PDB ID : 2VDD
Title : CRYSTAL STRUCTURE OF THE OPEN STATE OF TOLC OUTER MEMBRANE COMPONENT OF MUTLIDRUG EFFLUX PUMPS
Authors : Bavro, V.N.; Pietras, Z.; Furnham, N.; Perez-Cano, L.; Fernandez-Recio, J.; Pei, X.Y.; Truer, R.; Misra, R.; Luisi, B.
Deposited on : 2007-10-04
Resolution : 3.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1341 (3.40-3.20)
Clashscore	79885	1696 (3.40-3.20)
Ramachandran outliers	78287	1664 (3.40-3.20)
Sidechain outliers	78261	1662 (3.40-3.20)
RSRZ outliers	66119	1342 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	460	
1	B	460	
1	C	460	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9915 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OUTER MEMBRANE PROTEIN TOLC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	12	0	1
			3303	2037	584	677	5			
1	B	429	Total	C	N	O	S	31	0	1
			3302	2036	584	677	5			
1	C	429	Total	C	N	O	S	5	0	1
			3303	2037	584	677	5			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ARG	LYS	CONFLICT	UNP P02930
A	191	LEU	VAL	ENGINEERED MUTATION	UNP P02930
A	384	PHE	TYR	ENGINEERED MUTATION	UNP P02930
A	389	GLU	ARG	ENGINEERED MUTATION	UNP P02930
B	2	ARG	LYS	CONFLICT	UNP P02930
B	191	LEU	VAL	ENGINEERED MUTATION	UNP P02930
B	384	PHE	TYR	ENGINEERED MUTATION	UNP P02930
B	389	GLU	ARG	ENGINEERED MUTATION	UNP P02930
C	2	ARG	LYS	CONFLICT	UNP P02930
C	191	LEU	VAL	ENGINEERED MUTATION	UNP P02930
C	384	PHE	TYR	ENGINEERED MUTATION	UNP P02930
C	389	GLU	ARG	ENGINEERED MUTATION	UNP P02930

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		
2	A	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	K	0	0
			2	2		
3	C	1	Total	K	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	O	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: OUTER MEMBRANE PROTEIN TOLC

Chain A:




• Molecule 1: OUTER MEMBRANE PROTEIN TOLC

Chain B:



• Molecule 1: OUTER MEMBRANE PROTEIN TOLC

Chain C:



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.48Å 70.97Å 219.95Å 90.00° 100.61° 90.00°	Depositor
Resolution (Å)	29.80 – 3.30 29.80 – 3.28	Depositor EDS
% Data completeness (in resolution range)	97.2 (29.80-3.30) 96.6 (29.80-3.28)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 3.24Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.236 , 0.295 0.208 , 0.214	Depositor DCC
R_{free} test set	1410 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , -5.6	EDS
Estimated twinning fraction	0.427 for $1/2^*h+3/2^*k, 1/2^*h-1/2^*k, -1/2^*h-1/2^*k-l$ 0.429 for $1/2^*h-3/2^*k, -1/2^*h-1/2^*k, -1/2^*h+1/2^*k-l$	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 28146 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9915	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: K, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/3344	0.56	0/4543
1	B	0.54	1/3343 (0.0%)	0.58	3/4541 (0.1%)
1	C	0.57	0/3344	0.58	2/4543 (0.0%)
All	All	0.56	1/10031 (0.0%)	0.57	5/13627 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	392	VAL	CA-CB	5.10	1.65	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	224	LYS	CB-CA-C	10.90	132.20	110.40
1	C	224	LYS	CA-CB-CG	6.97	128.74	113.40
1	B	393	ASP	CB-CG-OD2	5.17	122.95	118.30
1	B	392	VAL	CB-CA-C	-5.12	101.68	111.40
1	B	226	ASP	CA-CB-CG	-5.04	102.32	113.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	292	ALA	Peptide
1	B	79	ALA	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3303	0	3246	29	0
1	B	3302	0	3242	22	0
1	C	3303	0	3246	29	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	2	0	0	0	0
3	C	1	0	0	0	0
4	C	1	0	0	0	0
All	All	9915	0	9734	76	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

The worst 5 of 76 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:205:ARG:HH21	1:C:211:TYR:HB2	1.43	0.84
1:A:104:ARG:HG3	1:A:264:ASP:OD2	1.88	0.72
1:C:104:ARG:HG3	1:C:264:ASP:OD2	1.93	0.69
1:C:26:MET:O	1:C:30:GLN:HG2	1.94	0.67
1:B:204:LEU:O	1:B:208:THR:HG22	1.95	0.67

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/460 (93%)	397 (93%)	24 (6%)	6 (1%)	16	67
1	B	427/460 (93%)	414 (97%)	12 (3%)	1 (0%)	56	94
1	C	427/460 (93%)	395 (92%)	26 (6%)	6 (1%)	16	67
All	All	1281/1380 (93%)	1206 (94%)	62 (5%)	13 (1%)	22	74

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	78	ASP
1	C	289	ARG
1	C	386	VAL
1	C	442	SER
1	A	292	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/387 (92%)	340 (95%)	18 (5%)	34	78
1	B	358/387 (92%)	343 (96%)	15 (4%)	40	82
1	C	358/387 (92%)	343 (96%)	15 (4%)	40	82
All	All	1074/1161 (92%)	1026 (96%)	48 (4%)	38	81

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	96	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	151	GLN
1	C	303	GLN
1	B	100	MET
1	B	126	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	266	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	429/460 (93%)	-0.03	1 (0%)	93 70	20, 47, 85, 93	2 (0%)
1	B	428/460 (93%)	-0.04	2 (0%)	88 51	18, 48, 81, 90	7 (1%)
1	C	429/460 (93%)	-0.02	0	100 100	18, 48, 85, 97	1 (0%)
All	All	1286/1380 (93%)	-0.03	3 (0%)	93 70	18, 48, 84, 97	10 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	388	THR	2.8
1	A	379	ALA	2.4
1	B	293	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CL	A	1454	1/1	0.19	-0.46	51,51,51,51	0
3	K	A	1455	1/1	0.22	-0.61	66,66,66,66	0
2	CL	C	1454	1/1	0.18	-1.01	42,42,42,42	0
2	CL	B	1454	1/1	0.13	-1.86	31,31,31,31	0
3	K	A	1456	1/1	0.15	-2.45	68,68,68,68	0
3	K	C	1455	1/1	0.17	-52.00	79,79,79,79	0

6.5 Other polymers ⓘ

There are no such residues in this entry.