



wwPDB X-ray Structure Validation Summary Report i

Oct 3, 2014 – 03:33 AM EDT

PDB ID : 3VD1
Title : structure of p73 DNA binding domain tetramer modulates p73 transactivation
Authors : Ethayathulla, A.S.; Tse, P.W.; Nguyen, S.; Viadiu, H.
Deposited on : 2012-01-04
Resolution : 2.95 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.16 November 2013
Xtriage (Phenix)	:	dev-1439
EDS	:	stable23828
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable23828

1 Overall quality at a glance

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1587 (3.00-2.92)
Clashscore	79885	2029 (3.00-2.92)
Ramachandran outliers	78287	1955 (3.00-2.92)
Sidechain outliers	78261	1958 (3.00-2.92)
RSRZ outliers	66119	1588 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	210	
1	B	210	
1	C	210	
1	D	210	
1	I	210	
1	J	210	
1	K	210	
1	L	210	
2	E	12	
2	F	12	
2	G	12	
2	H	12	
2	M	12	
2	N	12	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	O	12	
2	P	12	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14716 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor protein p73.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	202	Total	C	N	O	S	0	0	0
			1581	991	284	295	11			
1	B	199	Total	C	N	O	S	0	0	0
			1560	976	281	292	11			
1	C	201	Total	C	N	O	S	0	0	0
			1575	985	285	294	11			
1	D	199	Total	C	N	O	S	0	0	0
			1560	976	281	292	11			
1	I	199	Total	C	N	O	S	0	0	0
			1560	976	281	292	11			
1	J	199	Total	C	N	O	S	0	0	0
			1560	976	281	292	11			
1	K	202	Total	C	N	O	S	0	0	0
			1590	996	286	297	11			
1	L	202	Total	C	N	O	S	0	0	0
			1590	996	286	297	11			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	103	MET	-	INITIATING METHIONINE	UNP O15350
A	104	GLY	-	EXPRESSION TAG	UNP O15350
A	105	HIS	-	EXPRESSION TAG	UNP O15350
A	106	HIS	-	EXPRESSION TAG	UNP O15350
A	107	HIS	-	EXPRESSION TAG	UNP O15350
A	108	HIS	-	EXPRESSION TAG	UNP O15350
A	109	HIS	-	EXPRESSION TAG	UNP O15350
A	110	HIS	-	EXPRESSION TAG	UNP O15350
A	111	HIS	-	EXPRESSION TAG	UNP O15350
A	112	HIS	-	EXPRESSION TAG	UNP O15350
A	113	GLU	-	EXPRESSION TAG	UNP O15350
A	114	PHE	-	EXPRESSION TAG	UNP O15350
B	103	MET	-	INITIATING METHIONINE	UNP O15350

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	104	GLY	-	EXPRESSION TAG	UNP O15350
B	105	HIS	-	EXPRESSION TAG	UNP O15350
B	106	HIS	-	EXPRESSION TAG	UNP O15350
B	107	HIS	-	EXPRESSION TAG	UNP O15350
B	108	HIS	-	EXPRESSION TAG	UNP O15350
B	109	HIS	-	EXPRESSION TAG	UNP O15350
B	110	HIS	-	EXPRESSION TAG	UNP O15350
B	111	HIS	-	EXPRESSION TAG	UNP O15350
B	112	HIS	-	EXPRESSION TAG	UNP O15350
B	113	GLU	-	EXPRESSION TAG	UNP O15350
B	114	PHE	-	EXPRESSION TAG	UNP O15350
C	103	MET	-	INITIATING METHIONINE	UNP O15350
C	104	GLY	-	EXPRESSION TAG	UNP O15350
C	105	HIS	-	EXPRESSION TAG	UNP O15350
C	106	HIS	-	EXPRESSION TAG	UNP O15350
C	107	HIS	-	EXPRESSION TAG	UNP O15350
C	108	HIS	-	EXPRESSION TAG	UNP O15350
C	109	HIS	-	EXPRESSION TAG	UNP O15350
C	110	HIS	-	EXPRESSION TAG	UNP O15350
C	111	HIS	-	EXPRESSION TAG	UNP O15350
C	112	HIS	-	EXPRESSION TAG	UNP O15350
C	113	GLU	-	EXPRESSION TAG	UNP O15350
C	114	PHE	-	EXPRESSION TAG	UNP O15350
D	103	MET	-	INITIATING METHIONINE	UNP O15350
D	104	GLY	-	EXPRESSION TAG	UNP O15350
D	105	HIS	-	EXPRESSION TAG	UNP O15350
D	106	HIS	-	EXPRESSION TAG	UNP O15350
D	107	HIS	-	EXPRESSION TAG	UNP O15350
D	108	HIS	-	EXPRESSION TAG	UNP O15350
D	109	HIS	-	EXPRESSION TAG	UNP O15350
D	110	HIS	-	EXPRESSION TAG	UNP O15350
D	111	HIS	-	EXPRESSION TAG	UNP O15350
D	112	HIS	-	EXPRESSION TAG	UNP O15350
D	113	GLU	-	EXPRESSION TAG	UNP O15350
D	114	PHE	-	EXPRESSION TAG	UNP O15350
I	103	MET	-	INITIATING METHIONINE	UNP O15350
I	104	GLY	-	EXPRESSION TAG	UNP O15350
I	105	HIS	-	EXPRESSION TAG	UNP O15350
I	106	HIS	-	EXPRESSION TAG	UNP O15350
I	107	HIS	-	EXPRESSION TAG	UNP O15350
I	108	HIS	-	EXPRESSION TAG	UNP O15350
I	109	HIS	-	EXPRESSION TAG	UNP O15350

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	110	HIS	-	EXPRESSION TAG	UNP O15350
I	111	HIS	-	EXPRESSION TAG	UNP O15350
I	112	HIS	-	EXPRESSION TAG	UNP O15350
I	113	GLU	-	EXPRESSION TAG	UNP O15350
I	114	PHE	-	EXPRESSION TAG	UNP O15350
J	103	MET	-	INITIATING METHIONINE	UNP O15350
J	104	GLY	-	EXPRESSION TAG	UNP O15350
J	105	HIS	-	EXPRESSION TAG	UNP O15350
J	106	HIS	-	EXPRESSION TAG	UNP O15350
J	107	HIS	-	EXPRESSION TAG	UNP O15350
J	108	HIS	-	EXPRESSION TAG	UNP O15350
J	109	HIS	-	EXPRESSION TAG	UNP O15350
J	110	HIS	-	EXPRESSION TAG	UNP O15350
J	111	HIS	-	EXPRESSION TAG	UNP O15350
J	112	HIS	-	EXPRESSION TAG	UNP O15350
J	113	GLU	-	EXPRESSION TAG	UNP O15350
J	114	PHE	-	EXPRESSION TAG	UNP O15350
K	103	MET	-	INITIATING METHIONINE	UNP O15350
K	104	GLY	-	EXPRESSION TAG	UNP O15350
K	105	HIS	-	EXPRESSION TAG	UNP O15350
K	106	HIS	-	EXPRESSION TAG	UNP O15350
K	107	HIS	-	EXPRESSION TAG	UNP O15350
K	108	HIS	-	EXPRESSION TAG	UNP O15350
K	109	HIS	-	EXPRESSION TAG	UNP O15350
K	110	HIS	-	EXPRESSION TAG	UNP O15350
K	111	HIS	-	EXPRESSION TAG	UNP O15350
K	112	HIS	-	EXPRESSION TAG	UNP O15350
K	113	GLU	-	EXPRESSION TAG	UNP O15350
K	114	PHE	-	EXPRESSION TAG	UNP O15350
L	103	MET	-	INITIATING METHIONINE	UNP O15350
L	104	GLY	-	EXPRESSION TAG	UNP O15350
L	105	HIS	-	EXPRESSION TAG	UNP O15350
L	106	HIS	-	EXPRESSION TAG	UNP O15350
L	107	HIS	-	EXPRESSION TAG	UNP O15350
L	108	HIS	-	EXPRESSION TAG	UNP O15350
L	109	HIS	-	EXPRESSION TAG	UNP O15350
L	110	HIS	-	EXPRESSION TAG	UNP O15350
L	111	HIS	-	EXPRESSION TAG	UNP O15350
L	112	HIS	-	EXPRESSION TAG	UNP O15350
L	113	GLU	-	EXPRESSION TAG	UNP O15350
L	114	PHE	-	EXPRESSION TAG	UNP O15350

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*

CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
2	F	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
2	G	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
2	H	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
2	M	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
2	N	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
2	O	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
2	P	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	J	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		
3	K	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	I	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		
3	L	1	Total	Zn	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	25	Total	O	0	0
			25	25		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	20	Total	O	0	0
			20	20		
4	C	28	Total	O	0	0
			28	28		
4	D	23	Total	O	0	0
			23	23		
4	I	26	Total	O	0	0
			26	26		
4	J	14	Total	O	0	0
			14	14		
4	K	20	Total	O	0	0
			20	20		
4	L	19	Total	O	0	0
			19	19		
4	E	3	Total	O	0	0
			3	3		
4	F	2	Total	O	0	0
			2	2		
4	H	1	Total	O	0	0
			1	1		
4	M	1	Total	O	0	0
			1	1		
4	N	3	Total	O	0	0
			3	3		
4	O	2	Total	O	0	0
			2	2		
4	P	1	Total	O	0	0
			1	1		




• Molecule 1: Tumor protein p73

Chain D:




• Molecule 1: Tumor protein p73

Chain I:




• Molecule 1: Tumor protein p73

Chain J:




• Molecule 1: Tumor protein p73

Chain K:




- Molecule 1: Tumor protein p73

Chain L:



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain E:



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain F:




- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain G:



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain H:




- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain M: 



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain N: 



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain O: 



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')

Chain P: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.54Å 104.20Å 123.22Å 90.00° 96.55° 90.00°	Depositor
Resolution (Å)	19.99 – 2.95 19.99 – 2.95	Depositor EDS
% Data completeness (in resolution range)	95.6 (19.99-2.95) 95.8 (19.99-2.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.93Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.239 , 0.285 0.248 , 0.294	Depositor DCC
R_{free} test set	843 reflections (2.02%)	DCC
Wilson B-factor (Å ²)	68.6	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 32.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 41779 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14716	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/1621	0.79	1/2204 (0.0%)
1	B	0.56	0/1599	0.81	3/2174 (0.1%)
1	C	0.50	0/1615	0.77	2/2196 (0.1%)
1	D	0.48	0/1599	0.75	0/2174
1	I	0.46	0/1599	0.72	0/2174
1	J	0.40	0/1599	0.73	1/2174 (0.0%)
1	K	0.52	0/1631	0.84	2/2217 (0.1%)
1	L	0.53	0/1631	0.85	1/2217 (0.0%)
2	E	0.54	0/272	0.84	0/418
2	F	0.77	0/272	1.29	2/418 (0.5%)
2	G	0.40	0/272	0.67	0/418
2	H	0.47	0/272	0.86	0/418
2	M	0.51	0/272	0.89	0/418
2	N	0.55	0/272	0.94	1/418 (0.2%)
2	O	0.51	0/272	0.88	0/418
2	P	0.45	0/272	0.84	0/418
All	All	0.50	0/15070	0.81	13/20874 (0.1%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	415	DA	P-O3'-C3'	10.24	131.98	119.70
1	K	284	VAL	N-CA-C	5.80	126.65	111.00
1	B	168	PRO	C-N-CD	-5.70	108.07	120.60
2	N	623	DG	C4'-C3'-O3'	5.65	123.83	109.70
1	A	125	HIS	N-CA-C	-5.65	95.75	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1581	0	1544	124	0
1	B	1560	0	1531	142	0
1	C	1575	0	1540	82	0
1	D	1560	0	1531	129	0
1	I	1560	0	1531	100	0
1	J	1560	0	1531	125	0
1	K	1590	0	1553	139	0
1	L	1590	0	1553	143	0
2	E	243	0	135	8	0
2	F	243	0	135	12	0
2	G	243	0	135	5	0
2	H	243	0	135	6	0
2	M	243	0	135	2	0
2	N	243	0	135	6	0
2	O	243	0	135	6	0
2	P	243	0	135	8	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
4	A	25	0	0	3	0
4	B	20	0	0	3	0
4	C	28	0	0	1	0
4	D	23	0	0	1	0
4	E	3	0	0	0	0
4	F	2	0	0	0	0
4	H	1	0	0	0	0
4	I	26	0	0	1	0
4	J	14	0	0	0	0
4	K	20	0	0	1	0
4	L	19	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	M	1	0	0	0	0
4	N	3	0	0	0	0
4	O	2	0	0	0	0
4	P	1	0	0	0	0
All	All	14716	0	13394	1003	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 36.

The worst 5 of 1003 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:169:PRO:CD	1:B:170:PRO:HD3	1.62	1.28
1:D:115:ILE:CB	1:D:116:PRO:HD3	1.66	1.22
1:B:169:PRO:HD2	1:B:170:PRO:CD	1.75	1.15
1:D:310:ARG:HH11	1:D:310:ARG:HA	1.15	1.11
1:C:138:LYS:HA	1:C:299:GLY:HA3	1.34	1.10

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/210 (95%)	173 (86%)	21 (10%)	6 (3%)	7	32
1	B	197/210 (94%)	165 (84%)	26 (13%)	6 (3%)	7	32
1	C	199/210 (95%)	170 (85%)	26 (13%)	3 (2%)	15	56
1	D	197/210 (94%)	153 (78%)	30 (15%)	14 (7%)	2	7
1	I	197/210 (94%)	166 (84%)	24 (12%)	7 (4%)	5	26
1	J	197/210 (94%)	161 (82%)	28 (14%)	8 (4%)	4	22
1	K	200/210 (95%)	158 (79%)	27 (14%)	15 (8%)	2	7
1	L	200/210 (95%)	167 (84%)	25 (12%)	8 (4%)	5	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1587/1680 (94%)	1313 (83%)	207 (13%)	67 (4%)	4	21

5 of 67 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	PHE
1	B	168	PRO
1	B	169	PRO
1	C	243	PRO
1	D	115	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/186 (95%)	142 (81%)	34 (19%)	2	10
1	B	175/186 (94%)	147 (84%)	28 (16%)	3	15
1	C	176/186 (95%)	146 (83%)	30 (17%)	3	13
1	D	175/186 (94%)	151 (86%)	24 (14%)	5	22
1	I	175/186 (94%)	156 (89%)	19 (11%)	9	33
1	J	175/186 (94%)	148 (85%)	27 (15%)	4	17
1	K	178/186 (96%)	150 (84%)	28 (16%)	4	16
1	L	178/186 (96%)	148 (83%)	30 (17%)	3	13
All	All	1408/1488 (95%)	1188 (84%)	220 (16%)	4	16

5 of 220 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	190	VAL
1	I	204	ASN
1	L	215	ILE
1	D	231	THR
1	D	287	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such

sidechains are listed below:

Mol	Chain	Res	Type
1	C	267	ASN
1	I	125	HIS
1	K	267	ASN
1	D	154	GLN
1	I	126	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	202/210 (96%)	0.17	5 (2%)	54	24	42, 74, 109, 114	2 (0%)
1	B	199/210 (94%)	0.04	4 (2%)	62	27	43, 73, 108, 129	2 (1%)
1	C	201/210 (95%)	-0.15	4 (1%)	62	27	41, 62, 99, 114	2 (0%)
1	D	199/210 (94%)	-0.02	6 (3%)	48	21	40, 69, 102, 120	1 (0%)
1	I	199/210 (94%)	-0.12	6 (3%)	48	21	42, 63, 102, 113	2 (1%)
1	J	199/210 (94%)	0.35	10 (5%)	28	14	52, 86, 112, 119	2 (1%)
1	K	202/210 (96%)	0.16	9 (4%)	32	15	42, 72, 100, 118	0
1	L	202/210 (96%)	-0.23	0	100	100	29, 53, 83, 96	1 (0%)
2	E	12/12 (100%)	0.80	1 (8%)	11	6	68, 79, 129, 155	0
2	F	12/12 (100%)	0.31	2 (16%)	2	2	57, 84, 126, 144	0
2	G	12/12 (100%)	0.07	0	100	100	66, 87, 99, 115	0
2	H	12/12 (100%)	-0.12	0	100	100	61, 85, 113, 121	0
2	M	12/12 (100%)	-0.10	0	100	100	51, 73, 100, 117	0
2	N	12/12 (100%)	-0.19	0	100	100	53, 76, 92, 95	0
2	O	12/12 (100%)	-0.19	0	100	100	64, 75, 108, 115	0
2	P	12/12 (100%)	-0.07	0	100	100	54, 70, 120, 128	0
All	All	1699/1776 (95%)	0.03	47 (2%)	50	22	29, 70, 107, 155	12 (0%)

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	409	DG	12.0
1	K	205	GLU	5.3
1	A	111	HIS	5.0
1	D	308	HIS	4.5
2	F	410	DC	4.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ZN	C	401	1/1	0.15	0.09	46,46,46,46	0
3	ZN	L	401	1/1	0.15	-0.44	57,57,57,57	0
3	ZN	I	401	1/1	0.12	-1.57	66,66,66,66	0
3	ZN	B	401	1/1	0.09	-1.57	46,46,46,46	0
3	ZN	A	401	1/1	0.12	-1.59	72,72,72,72	0
3	ZN	K	401	1/1	0.08	-1.78	59,59,59,59	0
3	ZN	J	401	1/1	0.10	-1.89	64,64,64,64	0
3	ZN	D	401	1/1	0.08	-2.14	53,53,53,53	0

6.5 Other polymers ⓘ

There are no such residues in this entry.