



wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 03:47 PM GMT

PDB ID : 1VHC
Title : Crystal structure of a putative KHG/KDPG aldolase
Authors : Structural GenomiX
Deposited on : 2003-12-01
Resolution : 1.89 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	224	
1	B	224	
1	C	224	
1	D	224	
1	E	224	
1	F	224	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 10319 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative KHG/KDPG aldolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	212	Total 1600	C 1029	N 268	O 295	S 2	Se 6	0	2	0
1	B	212	Total 1576	C 1013	N 262	O 293	S 2	Se 6	0	2	0
1	C	212	Total 1600	C 1029	N 268	O 295	S 2	Se 6	0	2	0
1	D	212	Total 1596	C 1028	N 268	O 292	S 2	Se 6	0	2	0
1	E	212	Total 1594	C 1025	N 268	O 293	S 2	Se 6	0	2	0
1	F	213	Total 1609	C 1035	N 270	O 296	S 2	Se 6	0	2	0

There are 102 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	cloning artifact	UNP P44480
A	0	SER	-	cloning artifact	UNP P44480
A	1	LEU	-	cloning artifact	UNP P44480
A	119	MSE	MET	modified residue	UNP P44480
A	127	MSE	MET	modified residue	UNP P44480
A	145	MSE	MET	modified residue	UNP P44480
A	159	MSE	MET	modified residue	UNP P44480
A	213	GLU	-	cloning artifact	UNP P44480
A	214	GLY	-	cloning artifact	UNP P44480
A	215	GLY	-	cloning artifact	UNP P44480
A	216	SER	-	cloning artifact	UNP P44480
A	217	HIS	-	cloning artifact	UNP P44480
A	218	HIS	-	cloning artifact	UNP P44480
A	219	HIS	-	cloning artifact	UNP P44480
A	220	HIS	-	cloning artifact	UNP P44480
A	221	HIS	-	cloning artifact	UNP P44480
A	222	HIS	-	cloning artifact	UNP P44480

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	MSE	-	cloning artifact	UNP P44480
B	0	SER	-	cloning artifact	UNP P44480
B	1	LEU	-	cloning artifact	UNP P44480
B	119	MSE	MET	modified residue	UNP P44480
B	127	MSE	MET	modified residue	UNP P44480
B	145	MSE	MET	modified residue	UNP P44480
B	159	MSE	MET	modified residue	UNP P44480
B	213	GLU	-	cloning artifact	UNP P44480
B	214	GLY	-	cloning artifact	UNP P44480
B	215	GLY	-	cloning artifact	UNP P44480
B	216	SER	-	cloning artifact	UNP P44480
B	217	HIS	-	cloning artifact	UNP P44480
B	218	HIS	-	cloning artifact	UNP P44480
B	219	HIS	-	cloning artifact	UNP P44480
B	220	HIS	-	cloning artifact	UNP P44480
B	221	HIS	-	cloning artifact	UNP P44480
B	222	HIS	-	cloning artifact	UNP P44480
C	-1	MSE	-	cloning artifact	UNP P44480
C	0	SER	-	cloning artifact	UNP P44480
C	1	LEU	-	cloning artifact	UNP P44480
C	119	MSE	MET	modified residue	UNP P44480
C	127	MSE	MET	modified residue	UNP P44480
C	145	MSE	MET	modified residue	UNP P44480
C	159	MSE	MET	modified residue	UNP P44480
C	213	GLU	-	cloning artifact	UNP P44480
C	214	GLY	-	cloning artifact	UNP P44480
C	215	GLY	-	cloning artifact	UNP P44480
C	216	SER	-	cloning artifact	UNP P44480
C	217	HIS	-	cloning artifact	UNP P44480
C	218	HIS	-	cloning artifact	UNP P44480
C	219	HIS	-	cloning artifact	UNP P44480
C	220	HIS	-	cloning artifact	UNP P44480
C	221	HIS	-	cloning artifact	UNP P44480
C	222	HIS	-	cloning artifact	UNP P44480
D	-1	MSE	-	cloning artifact	UNP P44480
D	0	SER	-	cloning artifact	UNP P44480
D	1	LEU	-	cloning artifact	UNP P44480
D	119	MSE	MET	modified residue	UNP P44480
D	127	MSE	MET	modified residue	UNP P44480
D	145	MSE	MET	modified residue	UNP P44480
D	159	MSE	MET	modified residue	UNP P44480
D	213	GLU	-	cloning artifact	UNP P44480

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	214	GLY	-	cloning artifact	UNP P44480
D	215	GLY	-	cloning artifact	UNP P44480
D	216	SER	-	cloning artifact	UNP P44480
D	217	HIS	-	cloning artifact	UNP P44480
D	218	HIS	-	cloning artifact	UNP P44480
D	219	HIS	-	cloning artifact	UNP P44480
D	220	HIS	-	cloning artifact	UNP P44480
D	221	HIS	-	cloning artifact	UNP P44480
D	222	HIS	-	cloning artifact	UNP P44480
E	-1	MSE	-	cloning artifact	UNP P44480
E	0	SER	-	cloning artifact	UNP P44480
E	1	LEU	-	cloning artifact	UNP P44480
E	119	MSE	MET	modified residue	UNP P44480
E	127	MSE	MET	modified residue	UNP P44480
E	145	MSE	MET	modified residue	UNP P44480
E	159	MSE	MET	modified residue	UNP P44480
E	213	GLU	-	cloning artifact	UNP P44480
E	214	GLY	-	cloning artifact	UNP P44480
E	215	GLY	-	cloning artifact	UNP P44480
E	216	SER	-	cloning artifact	UNP P44480
E	217	HIS	-	cloning artifact	UNP P44480
E	218	HIS	-	cloning artifact	UNP P44480
E	219	HIS	-	cloning artifact	UNP P44480
E	220	HIS	-	cloning artifact	UNP P44480
E	221	HIS	-	cloning artifact	UNP P44480
E	222	HIS	-	cloning artifact	UNP P44480
F	-1	MSE	-	cloning artifact	UNP P44480
F	0	SER	-	cloning artifact	UNP P44480
F	1	LEU	-	cloning artifact	UNP P44480
F	119	MSE	MET	modified residue	UNP P44480
F	127	MSE	MET	modified residue	UNP P44480
F	145	MSE	MET	modified residue	UNP P44480
F	159	MSE	MET	modified residue	UNP P44480
F	213	GLU	-	cloning artifact	UNP P44480
F	214	GLY	-	cloning artifact	UNP P44480
F	215	GLY	-	cloning artifact	UNP P44480
F	216	SER	-	cloning artifact	UNP P44480
F	217	HIS	-	cloning artifact	UNP P44480
F	218	HIS	-	cloning artifact	UNP P44480
F	219	HIS	-	cloning artifact	UNP P44480
F	220	HIS	-	cloning artifact	UNP P44480
F	221	HIS	-	cloning artifact	UNP P44480

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	222	HIS	-	cloning artifact	UNP P44480

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	121	Total O 121 121	0	0
2	B	98	Total O 98 98	0	0
2	C	113	Total O 113 113	0	0
2	D	122	Total O 122 122	0	0
2	E	94	Total O 94 94	0	0
2	F	196	Total O 196 196	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative KHG/KDPG aldolase

Chain A:




- Molecule 1: Putative KHG/KDPG aldolase

Chain B:




- Molecule 1: Putative KHG/KDPG aldolase

Chain C:



- Molecule 1: Putative KHG/KDPG aldolase

Chain D:



- Molecule 1: Putative KHG/KDPG aldolase

Chain E: 

- Molecule 1: Putative KHG/KDPG aldolase

Chain F: 

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.27Å 133.23Å 89.73Å 90.00° 105.82° 90.00°	Depositor
Resolution (Å)	30.40 – 1.89 30.40 – 1.89	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.40-1.89) 91.2 (30.40-1.89)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 4	Depositor
R, R_{free}	0.219 , 0.256 0.203 , 0.236	Depositor DCC
R_{free} test set	6406 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	32.8	Xtrriage
Anisotropy	0.664	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.0	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	0 of 127796 reflections	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10319	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	0/1630	1.13	4/2212 (0.2%)
1	B	0.76	0/1606	1.17	7/2185 (0.3%)
1	C	0.81	0/1630	1.22	9/2212 (0.4%)
1	D	0.87	0/1626	1.29	10/2206 (0.5%)
1	E	0.71	0/1624	1.08	2/2204 (0.1%)
1	F	0.93	0/1639	1.31	9/2223 (0.4%)
All	All	0.81	0/9755	1.20	41/13242 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	64	ARG	NE-CZ-NH1	14.88	127.74	120.30
1	D	170	ARG	NE-CZ-NH1	12.40	126.50	120.30
1	D	58	ARG	CD-NE-CZ	10.88	138.83	123.60
1	F	49	ARG	CD-NE-CZ	10.45	138.22	123.60
1	D	64	ARG	NE-CZ-NH2	-10.44	115.08	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	168	ASN	Mainchain

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1600	0	1653	20	0
1	B	1576	0	1589	29	0
1	C	1600	0	1653	24	0
1	D	1596	0	1654	28	0
1	E	1594	0	1642	26	0
1	F	1609	0	1669	20	0
2	A	121	0	0	3	0
2	B	98	0	0	4	0
2	C	113	0	0	0	0
2	D	122	0	0	3	0
2	E	94	0	0	1	0
2	F	196	0	0	5	0
All	All	10319	0	9860	112	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

The worst 5 of 112 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:119[B]:MSE:HE1	1:B:119[B]:MSE:HG2	1.20	1.08
1:E:119[B]:MSE:SE	1:F:119[B]:MSE:SE	2.79	0.99
1:A:119[B]:MSE:SE	1:B:119[B]:MSE:SE	2.91	0.87
1:D:119[B]:MSE:SE	1:E:119[B]:MSE:HE2	2.28	0.83
1:D:144:LYS:HE2	1:E:139:ALA:HA	1.61	0.82

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/224 (95%)	205 (97%)	7 (3%)	0	100	100
1	B	212/224 (95%)	204 (96%)	8 (4%)	0	100	100
1	C	212/224 (95%)	207 (98%)	5 (2%)	0	100	100
1	D	212/224 (95%)	205 (97%)	7 (3%)	0	100	100
1	E	212/224 (95%)	205 (97%)	7 (3%)	0	100	100
1	F	213/224 (95%)	206 (97%)	7 (3%)	0	100	100
All	All	1273/1344 (95%)	1232 (97%)	41 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains i

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/178 (96%)	171 (100%)	0	100	100
1	B	163/178 (92%)	161 (99%)	2 (1%)	82	80
1	C	171/178 (96%)	170 (99%)	1 (1%)	92	92
1	D	170/178 (96%)	165 (97%)	5 (3%)	55	44
1	E	169/178 (95%)	165 (98%)	4 (2%)	61	53
1	F	172/178 (97%)	171 (99%)	1 (1%)	92	92
All	All	1016/1068 (95%)	1003 (99%)	13 (1%)	80	77

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	144	LYS
1	D	166	LEU
1	E	63	ASN
1	D	14	GLU
1	E	23	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	195	ASN
1	D	195	ASN
1	C	155	GLN
1	B	7	GLN
1	D	155	GLN

5.3.3 RNA [i](#)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/224 (94%)	-0.14	5 (2%) 56 57	23, 32, 46, 60	0
1	B	212/224 (94%)	0.27	14 (6%) 18 17	23, 36, 58, 71	0
1	C	212/224 (94%)	-0.08	4 (1%) 64 65	22, 32, 49, 60	0
1	D	212/224 (94%)	-0.12	3 (1%) 72 74	21, 31, 55, 67	0
1	E	212/224 (94%)	0.09	9 (4%) 35 35	23, 37, 56, 63	0
1	F	213/224 (95%)	-0.25	1 (0%) 88 90	20, 27, 42, 59	0
All	All	1273/1344 (94%)	-0.04	36 (2%) 50 51	20, 32, 54, 71	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	2	SER	5.0
1	E	3	TYR	4.8
1	B	3	TYR	4.6
1	D	190	LYS	4.5
1	B	51	GLU	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.