



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 01:38 PM GMT

PDB ID : 1VHG  
Title : Crystal structure of ADP compounds hydrolase  
Authors : Structural GenomiX  
Deposited on : 2003-12-01  
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1557 (2.70-2.70)
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)
RSRZ outliers	66119	1559 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	198	
1	B	198	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3149 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP compounds hydrolase nudE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	Se	0	1	0
			1480	941	250	282	7			
1	B	185	Total	C	N	O	Se	0	0	0
			1464	932	248	278	6			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	cloning artifact	UNP P45799
A	0	SER	-	cloning artifact	UNP P45799
A	1	LEU	-	cloning artifact	UNP P45799
A	42	MSE	MET	modified residue	UNP P45799
A	51	MSE	MET	modified residue	UNP P45799
A	115	MSE	MET	modified residue	UNP P45799
A	124	MSE	MET	modified residue	UNP P45799
A	155	MSE	MET	modified residue	UNP P45799
A	156	MSE	MET	modified residue	UNP P45799
A	187	GLU	-	cloning artifact	UNP P45799
A	188	GLY	-	cloning artifact	UNP P45799
A	189	GLY	-	cloning artifact	UNP P45799
A	190	SER	-	cloning artifact	UNP P45799
A	191	HIS	-	cloning artifact	UNP P45799
A	192	HIS	-	cloning artifact	UNP P45799
A	193	HIS	-	cloning artifact	UNP P45799
A	194	HIS	-	cloning artifact	UNP P45799
A	195	HIS	-	cloning artifact	UNP P45799
A	196	HIS	-	cloning artifact	UNP P45799
B	-1	MSE	-	cloning artifact	UNP P45799
B	0	SER	-	cloning artifact	UNP P45799
B	1	LEU	-	cloning artifact	UNP P45799
B	42	MSE	MET	modified residue	UNP P45799
B	51	MSE	MET	modified residue	UNP P45799
B	115	MSE	MET	modified residue	UNP P45799

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	124	MSE	MET	modified residue	UNP P45799
B	155	MSE	MET	modified residue	UNP P45799
B	156	MSE	MET	modified residue	UNP P45799
B	187	GLU	-	cloning artifact	UNP P45799
B	188	GLY	-	cloning artifact	UNP P45799
B	189	GLY	-	cloning artifact	UNP P45799
B	190	SER	-	cloning artifact	UNP P45799
B	191	HIS	-	cloning artifact	UNP P45799
B	192	HIS	-	cloning artifact	UNP P45799
B	193	HIS	-	cloning artifact	UNP P45799
B	194	HIS	-	cloning artifact	UNP P45799
B	195	HIS	-	cloning artifact	UNP P45799
B	196	HIS	-	cloning artifact	UNP P45799

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	97	Total O 97 97	0	0
2	B	108	Total O 108 108	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP compounds hydrolase nudE

Chain A: 



- Molecule 1: ADP compounds hydrolase nudE

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	193.17Å 193.17Å 193.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.17 – 2.70 41.18 – 2.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (41.17-2.70) 100.0 (41.18-2.70)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.59 (at 2.69Å)	Xtriage
Refinement program	REFMAC 4	Depositor
R, $R_{free}$	0.211 , 0.272 0.192 , 0.240	Depositor DCC
$R_{free}$ test set	868 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.2	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 17164 reflections (0.006%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3149	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/1504	1.18	8/2028 (0.4%)
1	B	0.57	0/1488	1.26	14/2009 (0.7%)
All	All	0.58	0/2992	1.22	22/4037 (0.5%)

There are no bond length outliers.

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	37	ARG	NE-CZ-NH2	-10.69	114.95	120.30
1	B	37	ARG	NE-CZ-NH1	10.12	125.36	120.30
1	A	185	ARG	NE-CZ-NH2	8.82	124.71	120.30
1	B	37	ARG	CD-NE-CZ	7.70	134.38	123.60
1	B	94	ARG	NE-CZ-NH2	-7.63	116.49	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1480	0	1465	49	1
1	B	1464	0	1441	38	1
2	A	97	0	0	6	0
2	B	108	0	0	4	0
All	All	3149	0	2906	78	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:115[A]:MSE:HE3	1:A:168:ARG:HD2	1.56	0.87
1:A:51:MSE:HE3	1:A:79:LYS:HE3	1.56	0.86
1:A:17:ALA:HB3	1:A:24:VAL:HG13	1.60	0.81
1:A:11:LEU:HB2	1:A:28:ASP:HB3	1.69	0.75
1:A:185:ARG:HH21	1:A:185:ARG:HB2	1.52	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:33:ASN:O	1:B:37:ARG:NH1[15_554]	2.10	0.10

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/198 (93%)	173 (94%)	11 (6%)	0	100	100
1	B	183/198 (92%)	176 (96%)	6 (3%)	1 (0%)	38	70
All	All	367/396 (93%)	349 (95%)	17 (5%)	1 (0%)	50	82

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	165	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/168 (97%)	139 (85%)	24 (15%)	4	11
1	B	159/168 (95%)	140 (88%)	19 (12%)	8	18
All	All	322/336 (96%)	279 (87%)	43 (13%)	6	14

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	137	LEU
1	B	2	SER
1	B	159	LEU
1	A	138	GLU
1	A	159	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	GLN
1	B	183	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	185/198 (93%)	-0.30	2 (1%) 77 82	22, 35, 66, 85	0
1	B	185/198 (93%)	-0.25	5 (2%) 52 57	20, 37, 75, 95	0
All	All	370/396 (93%)	-0.28	7 (1%) 64 70	20, 36, 72, 95	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	SER	4.7
1	B	119	TYR	3.9
1	B	17	ALA	2.7
1	A	22	PHE	2.4
1	B	185	ARG	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

### 6.5 Other polymers ⓘ

There are no such residues in this entry.