



# wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 12:08 AM GMT

PDB ID : 2VHW  
Title : CRYSTAL STRUCTURE OF HOLO L-ALANINE DEHYDROGENASE FROM MYCOBACTERIUM TUBERCULOSIS IN THE OPEN AND CLOSED CONFORMATION  
Authors : Agren, D.; Schneider, G.  
Deposited on : 2007-11-26  
Resolution : 2.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	377	
1	B	377	
1	C	377	
1	D	377	
1	E	377	
1	F	377	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 18159 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ALANINE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	371	2719	1710	479	517	13	0	0	0
1	B	370	2730	1717	479	521	13	0	3	0
1	C	371	2719	1710	479	517	13	0	0	0
1	D	370	2714	1707	478	516	13	0	0	0
1	E	373	2747	1727	488	519	13	0	1	0
1	F	373	2752	1730	488	521	13	0	2	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 1,4-DIHYDRONICOTINAMIDEADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: C<sub>21</sub>H<sub>29</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	44	21	7	14	2	0	0
3	B	1	44	21	7	14	2	0	0
3	C	1	44	21	7	14	2	0	0
3	D	1	44	21	7	14	2	0	0
3	E	1	44	21	7	14	2	0	0
3	F	1	44	21	7	14	2	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	219	219	219	0	0
4	B	310	310	310	0	0
4	C	211	211	211	0	0
4	D	297	297	297	0	0
4	E	195	195	195	0	0
4	F	280	280	280	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ALANINE DEHYDROGENASE

Chain A:




- Molecule 1: ALANINE DEHYDROGENASE

Chain B:




- Molecule 1: ALANINE DEHYDROGENASE

Chain C:




- Molecule 1: ALANINE DEHYDROGENASE

Chain D:




- Molecule 1: ALANINE DEHYDROGENASE

Chain E:



- Molecule 1: ALANINE DEHYDROGENASE

Chain F:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	177.03Å 172.01Å 98.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.00 49.56 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (40.00-2.00) 94.6 (49.56-2.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.176 , 0.212 0.219 , 0.247	Depositor DCC
$R_{free}$ test set	9700 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.0	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 41.7	EDS
Estimated twinning fraction	0.014 for k,h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 191725 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	18159	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NAI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.56	1/2769 (0.0%)	0.72	3/3773 (0.1%)
1	B	0.60	0/2789	0.76	4/3800 (0.1%)
1	C	0.53	0/2769	0.66	2/3773 (0.1%)
1	D	0.58	0/2764	0.67	1/3766 (0.0%)
1	E	0.52	0/2802	0.67	3/3817 (0.1%)
1	F	0.56	0/2810	0.68	1/3828 (0.0%)
All	All	0.56	1/16703 (0.0%)	0.69	14/22757 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	46	THR	CB-OG1	-5.76	1.31	1.43

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	272	GLY	N-CA-C	-10.27	87.42	113.10
1	B	272	GLY	N-CA-C	-8.68	91.39	113.10
1	E	205	ARG	NE-CZ-NH2	-8.31	116.14	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	272	GLY	N-CA-C	-7.83	93.51	113.10
1	E	205	ARG	NE-CZ-NH1	7.65	124.13	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	271	GLN	Peptide
1	B	271	GLN	Peptide
1	C	271	GLN	Peptide

## 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2719	0	2732	21	0
1	B	2730	0	2743	14	0
1	C	2719	0	2732	32	0
1	D	2714	0	2727	23	0
1	E	2747	0	2759	19	0
1	F	2752	0	2763	20	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	44	0	27	1	0
3	B	44	0	27	1	0
3	C	44	0	27	3	0
3	D	44	0	27	3	0
3	E	44	0	27	2	0
3	F	44	0	27	1	0
4	A	219	0	0	0	0
4	B	310	0	0	1	0
4	C	211	0	0	1	0
4	D	297	0	0	3	0
4	E	195	0	0	3	0
4	F	280	0	0	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	18159	0	16618	120	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

The worst 5 of 120 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:101:ARG:HG3	1:C:101:ARG:HH21	1.18	1.04
1:C:241:VAL:HG23	1:C:244:ALA:HB3	1.57	0.87
1:C:101:ARG:CG	1:C:101:ARG:HH21	1.87	0.86
1:D:282:THR:HG22	1:D:284:ASP:H	1.43	0.83
1:B:255:VAL:HG11	1:B:290:VAL:HG22	1.63	0.78

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/377 (98%)	362 (98%)	6 (2%)	1 (0%)	50	44
1	B	371/377 (98%)	364 (98%)	6 (2%)	1 (0%)	50	44
1	C	369/377 (98%)	361 (98%)	8 (2%)	0	100	100
1	D	368/377 (98%)	364 (99%)	4 (1%)	0	100	100
1	E	372/377 (99%)	366 (98%)	6 (2%)	0	100	100
1	F	373/377 (99%)	368 (99%)	5 (1%)	0	100	100
All	All	2222/2262 (98%)	2185 (98%)	35 (2%)	2 (0%)	59	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	272	GLY
1	B	272	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/282 (98%)	265 (96%)	11 (4%)	42	36
1	B	279/282 (99%)	273 (98%)	6 (2%)	64	65
1	C	276/282 (98%)	264 (96%)	12 (4%)	40	33
1	D	276/282 (98%)	270 (98%)	6 (2%)	64	65
1	E	279/282 (99%)	273 (98%)	6 (2%)	64	65
1	F	280/282 (99%)	274 (98%)	6 (2%)	66	67
All	All	1666/1692 (98%)	1619 (97%)	47 (3%)	56	54

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	130	LEU
1	C	339	LEU
1	F	71	LEU
1	C	290	VAL
1	C	361	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	121	GLN
1	C	148	HIS
1	E	148	HIS
1	C	31	HIS
1	E	56	GLN

### 5.3.3 RNA [i](#)

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAI	A	1373	-	48,48,48	1.45	6 (12%)	73,73,73	1.92	14 (19%)
3	NAI	B	1372	-	48,48,48	1.44	6 (12%)	73,73,73	1.98	16 (21%)
3	NAI	C	1372	-	48,48,48	1.40	6 (12%)	73,73,73	1.93	12 (16%)
3	NAI	D	1371	-	48,48,48	1.46	6 (12%)	73,73,73	2.05	13 (17%)
3	NAI	E	1374	-	48,48,48	1.41	6 (12%)	73,73,73	1.72	12 (16%)
3	NAI	F	1374	-	48,48,48	1.40	6 (12%)	73,73,73	1.71	14 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAI	A	1373	-	-	0/30/72/72	0/3/5/5
3	NAI	B	1372	-	-	0/30/72/72	0/3/5/5
3	NAI	C	1372	-	-	0/30/72/72	0/3/5/5
3	NAI	D	1371	-	-	0/30/72/72	0/3/5/5
3	NAI	E	1374	-	-	0/30/72/72	0/3/5/5
3	NAI	F	1374	-	-	0/30/72/72	0/3/5/5

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1373	NAI	C4N-C3N	-5.75	1.39	1.50
3	B	1372	NAI	C4N-C3N	-5.56	1.39	1.50
3	C	1372	NAI	C4N-C3N	-5.20	1.40	1.50
3	D	1371	NAI	C4N-C3N	-5.16	1.40	1.50
3	E	1374	NAI	C4N-C3N	-5.04	1.40	1.50

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1371	NAI	O4B-C1B-N9A	8.64	116.47	108.44
3	C	1372	NAI	O4B-C1B-N9A	8.54	116.39	108.44
3	B	1372	NAI	O4B-C1B-N9A	7.98	115.86	108.44
3	A	1373	NAI	N3A-C2A-N1A	-7.39	122.53	128.71
3	E	1374	NAI	N3A-C2A-N1A	-7.03	122.83	128.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	371/377 (98%)	0.51	16 (4%) 34 32	26, 46, 56, 68	0
1	B	370/377 (98%)	0.16	4 (1%) 77 78	25, 40, 51, 58	0
1	C	371/377 (98%)	0.57	29 (7%) 13 12	26, 47, 61, 70	0
1	D	370/377 (98%)	0.14	5 (1%) 72 72	25, 40, 52, 57	0
1	E	373/377 (98%)	0.76	34 (9%) 9 9	27, 50, 65, 76	0
1	F	373/377 (98%)	0.45	16 (4%) 34 32	26, 42, 49, 66	0
All	All	2228/2262 (98%)	0.43	104 (4%) 30 29	25, 44, 57, 76	0

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	283	TYR	5.8
1	E	372	HIS	4.8
1	C	371	ALA	4.2
1	A	57	LEU	4.2
1	F	34	LEU	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAI	C	1372	44/44	0.13	-0.20	36,41,46,47	0
3	NAI	D	1371	44/44	0.10	-0.84	30,34,39,43	0
3	NAI	A	1373	44/44	0.09	-1.51	30,34,39,43	0
3	NAI	E	1374	44/44	0.10	-1.52	27,31,40,41	0
3	NAI	F	1374	44/44	0.10	-1.76	20,24,29,31	0
3	NAI	B	1372	44/44	0.07	-2.10	23,29,33,36	0
2	MG	B	1371	1/1	0.03	-2.24	45,45,45,45	0
2	MG	A	1372	1/1	0.11	-2.58	48,48,48,48	0

## 6.5 Other polymers

There are no such residues in this entry.