



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 02:03 AM GMT

PDB ID : 1VKF
Title : CRYSTAL STRUCTURE OF A GLYCEROL UPTAKE OPERON ANTITERMINATOR-RELATED PROTEIN (TM1436) FROM THERMOTOGA MARITIMA MSB8 AT 1.65 Å RESOLUTION
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2004-05-13
Resolution : 1.65 Å (reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1404 (1.68-1.64)
Clashscore	79885	1001 (1.66-1.66)
Ramachandran outliers	78287	1581 (1.68-1.64)
Sidechain outliers	78261	1580 (1.68-1.64)
RSRZ outliers	66119	1404 (1.68-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	188	
1	B	188	
1	C	188	
1	D	188	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	CIT	C	502	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5716 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glycerol uptake operon antiterminator-related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	172	Total	C	N	O	Se	0	2	0
			1334	870	221	239	4			
1	B	173	Total	C	N	O	Se	0	1	0
			1338	870	223	241	4			
1	C	173	Total	C	N	O	Se	0	0	0
			1311	853	216	238	4			
1	D	172	Total	C	N	O	Se	0	1	0
			1298	844	212	238	4			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	LEADER SEQUENCE	UNP Q9X1F0
A	-10	GLY	-	LEADER SEQUENCE	UNP Q9X1F0
A	-9	SER	-	LEADER SEQUENCE	UNP Q9X1F0
A	-8	ASP	-	LEADER SEQUENCE	UNP Q9X1F0
A	-7	LYS	-	LEADER SEQUENCE	UNP Q9X1F0
A	-6	ILE	-	LEADER SEQUENCE	UNP Q9X1F0
A	-5	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
A	-4	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
A	-3	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
A	-2	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
A	-1	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
A	0	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
A	12	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
A	51	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
A	172	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
B	-11	MET	-	LEADER SEQUENCE	UNP Q9X1F0
B	-10	GLY	-	LEADER SEQUENCE	UNP Q9X1F0
B	-9	SER	-	LEADER SEQUENCE	UNP Q9X1F0
B	-8	ASP	-	LEADER SEQUENCE	UNP Q9X1F0
B	-7	LYS	-	LEADER SEQUENCE	UNP Q9X1F0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	ILE	-	LEADER SEQUENCE	UNP Q9X1F0
B	-5	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
B	-4	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
B	-3	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
B	-2	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
B	-1	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
B	0	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
B	12	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
B	51	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
B	172	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
C	-11	MET	-	LEADER SEQUENCE	UNP Q9X1F0
C	-10	GLY	-	LEADER SEQUENCE	UNP Q9X1F0
C	-9	SER	-	LEADER SEQUENCE	UNP Q9X1F0
C	-8	ASP	-	LEADER SEQUENCE	UNP Q9X1F0
C	-7	LYS	-	LEADER SEQUENCE	UNP Q9X1F0
C	-6	ILE	-	LEADER SEQUENCE	UNP Q9X1F0
C	-5	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
C	-4	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
C	-3	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
C	-2	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
C	-1	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
C	0	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
C	12	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
C	51	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
C	172	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
D	-11	MET	-	LEADER SEQUENCE	UNP Q9X1F0
D	-10	GLY	-	LEADER SEQUENCE	UNP Q9X1F0
D	-9	SER	-	LEADER SEQUENCE	UNP Q9X1F0
D	-8	ASP	-	LEADER SEQUENCE	UNP Q9X1F0
D	-7	LYS	-	LEADER SEQUENCE	UNP Q9X1F0
D	-6	ILE	-	LEADER SEQUENCE	UNP Q9X1F0
D	-5	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
D	-4	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
D	-3	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
D	-2	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
D	-1	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
D	0	HIS	-	LEADER SEQUENCE	UNP Q9X1F0
D	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
D	12	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0
D	51	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	172	MSE	MET	MODIFIED RESIDUE	UNP Q9X1F0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	C	1	Total	C	O	0	0
			13	6	7		
3	D	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	148	Total	O	0	0
			148	148		
4	B	119	Total	O	0	0
			119	119		
4	C	74	Total	O	0	0
			74	74		
4	D	50	Total	O	0	0
			50	50		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: glycerol uptake operon antiterminator-relatedprotein

Chain A: 



- Molecule 1: glycerol uptake operon antiterminator-relatedprotein

Chain B: 




- Molecule 1: glycerol uptake operon antiterminator-relatedprotein

Chain C: 



- Molecule 1: glycerol uptake operon antiterminator-relatedprotein

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	84.53Å 138.78Å 160.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.19 – 1.65 72.19 – 1.65	Depositor EDS
% Data completeness (in resolution range)	97.1 (72.19-1.65) 97.1 (72.19-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.185 , 0.211 0.197 , 0.222	Depositor DCC
R_{free} test set	5519 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.655	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.7	EDS
Estimated twinning fraction	0.017 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.023 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 109952 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5716	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/1352	0.86	4/1820 (0.2%)
1	B	0.72	1/1355 (0.1%)	0.91	4/1822 (0.2%)
1	C	0.62	0/1328	0.83	4/1790 (0.2%)
1	D	1.39	12/1315 (0.9%)	1.01	4/1774 (0.2%)
All	All	0.92	13/5350 (0.2%)	0.90	16/7206 (0.2%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	126	ALA	C-O	24.62	1.70	1.23
1	D	128	ALA	C-O	12.04	1.46	1.23
1	D	99	ALA	C-O	9.58	1.41	1.23
1	D	153	ARG	C-O	8.49	1.39	1.23
1	D	102	SER	C-O	8.41	1.39	1.23
1	D	169	LEU	C-O	7.36	1.37	1.23
1	B	172	MSE	SE-CE	-7.24	1.52	1.95
1	D	155	ILE	C-O	6.48	1.35	1.23
1	D	168	ILE	C-N	5.69	1.47	1.34
1	D	38	LYS	C-O	-5.63	1.12	1.23
1	D	152	ALA	C-O	5.38	1.33	1.23
1	D	124	PRO	CA-CB	5.23	1.64	1.53
1	D	144	GLY	N-CA	5.03	1.53	1.46

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	96	ARG	NE-CZ-NH2	-11.36	114.62	120.30
1	B	96	ARG	NE-CZ-NH1	11.33	125.97	120.30
1	A	11	ASP	CB-CG-OD2	7.32	124.89	118.30
1	C	101	ASP	CB-CG-OD2	6.53	124.17	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	73	ASP	CB-CG-OD2	6.02	123.72	118.30
1	D	73	ASP	CB-CG-OD2	5.92	123.62	118.30
1	B	118	ASP	CB-CG-OD2	5.76	123.48	118.30
1	C	118	ASP	CB-CG-OD2	5.71	123.44	118.30
1	D	118	ASP	CB-CG-OD2	5.66	123.40	118.30
1	A	13	ASP	CB-CG-OD2	5.62	123.36	118.30
1	D	24	PHE	CB-CG-CD2	-5.60	116.88	120.80
1	C	96	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	A	50	ASP	CB-CG-OD2	5.47	123.22	118.30
1	A	42	ASP	CB-CG-OD2	5.41	123.17	118.30
1	D	128	ALA	N-CA-C	5.38	125.51	111.00
1	B	101	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1334	0	1405	4	0
1	B	1338	0	1421	8	0
1	C	1311	0	1369	6	0
1	D	1298	0	1338	16	0
2	B	5	0	0	0	0
3	A	13	0	5	0	0
3	C	13	0	5	2	0
3	D	13	0	5	0	0
4	A	148	0	0	3	0
4	B	119	0	0	1	0
4	C	74	0	0	0	0
4	D	50	0	0	0	0
All	All	5716	0	5548	33	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 3.

All (33) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:126:ALA:O	1:D:126:ALA:C	1.70	1.28
1:B:172:MSE:HG2	1:B:173:LYS:HA	1.76	0.67
3:C:502:CIT:C6	3:C:502:CIT:O2	2.43	0.65
1:D:131:VAL:O	1:D:135:ILE:HG12	1.96	0.65
1:D:128:ALA:O	1:D:140:VAL:HG11	2.03	0.58
1:B:169:LEU:HD22	1:B:172:MSE:HE2	1.87	0.56
1:A:96:ARG:CZ	4:A:536:HOH:O	2.55	0.55
1:D:101:ASP:O	1:D:105:VAL:HG23	2.07	0.54
1:B:69:LYS:HD2	1:C:16:GLY:HA2	1.90	0.53
1:C:153:ARG:CD	1:D:153:ARG:HD3	2.41	0.51
1:D:125:GLY:HA2	1:D:142:ALA:HB1	1.95	0.47
1:D:3:LYS:HA	1:D:160:SER:O	2.14	0.47
1:C:12:MSE:HA	1:C:15:ILE:HD11	1.97	0.47
1:D:155:ILE:O	1:D:159:VAL:HG22	2.15	0.46
1:B:43:ARG:NH1	4:B:493:HOH:O	2.50	0.45
1:D:5:ILE:HG23	1:D:164:THR:HG22	1.98	0.44
1:A:96:ARG:NE	4:A:536:HOH:O	2.51	0.44
3:C:502:CIT:O4	3:C:502:CIT:O7	2.36	0.44
1:D:19:GLU:N	1:D:20:PRO:CD	2.81	0.44
1:C:172:MSE:O	1:C:173:LYS:C	2.56	0.44
1:C:12:MSE:HG3	1:C:36:HIS:CD2	2.54	0.43
1:D:112:ILE:HD11	1:D:120:VAL:HG11	2.00	0.43
1:B:172:MSE:HB3	1:B:172:MSE:HE3	1.80	0.43
1:D:168:ILE:H	1:D:168:ILE:HD12	1.84	0.42
1:D:51:MSE:HE1	1:D:64:ILE:HG13	2.02	0.42
1:D:12:MSE:HE3	1:D:12:MSE:HB3	1.83	0.42
1:A:130:LYS:NZ	1:B:145:LEU:O	2.53	0.42
1:D:12:MSE:HG3	1:D:36:HIS:CD2	2.54	0.42
1:B:78:ILE:O	1:B:80:PRO:HD3	2.20	0.42
1:A:96:ARG:NH2	4:A:536:HOH:O	2.53	0.41
1:B:18:ILE:HD11	1:B:43:ARG:HD2	2.01	0.41
1:C:128:ALA:N	1:C:129:PRO:CD	2.83	0.41
1:D:123:LEU:HA	1:D:124:PRO:C	2.41	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	172/188 (92%)	171 (99%)	1 (1%)	0	100	100
1	B	172/188 (92%)	170 (99%)	1 (1%)	1 (1%)	33	10
1	C	171/188 (91%)	170 (99%)	1 (1%)	0	100	100
1	D	171/188 (91%)	170 (99%)	1 (1%)	0	100	100
All	All	686/752 (91%)	681 (99%)	4 (1%)	1 (0%)	59	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	18	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/153 (92%)	139 (99%)	2 (1%)	78	57
1	B	143/153 (94%)	141 (99%)	2 (1%)	78	57
1	C	137/153 (90%)	135 (98%)	2 (2%)	76	54
1	D	134/153 (88%)	130 (97%)	4 (3%)	53	21
All	All	555/612 (91%)	545 (98%)	10 (2%)	71	46

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ASP
1	A	97	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	97	PHE
1	B	172	MSE
1	C	1	MSE
1	C	97	PHE
1	D	14	SER
1	D	53	PHE
1	D	97	PHE
1	D	172	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CIT	A	501	-	12,12,12	0.94	1 (8%)	17,17,17	1.67	6 (35%)
2	PO4	B	401	-	4,4,4	0.31	0	6,6,6	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CIT	C	502	-	12,12,12	0.89	0	17,17,17	3.59	10 (58%)
3	CIT	D	503	-	12,12,12	1.04	0	17,17,17	1.66	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIT	A	501	-	-	0/16/16/16	0/0/0/0
2	PO4	B	401	-	-	0/0/0/0	0/0/0/0
3	CIT	C	502	-	-	0/16/16/16	0/0/0/0
3	CIT	D	503	-	-	0/16/16/16	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	CIT	O5-C6	2.09	1.29	1.22

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	CIT	O6-C6-C3	9.77	127.09	112.89
3	C	502	CIT	O5-C6-C3	-6.13	113.74	122.20
3	C	502	CIT	O7-C3-C4	-4.20	100.75	109.22
3	C	502	CIT	C4-C3-C6	3.66	118.60	110.12
3	D	503	CIT	O6-C6-C3	3.22	117.58	112.89
3	C	502	CIT	O4-C5-C4	3.09	125.50	114.63
3	D	503	CIT	O2-C1-O1	-2.67	116.50	123.30
3	C	502	CIT	O2-C1-C2	2.64	123.95	114.63
3	C	502	CIT	O2-C1-O1	-2.61	116.66	123.30
3	D	503	CIT	O2-C1-C2	2.54	123.58	114.63
3	A	501	CIT	O7-C3-C2	2.52	114.30	109.22
3	D	503	CIT	O4-C5-C4	2.39	123.03	114.63
3	D	503	CIT	O4-C5-O3	-2.36	117.29	123.30
3	A	501	CIT	O6-C6-C3	2.33	116.28	112.89
3	A	501	CIT	O4-C5-O3	-2.32	117.39	123.30
3	A	501	CIT	O2-C1-C2	2.27	122.61	114.63
3	A	501	CIT	O7-C3-C6	-2.23	105.73	108.95
3	A	501	CIT	C3-C2-C1	2.16	119.01	113.77
3	C	502	CIT	C2-C3-C6	-2.15	105.13	110.12
3	C	502	CIT	O4-C5-O3	-2.11	117.93	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	CIT	C4-C3-C2	2.01	114.21	109.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/188 (91%)	-0.02	1 (0%) 86 90	7, 12, 25, 42	0
1	B	173/188 (92%)	0.04	2 (1%) 75 79	8, 13, 27, 64	0
1	C	173/188 (92%)	-0.02	3 (1%) 67 69	13, 19, 33, 63	0
1	D	172/188 (91%)	0.79	35 (20%) 1 1	10, 22, 36, 60	0
All	All	690/752 (91%)	0.20	41 (5%) 22 20	7, 17, 33, 64	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	128	ALA	6.8
1	D	1	MSE	4.3
1	D	17	GLU	4.2
1	D	15	ILE	4.2
1	D	152	ALA	4.1
1	D	125	GLY	3.9
1	D	53	PHE	3.9
1	D	131	VAL	3.9
1	D	127	VAL	3.7
1	D	172	MSE	3.5
1	D	14	SER	3.4
1	D	147	GLU	3.3
1	D	13	ASP	3.2
1	D	123	LEU	3.2
1	C	2	PHE	3.2
1	B	1	MSE	2.9
1	D	4	GLY	2.9
1	D	144	GLY	2.9
1	D	148	THR	2.9
1	A	14	SER	2.9
1	C	1	MSE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	5	ILE	2.8
1	C	16	GLY	2.7
1	D	150	GLU	2.7
1	D	155	ILE	2.6
1	D	149	GLU	2.6
1	D	136	PRO	2.5
1	D	168	ILE	2.5
1	D	35	PHE	2.4
1	D	153	ARG	2.4
1	D	154	GLU	2.4
1	D	101	ASP	2.4
1	D	137	GLY	2.4
1	B	173	LYS	2.3
1	D	166	SER	2.3
1	D	169	LEU	2.3
1	D	129	PRO	2.3
1	D	126	ALA	2.2
1	D	145	LEU	2.2
1	D	100	LEU	2.1
1	D	135	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CIT	C	502	13/13	0.16	2.76	37,44,54,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CIT	D	503	13/13	0.18	0.19	43,50,52,54	0
3	CIT	A	501	13/13	0.08	-0.87	22,27,34,35	0
2	PO4	B	401	5/5	0.07	-1.49	22,23,25,29	0

6.5 Other polymers

There are no such residues in this entry.