



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 01:02 AM GMT

PDB ID : 3VPS  
Title : Structure of a novel NAD dependent-NDP-hexosamine5,6-dehydratase, TunA, involved in tunicamycin biosynthesis  
Authors : Wyszynski, F.J.; Lee, S.S.; Yabe, T.; Wang, H.; Gomez-Escribano, J.P.; Bibb, M.J.; Lee, S.J.; Davies, G.J.; Davis, B.G.  
Deposited on : 2012-03-12  
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	321	
1	B	321	

## 2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 5003 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent epimerase/dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2315	1453	413	444	5			
1	B	301	Total	C	N	O	S	0	0	0
			2303	1447	410	441	5			

- Molecule 2 is URIDINE-DIPHOSPHATE-N-ACETYLGLUCOSAMINE (three-letter code: UD1) (formula:  $C_{17}H_{27}N_3O_{17}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			39	17	3	17	2		
2	B	1	Total	C	N	O	P	0	0
			39	17	3	17	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	125	Total	O	0	0
			125	125		
4	B	94	Total	O	0	0
			94	94		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NAD-dependent epimerase/dehydratase

Chain A: 



- Molecule 1: NAD-dependent epimerase/dehydratase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.45Å 51.06Å 67.80Å 98.13° 106.65° 94.36°	Depositor
Resolution (Å)	22.26 – 1.90 22.26 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (22.26-1.90) 99.4 (22.26-1.90)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.99 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.186 , 0.235 0.187 , 0.235	Depositor DCC
$R_{free}$ test set	2283 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.5	Xtriage
Anisotropy	0.084	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 63.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 44950 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5003	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UD1, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.19	10/2361 (0.4%)	0.98	8/3215 (0.2%)
1	B	1.05	7/2348 (0.3%)	1.02	13/3197 (0.4%)
All	All	1.12	17/4709 (0.4%)	1.00	21/6412 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	284	GLU	CG-CD	-9.57	1.37	1.51
1	A	262	GLN	CG-CD	-8.29	1.31	1.51
1	A	309	ASP	CB-CG	-7.26	1.36	1.51
1	A	36	VAL	CB-CG2	-7.15	1.37	1.52
1	A	262	GLN	CB-CG	-7.15	1.33	1.52
1	A	63	GLU	CD-OE1	-6.99	1.18	1.25
1	B	36	VAL	CB-CG1	-6.55	1.39	1.52
1	A	63	GLU	CB-CG	-6.24	1.40	1.52
1	A	49	GLU	CB-CG	-6.18	1.40	1.52
1	B	71	ARG	NE-CZ	-6.17	1.25	1.33
1	B	257	GLU	CG-CD	-6.12	1.42	1.51
1	B	187	ARG	CZ-NH1	-5.96	1.25	1.33
1	A	308	ASP	CB-CG	-5.62	1.40	1.51
1	A	187	ARG	CZ-NH1	-5.37	1.26	1.33
1	B	71	ARG	CB-CG	-5.32	1.38	1.52
1	A	143	TYR	CD1-CE1	5.18	1.47	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	292	GLU	CG-CD	5.17	1.59	1.51

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	187	ARG	NE-CZ-NH2	14.22	127.41	120.30
1	B	187	ARG	NE-CZ-NH1	-12.52	114.04	120.30
1	A	187	ARG	NE-CZ-NH2	11.69	126.15	120.30
1	A	187	ARG	NE-CZ-NH1	-10.08	115.26	120.30
1	A	9	ARG	NE-CZ-NH2	-7.95	116.33	120.30
1	A	36	VAL	CG1-CB-CG2	-7.37	99.11	110.90
1	A	262	GLN	N-CA-CB	-7.13	97.76	110.60
1	B	264	ARG	NE-CZ-NH1	-6.82	116.89	120.30
1	B	71	ARG	CG-CD-NE	-6.72	97.68	111.80
1	B	169	VAL	CG1-CB-CG2	-6.46	100.56	110.90
1	B	71	ARG	NE-CZ-NH1	-6.13	117.23	120.30
1	B	71	ARG	CD-NE-CZ	-6.13	115.02	123.60
1	A	169	VAL	CG1-CB-CG2	-5.72	101.75	110.90
1	B	247	ARG	NE-CZ-NH1	5.61	123.10	120.30
1	B	247	ARG	CG-CD-NE	5.57	123.49	111.80
1	A	262	GLN	CB-CG-CD	-5.56	97.13	111.60
1	B	284	GLU	OE1-CD-OE2	5.54	129.95	123.30
1	A	127	ASP	CB-CG-OD1	5.37	123.14	118.30
1	B	127	ASP	CB-CG-OD1	5.29	123.06	118.30
1	B	91	ASP	CB-CG-OD1	5.27	123.04	118.30
1	B	247	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	78	SER	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2315	0	0	19	0
1	B	2303	0	0	18	0
2	A	39	0	0	2	0
2	B	39	0	0	2	0
3	A	44	0	0	0	0
3	B	44	0	0	0	0
4	A	125	0	0	7	2
4	B	94	0	0	3	1
All	All	5003	0	0	37	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (37) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:308:ASP:OD1	1:A:309:ASP:N	1.83	1.10
1:A:309:ASP:OD1	1:A:309:ASP:O	1.77	1.00
1:B:262:GLN:N	1:B:262:GLN:OE1	2.00	0.95
1:B:309:ASP:O	1:B:310:ILE:CG1	2.30	0.79
1:A:8:HIS:NE2	4:A:618:HOH:O	2.23	0.71
1:A:260:ARG:NH2	4:A:620:HOH:O	2.24	0.71
1:B:9:ARG:NH2	4:B:530:HOH:O	2.24	0.69
1:B:308:ASP:C	1:B:308:ASP:OD1	2.30	0.69
1:A:309:ASP:CG	1:A:309:ASP:O	2.38	0.62
1:B:219:LYS:NZ	4:B:551:HOH:O	2.34	0.60
1:A:306:ASP:C	1:A:306:ASP:OD1	2.40	0.60
1:A:51:THR:CG2	4:A:610:HOH:O	2.50	0.59
1:A:71:ARG:NE	4:A:612:HOH:O	2.36	0.59
1:B:306:ASP:C	1:B:306:ASP:OD1	2.45	0.56
1:B:279:THR:O	1:B:283:GLY:CA	2.55	0.55
1:A:309:ASP:OD1	1:A:309:ASP:C	2.37	0.52
1:B:257:GLU:OE1	1:B:257:GLU:N	2.43	0.52
1:B:71:ARG:CG	1:B:71:ARG:NH1	2.68	0.49
1:B:308:ASP:O	1:B:308:ASP:OD1	2.30	0.49
1:A:305:ARG:NH2	4:A:559:HOH:O	2.46	0.48
1:A:25:ARG:NH2	4:A:533:HOH:O	2.47	0.48
1:B:173:ASN:ND2	4:B:506:HOH:O	2.48	0.47
1:B:84:ARG:O	1:B:87:LYS:CG	2.63	0.46
1:A:279:THR:O	1:A:283:GLY:CA	2.64	0.45
1:A:120:CYS:SG	2:A:400:UD1:O6'	2.74	0.45
1:A:184:LEU:N	2:A:400:UD1:O1A	2.50	0.44
1:B:184:LEU:N	2:B:400:UD1:O1A	2.51	0.44
1:B:205:GLU:OE2	1:B:243:ASN:ND2	2.51	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:9:ARG:NE	1:B:35:THR:OG1	2.51	0.43
1:A:57:LYS:NZ	1:A:66:ASP:OD2	2.51	0.43
1:B:8:HIS:CD2	1:B:33:GLU:OE1	2.72	0.43
1:A:69:ASP:OD2	4:A:618:HOH:O	2.22	0.42
1:B:264:ARG:NH1	2:B:400:UD1:O2B	2.52	0.42
1:B:57:LYS:NZ	1:B:66:ASP:OD2	2.53	0.42
1:A:296:ARG:NH1	1:A:300:GLU:OE1	2.53	0.42
1:A:308:ASP:CG	1:A:309:ASP:N	2.65	0.41
1:A:13:THR:OG1	1:A:75:HIS:CD2	2.73	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:A:597:HOH:O	4:A:617:HOH:O[1_655]	1.79	0.41
4:A:595:HOH:O	4:B:581:HOH:O[1_656]	1.98	0.22

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/321 (94%)	291 (97%)	9 (3%)	1 (0%)	50	37
1	B	297/321 (92%)	292 (98%)	5 (2%)	0	100	100
All	All	598/642 (93%)	583 (98%)	14 (2%)	1 (0%)	56	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	287	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of

similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/270 (94%)	240 (95%)	13 (5%)	33	19
1	B	252/270 (93%)	238 (94%)	14 (6%)	30	16
All	All	505/540 (94%)	478 (95%)	27 (5%)	32	18

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	ILE
1	A	161	SER
1	A	247	ARG
1	A	250	GLN
1	A	252	THR
1	A	253	SER
1	A	262	GLN
1	A	286	SER
1	A	292	GLU
1	A	295	ILE
1	A	306	ASP
1	A	307	LEU
1	A	309	ASP
1	B	51	THR
1	B	65	ARG
1	B	187	ARG
1	B	193	LEU
1	B	241	SER
1	B	247	ARG
1	B	252	THR
1	B	253	SER
1	B	281	GLN
1	B	284	GLU
1	B	292	GLU
1	B	306	ASP
1	B	308	ASP
1	B	310	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	UD1	A	400	-	41,41,41	1.17	2 (4%)	58,62,62	1.40	9 (15%)
3	NAD	A	401	-	48,48,48	1.62	7 (14%)	73,73,73	2.43	11 (15%)
2	UD1	B	400	-	41,41,41	1.12	3 (7%)	58,62,62	1.30	5 (8%)
3	NAD	B	401	-	48,48,48	1.32	7 (14%)	73,73,73	2.82	19 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UD1	A	400	-	-	0/25/63/63	0/3/3/3
3	NAD	A	401	-	-	0/30/62/62	0/3/5/5
2	UD1	B	400	-	-	0/25/63/63	0/3/3/3
3	NAD	B	401	-	-	0/30/62/62	0/3/5/5

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	NAD	O7N-C7N	6.44	1.39	1.24
2	A	400	UD1	C2-N1	5.17	1.44	1.38
2	B	400	UD1	C2-N1	4.04	1.42	1.38
3	B	401	NAD	O7N-C7N	3.77	1.33	1.24
3	A	401	NAD	C7N-N7N	-3.53	1.25	1.33
3	A	401	NAD	C2A-N3A	3.07	1.38	1.32
3	A	401	NAD	O4B-C1B	2.96	1.45	1.41
3	B	401	NAD	C2A-N1A	2.86	1.39	1.33
3	A	401	NAD	C2B-C1B	-2.82	1.49	1.53
3	A	401	NAD	C2A-N1A	2.72	1.39	1.33
2	B	400	UD1	O4B-C1B	2.72	1.45	1.41
3	B	401	NAD	O4D-C1D	2.54	1.45	1.41
3	B	401	NAD	C2D-C1D	-2.49	1.49	1.53
3	B	401	NAD	C2N-N1N	2.42	1.38	1.35
2	A	400	UD1	O1'-C1'	2.25	1.46	1.42
3	B	401	NAD	C4N-C3N	2.19	1.43	1.39
3	A	401	NAD	C2N-N1N	2.18	1.38	1.35
3	B	401	NAD	C2A-N3A	2.17	1.36	1.32
2	B	400	UD1	PB-O3A	2.11	1.63	1.59

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	NAD	N3A-C2A-N1A	-11.98	118.69	128.71
3	A	401	NAD	O4B-C1B-N9A	-11.55	97.70	108.44
3	B	401	NAD	C3N-C7N-N7N	10.65	129.89	117.77
3	A	401	NAD	N3A-C2A-N1A	-10.36	120.05	128.71
3	B	401	NAD	O4B-C1B-N9A	-9.87	99.26	108.44
3	B	401	NAD	O7N-C7N-N7N	-7.40	111.90	122.59
3	A	401	NAD	O7N-C7N-C3N	7.32	127.83	119.58
3	A	401	NAD	O7N-C7N-N7N	-6.54	113.15	122.59
2	B	400	UD1	N3-C2-N1	5.29	120.39	115.97
3	B	401	NAD	C4A-C5A-N7A	-4.40	105.75	109.52
2	A	400	UD1	N3-C2-N1	3.38	118.80	115.97
2	A	400	UD1	C5-C4-N3	3.26	122.84	116.70
3	B	401	NAD	C8A-N9A-C4A	3.20	109.34	106.90
2	A	400	UD1	O5'-C1'-O1'	-3.03	107.40	111.36
3	B	401	NAD	C2D-C1D-N1N	2.99	118.92	113.86
3	A	401	NAD	C8A-N9A-C4A	2.88	109.10	106.90
3	B	401	NAD	O4D-C1D-C2D	-2.87	102.37	106.77
3	B	401	NAD	C4D-O4D-C1D	2.84	112.83	109.75
3	A	401	NAD	O4D-C1D-C2D	-2.81	102.47	106.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	NAD	N3A-C4A-N9A	2.78	130.45	125.43
3	B	401	NAD	N7A-C8A-N9A	-2.74	106.60	114.36
2	A	400	UD1	O6'-C6'-C5'	-2.73	101.95	111.36
3	A	401	NAD	C1B-N9A-C4A	-2.66	122.05	126.64
3	B	401	NAD	C1B-N9A-C4A	-2.65	122.06	126.64
2	B	400	UD1	O6'-C6'-C5'	-2.65	102.24	111.36
2	A	400	UD1	O3A-PB-O1'	-2.58	96.97	103.48
2	B	400	UD1	C5-C4-N3	2.56	121.53	116.70
3	B	401	NAD	C2N-C3N-C4N	2.55	121.20	118.31
3	B	401	NAD	C8A-N7A-C5A	2.47	111.24	103.58
2	B	400	UD1	C5-C6-N1	-2.46	118.42	121.21
3	B	401	NAD	C4B-O4B-C1B	-2.39	107.16	109.75
3	A	401	NAD	N7A-C8A-N9A	-2.37	107.65	114.36
2	B	400	UD1	C1'-O5'-C5'	-2.37	109.12	113.73
3	B	401	NAD	O3D-C3D-C4D	-2.36	104.11	111.08
2	A	400	UD1	C2B-C1B-N1	2.34	119.29	113.26
3	B	401	NAD	C6N-N1N-C2N	-2.33	119.41	122.04
2	A	400	UD1	C2-N1-C1B	2.32	119.66	118.21
2	A	400	UD1	O5'-C5'-C4'	2.31	114.04	109.76
3	B	401	NAD	C2A-N1A-C6A	2.29	122.91	118.77
3	A	401	NAD	N3A-C4A-N9A	2.29	129.56	125.43
3	A	401	NAD	C2B-C1B-N9A	2.22	118.96	113.27
2	A	400	UD1	C1'-C2'-N2'	-2.13	106.98	111.02
3	B	401	NAD	C5N-C4N-C3N	-2.04	117.67	120.32
3	A	401	NAD	C4B-O4B-C1B	-2.03	107.55	109.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	303/321 (94%)	0.47	29 (9%) 8 8	21, 36, 65, 84	0
1	B	301/321 (93%)	0.50	32 (10%) 7 6	23, 39, 69, 83	0
All	All	604/642 (94%)	0.48	61 (10%) 8 7	21, 38, 68, 84	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	286	SER	7.9
1	B	51	THR	5.8
1	B	50	GLY	5.8
1	A	87	LYS	5.7
1	A	287	GLY	5.7
1	B	283	GLY	5.7
1	B	87	LYS	5.6
1	A	257	GLU	5.5
1	B	287	GLY	5.1
1	B	47	PRO	4.9
1	B	310	ILE	4.5
1	A	285	ARG	4.4
1	A	284	GLU	4.3
1	A	161	SER	4.3
1	B	284	GLU	4.2
1	A	308	ASP	4.1
1	A	256	ALA	3.9
1	B	49	GLU	3.9
1	B	254	PRO	3.8
1	B	262	GLN	3.8
1	A	254	PRO	3.7
1	B	88	GLN	3.6
1	B	308	ASP	3.5
1	B	309	ASP	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	288	GLY	3.4
1	B	162	VAL	3.4
1	A	307	LEU	3.4
1	B	48	PRO	3.4
1	A	306	ASP	3.3
1	A	88	GLN	3.1
1	B	307	LEU	3.1
1	B	257	GLU	3.1
1	B	265	PRO	3.0
1	A	262	GLN	3.0
1	B	306	ASP	2.9
1	B	251	ALA	2.9
1	B	260	ARG	2.9
1	B	197	GLU	2.8
1	A	195	ARG	2.7
1	A	185	VAL	2.7
1	B	266	ASN	2.6
1	A	309	ASP	2.6
1	B	255	ALA	2.5
1	A	197	GLU	2.5
1	B	259	ALA	2.4
1	A	186	PRO	2.3
1	B	43	PRO	2.3
1	B	304	SER	2.3
1	A	50	GLY	2.2
1	A	84	ARG	2.2
1	B	161	SER	2.2
1	A	255	ALA	2.2
1	B	303	GLN	2.2
1	B	185	VAL	2.2
1	A	188	LEU	2.2
1	B	188	LEU	2.2
1	A	12	ILE	2.1
1	A	20	GLY	2.0
1	A	183	ALA	2.0
1	A	259	ALA	2.0
1	A	263	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAD	A	401	44/44	0.11	-0.54	21,25,29,34	0
3	NAD	B	401	44/44	0.10	-0.56	28,31,33,38	0
2	UD1	A	400	39/39	0.10	-0.71	33,40,50,53	0
2	UD1	B	400	39/39	0.09	-1.16	36,43,53,56	0

### 6.5 Other polymers

There are no such residues in this entry.