



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 02:45 AM GMT

PDB ID : 2W2K
Title : CRYSTAL STRUCTURE OF THE APO FORMS OF RHODOTORULA GRAMINIS D-MANDELATE DEHYDROGENASE AT 1.8Å.
Authors : Vachieri, S.G.; Cole, A.R.; Bagneris, C.; Baker, D.P.; Fewson, C.A.; Basak, A.K.
Deposited on : 2008-11-02
Resolution : 1.85 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1269 (1.86-1.86)
Clashscore	79885	1470 (1.86-1.86)
Ramachandran outliers	78287	1451 (1.86-1.86)
Sidechain outliers	78261	1451 (1.86-1.86)
RSRZ outliers	66119	1269 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	348	
2	B	348	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6062 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-MANDELATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	50	1	0
			2658	1688	468	495	7			

- Molecule 2 is a protein called D-MANDELATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	344	Total	C	N	O	S	41	2	0
			2663	1693	470	493	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	288	ASN	GLN	CONFLICT	UNP Q7LLW9


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	357	Total	O	0	0
			357	357		
3	B	384	Total	O	0	0
			384	384		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: D-MANDELATE DEHYDROGENASE



• Molecule 2: D-MANDELATE DEHYDROGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.03Å 81.07Å 93.95Å 90.00° 96.27° 90.00°	Depositor
Resolution (Å)	25.96 – 1.85 25.27 – 1.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (25.96-1.85) 94.0 (25.27-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.27 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.154 , 0.199 0.154 , 0.200	Depositor DCC
R_{free} test set	3020 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	14.4	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 57.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 59832 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6062	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.38	9/2718 (0.3%)	1.24	15/3682 (0.4%)
2	B	1.18	6/2725 (0.2%)	0.77	8/3691 (0.2%)
All	All	1.29	15/5443 (0.3%)	1.03	23/7373 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	162	ARG	CZ-NH2	46.71	1.93	1.33
2	B	92	ASP	CB-CG	-35.26	0.77	1.51
1	A	156	LYS	CE-NZ	-18.15	1.03	1.49
1	A	176	LYS	CE-NZ	-13.60	1.15	1.49
1	A	204	LYS	CB-CG	-10.98	1.23	1.52
1	A	41	LYS	CD-CE	-10.69	1.24	1.51
1	A	14	ARG	CG-CD	8.92	1.74	1.51
2	B	189	LYS	CE-NZ	-8.78	1.27	1.49
2	B	48	ARG	CG-CD	-8.54	1.30	1.51
1	A	291	LYS	CB-CG	-7.83	1.31	1.52
2	B	295	GLU	CG-CD	-7.39	1.40	1.51
1	A	162	ARG	CZ-NH1	7.39	1.42	1.33
2	B	240	GLU	CG-CD	-6.37	1.42	1.51
1	A	271	LYS	CE-NZ	6.35	1.65	1.49
2	B	145	GLU	CG-CD	-5.64	1.43	1.51

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	ARG	NE-CZ-NH1	47.76	144.18	120.30
1	A	162	ARG	NH1-CZ-NH2	-31.45	84.81	119.40
2	B	92	ASP	CA-CB-CG	11.68	139.09	113.40
1	A	45	ARG	NE-CZ-NH1	10.18	125.39	120.30
2	B	92	ASP	CB-CG-OD2	10.09	127.38	118.30
1	A	45	ARG	NE-CZ-NH2	-9.98	115.31	120.30
1	A	156	LYS	CD-CE-NZ	9.49	133.53	111.70
1	A	200	ALA	CB-CA-C	7.60	121.50	110.10
2	B	162	ARG	NE-CZ-NH2	-7.41	116.60	120.30
1	A	176	LYS	CD-CE-NZ	7.36	128.63	111.70
1	A	14	ARG	CB-CG-CD	-6.91	93.63	111.60
1	A	271	LYS	CD-CE-NZ	-6.61	96.51	111.70
2	B	156	LYS	CD-CE-NZ	-6.54	96.67	111.70
1	A	257	ARG	CB-CG-CD	6.46	128.40	111.60
1	A	2	PRO	N-CA-CB	6.36	110.93	103.30
1	A	291	LYS	CA-CB-CG	6.16	126.96	113.40
2	B	2	PRO	N-CA-CB	6.00	110.50	103.30
2	B	162	ARG	NE-CZ-NH1	5.95	123.27	120.30
1	A	232	LYS	CB-CA-C	-5.73	98.94	110.40
2	B	189	LYS	CD-CE-NZ	5.35	124.00	111.70
1	A	204	LYS	CA-CB-CG	5.21	124.87	113.40
1	A	9	LEU	CA-CB-CG	5.12	127.07	115.30
2	B	92	ASP	CB-CG-OD1	-5.04	113.77	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162	ARG	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2658	0	2652	17	0
2	B	2663	0	2671	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	357	0	0	5	0
3	B	384	0	0	3	0
All	All	6062	0	5323	29	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (29) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:231:MET:HE2	2:B:233:LEU:H	1.55	0.71
1:A:311:GLU:OE2	2:B:159:HIS:HD2	1.80	0.64
2:B:323:ASN:ND2	2:B:335:LEU:H	1.99	0.59
2:B:323:ASN:HD21	2:B:335:LEU:H	1.49	0.58
1:A:270:LEU:HB3	1:A:296[B]:MET:CE	2.33	0.58
2:B:159:HIS:HE1	3:B:2101:HOH:O	1.88	0.57
1:A:323:ASN:ND2	1:A:335:LEU:H	2.03	0.57
1:A:159:HIS:HD2	2:B:311:GLU:OE2	1.89	0.55
1:A:159:HIS:HE1	3:A:2210:HOH:O	1.90	0.54
2:B:286:GLU:OE2	2:B:304:HIS:HD2	1.91	0.54
2:B:270:LEU:HB3	2:B:296[B]:MET:HE3	1.89	0.53
1:A:286:GLU:OE1	1:A:304:HIS:HD2	1.91	0.53
2:B:2:PRO:N	3:B:2001:HOH:O	2.42	0.52
1:A:298:HIS:HE1	3:A:2201:HOH:O	1.91	0.52
2:B:270:LEU:HB3	2:B:296[B]:MET:CE	2.40	0.51
1:A:2:PRO:HA	3:A:2001:HOH:O	2.10	0.51
2:B:221:SER:O	2:B:247:LYS:HE3	2.12	0.50
1:A:270:LEU:HB3	1:A:296[B]:MET:HE3	1.94	0.49
1:A:270:LEU:HB3	1:A:296[B]:MET:HE1	1.95	0.49
2:B:283:HIS:HE1	2:B:302:THR:O	1.96	0.49
1:A:323:ASN:HD21	1:A:335:LEU:H	1.61	0.47
1:A:37:HIS:HD2	1:A:66:SER:O	2.00	0.46
2:B:314:HIS:HD2	3:B:2078:HOH:O	2.00	0.44
1:A:184:HIS:HE1	3:A:2228:HOH:O	2.01	0.42
2:B:100:ASN:ND2	2:B:340:LYS:H	2.16	0.42
1:A:194:ASP:OD2	1:A:195:VAL:N	2.52	0.42
1:A:283:HIS:HE1	1:A:302:THR:O	2.03	0.42
1:A:247:LYS:HD3	3:A:2250:HOH:O	2.20	0.41
1:A:140:ARG:NH2	2:B:294:ILE:O	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/348 (99%)	335 (97%)	9 (3%)	0	100	100
2	B	344/348 (99%)	335 (97%)	9 (3%)	0	100	100
All	All	688/696 (99%)	670 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/279 (99%)	275 (99%)	2 (1%)	91	87
2	B	278/279 (100%)	276 (99%)	2 (1%)	91	87
All	All	555/558 (100%)	551 (99%)	4 (1%)	91	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	PHE
1	A	217	LEU
2	B	45	ARG
2	B	84	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	HIS
1	A	151	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	159	HIS
1	A	184	HIS
1	A	235	HIS
1	A	283	HIS
1	A	298	HIS
1	A	304	HIS
1	A	323	ASN
2	B	15	HIS
2	B	42	GLN
2	B	100	ASN
2	B	148	ASN
2	B	151	HIS
2	B	159	HIS
2	B	283	HIS
2	B	304	HIS
2	B	314	HIS
2	B	323	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/348 (99%)	-0.32	5 (1%) 72 71	6, 13, 28, 49	18 (5%)
2	B	344/348 (98%)	-0.48	0 100 100	5, 11, 23, 40	14 (4%)
All	All	689/696 (98%)	-0.40	5 (0%) 84 85	5, 12, 25, 49	32 (4%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	200	ALA	4.1
1	A	230	TYR	3.0
1	A	199	ASP	2.7
1	A	2	PRO	2.5
1	A	346	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.