



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 12:41 PM GMT

PDB ID : 2WIF  
Title : AGED FORM OF HUMAN BUTYRYLCHOLINESTERASEINHIBITED  
BY TABUN ANALOGUE TA1  
Authors : Carletti, E.; Aurbek, N.; Gillon, E.; Loiodice, M.; Nicolet, Y.; Fontecilla, J.;  
Masson, P.; Thiermann, H.; Nachon, F.; Worek, F.  
Deposited on : 2009-05-11  
Resolution : 2.25 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	529	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	SO4	A	1536	-	X
8	NAG	A	1543	-	X
8	NAG	A	1544	-	X

## 2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 4747 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHOLINESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	527	4213	2720	710	768	15	7	2	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	455	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	481	GLN	ASN	ENGINEERED MUTATION	UNP P06276


- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Cl	0	0
			3	3		

- Molecule 4 is DIETHYLPHOSPHORAMIDICACID (three-letter code: TN6) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	8	4	1	2	1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0
5	A	1	5	4	1	0	0

- Molecule 6 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	3	Total	C	N	O	0	0
			38	22	2	14		
6	A	3	Total	C	N	O	0	0
			38	22	2	14		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	455	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	481	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	17	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	455	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	481	GLN	ASN	ENGINEERED MUTATION	UNP P06276


- Molecule 7 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	2	Total	C	N	O	0	0
			24	14	1	9		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	455	GLN	ASN	ENGINEERED MUTATION	UNP P06276
A	481	GLN	ASN	ENGINEERED MUTATION	UNP P06276

- Molecule 8 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	370	Total	O	0	0
			370	370		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CHOLINESTERASE

Chain A:



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	154.39Å 154.39Å 126.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.67 – 2.25 47.67 – 2.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.67-2.25) 96.8 (47.67-2.25)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.88 (at 2.24Å)	Xtrriage
Refinement program	REFMAC 5.4.0069	Depositor
R, $R_{free}$	0.207 , 0.275 0.206 , 0.275	Depositor DCC
$R_{free}$ test set	1410 reflections (4.17%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.1	Xtrriage
Anisotropy	0.116	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 51.3	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Outliers	0 of 35241 reflections	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4747	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TN6, NAG, CL, NA, SO4, FUL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.21	12/4340 (0.3%)	1.03	9/5892 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	237	TYR	CB-CG	-10.71	1.35	1.51
1	A	497	GLU	CD-OE2	8.22	1.34	1.25
1	A	404	GLU	CG-CD	7.96	1.63	1.51
1	A	400	CYS	CB-SG	-7.70	1.69	1.82
1	A	367	GLU	CG-CD	6.02	1.60	1.51

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	515	ARG	NE-CZ-NH1	9.67	125.14	120.30
1	A	515	ARG	NE-CZ-NH2	-8.51	116.05	120.30
1	A	434	MET	CG-SD-CE	6.93	111.29	100.20
1	A	274	LEU	CA-CB-CG	6.05	129.21	115.30
1	A	424	ARG	NE-CZ-NH2	-5.38	117.61	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	361	VAL	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	360	GLY	Peptide

## 5.2 Close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4213	0	4110	146	0
2	A	1	0	0	0	0
3	A	3	0	0	1	0
4	A	8	0	11	3	0
5	A	10	0	0	0	0
6	A	76	0	68	1	0
7	A	24	0	22	2	0
8	A	42	0	39	5	0
9	A	370	0	0	25	0
All	All	4747	0	4250	151	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 18.

The worst 5 of 151 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:377:VAL:O	1:A:377:VAL:HG23	1.62	0.99
1:A:320:GLY:HA3	1:A:419:TYR:CE1	1.96	0.99
1:A:282:TYR:HD1	1:A:282:TYR:O	1.45	0.97
1:A:51:LYS:N	9:A:2044:HOH:O	1.96	0.96
1:A:518:GLN:H	1:A:518:GLN:HE21	1.00	0.96

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	527/529 (100%)	489 (93%)	33 (6%)	5 (1%)	25 20

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	361	VAL
1	A	453	ARG
1	A	496	THR
1	A	506	GLU
1	A	51	LYS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	453/454 (100%)	421 (93%)	32 (7%)	21 19

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	268	ASP
1	A	286	LEU
1	A	506	GLU
1	A	282	TYR
1	A	287	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	275	ASN
1	A	289	ASN
1	A	380	GLN
1	A	518	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
6	NAG	A	1537	1,6	12,14,15	1.34	2 (16%)	15,19,21	2.35	4 (26%)
6	NAG	A	1538	6	12,14,15	0.82	1 (8%)	15,19,21	2.08	3 (20%)
6	FUL	A	1539	6	9,10,11	1.11	0	10,14,16	1.80	2 (20%)
7	NAG	A	1540	1,7	12,14,15	0.57	0	15,19,21	2.80	8 (53%)
7	FUL	A	1541	7	9,10,11	0.99	1 (11%)	10,14,16	2.10	4 (40%)
6	NAG	A	1545	1,6	12,14,15	0.84	1 (8%)	15,19,21	1.08	1 (6%)
6	NAG	A	1546	6	12,14,15	0.45	0	15,19,21	1.42	3 (20%)
6	FUL	A	1547	6	9,10,11	0.92	1 (11%)	10,14,16	1.64	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	1537	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	1538	6	-	0/6/23/26	0/1/1/1
6	FUL	A	1539	6	-	0/0/17/20	0/1/1/1
7	NAG	A	1540	1,7	-	0/6/23/26	0/1/1/1
7	FUL	A	1541	7	-	0/0/17/20	0/1/1/1
6	NAG	A	1545	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	1546	6	-	0/6/23/26	0/1/1/1
6	FUL	A	1547	6	-	0/0/17/20	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1537	NAG	C2-N2	-3.43	1.42	1.46
7	A	1541	FUL	O5-C5	-2.16	1.41	1.45
6	A	1538	NAG	O5-C5	-2.13	1.41	1.45
6	A	1545	NAG	O5-C5	-2.11	1.41	1.45
6	A	1547	FUL	O5-C5	-2.08	1.41	1.45

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1537	NAG	C3-C2-N2	-7.52	100.31	111.76
7	A	1540	NAG	O5-C5-C6	6.73	114.05	106.98
6	A	1538	NAG	O5-C5-C6	5.43	112.68	106.98
7	A	1540	NAG	C3-C4-C5	-4.04	103.00	110.20
7	A	1541	FUL	C4-C3-C2	-3.97	105.18	110.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	TN6	A	1532	1	5,7,8	6.42	3 (60%)	5,8,11	1.76	1 (20%)
5	SO4	A	1534	-	4,4,4	0.25	0	6,6,6	0.32	0
5	SO4	A	1536	-	4,4,4	0.26	0	6,6,6	0.70	0
8	NAG	A	1542	1	12,14,15	0.63	0	15,19,21	1.83	4 (26%)
8	NAG	A	1543	1	12,14,15	0.94	0	15,19,21	1.24	1 (6%)
8	NAG	A	1544	1	12,14,15	0.66	0	15,19,21	2.17	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TN6	A	1532	1	-	0/4/8/10	0/0/0/0
5	SO4	A	1534	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1536	-	-	0/0/0/0	0/0/0/0
8	NAG	A	1542	1	-	0/6/23/26	0/1/1/1
8	NAG	A	1543	1	-	0/6/23/26	0/1/1/1
8	NAG	A	1544	1	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1532	TN6	P-N	-13.92	1.53	1.71
4	A	1532	TN6	C1-N	-2.52	1.44	1.50
4	A	1532	TN6	C4-C3	2.24	1.63	1.49

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	1544	NAG	O5-C5-C6	6.20	113.48	106.98
8	A	1542	NAG	O5-C5-C6	3.80	110.97	106.98
8	A	1544	NAG	C3-C4-C5	3.72	116.84	110.20
8	A	1542	NAG	C3-C4-C5	3.51	116.48	110.20
8	A	1542	NAG	C2-N2-C7	-3.37	117.44	123.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	527/529 (99%)	0.09	22 (4%) 35 40	17, 34, 58, 76	9 (1%)

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	282	TYR	4.7
1	A	50	THR	4.5
1	A	378	ASP	4.3
1	A	453	ARG	4.2
1	A	3	ASP	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	FUL	A	1541	10/11	0.30	7.67	40,46,47,49	10
6	NAG	A	1538	14/15	0.23	7.01	62,67,70,70	0
6	FUL	A	1539	10/11	0.23	3.94	65,69,73,73	0
7	NAG	A	1540	14/15	0.21	2.92	54,59,61,64	0
6	FUL	A	1547	10/11	0.27	1.81	71,73,75,77	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	NAG	A	1545	14/15	0.22	0.23	67,72,79,79	0
6	NAG	A	1537	14/15	0.16	0.07	46,52,58,64	0
6	NAG	A	1546	14/15	0.37	-	65,76,78,78	0

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
8	NAG	A	1544	14/15	0.47	8.11	82,85,88,88	0
8	NAG	A	1543	14/15	0.19	3.53	45,60,64,66	0
5	SO4	A	1536	5/5	0.17	2.60	40,42,43,45	5
2	NA	A	1530	1/1	0.20	1.44	41,41,41,41	1
8	NAG	A	1542	14/15	0.30	1.20	73,77,79,82	0
3	CL	A	1531	1/1	0.19	1.09	58,58,58,58	0
4	TN6	A	1532	8/9	0.15	0.32	24,27,28,29	0
5	SO4	A	1534	5/5	0.10	-1.10	52,52,57,58	5
3	CL	A	1535	1/1	0.08	-1.43	62,62,62,62	0
3	CL	A	1533	1/1	0.11	-1.46	71,71,71,71	0

## 6.5 Other polymers (i)

There are no such residues in this entry.