



# Full wwPDB X-ray Structure Validation Report i

Feb 28, 2014 – 07:31 AM GMT

PDB ID : 3WIQ  
Title : Crystal structure of kojibiose phosphorylase complexed with kojibiose  
Authors : Okada, S.; Yamamoto, T.; Watanabe, H.; Nishimoto, T.; Chaen, H.; Fukuda, S.; Wakagi, T.; Fushinobu, S.  
Deposited on : 2013-09-24  
Resolution : 2.80 Å (reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1799 (2.80-2.80)
Clashscore	79885	2295 (2.80-2.80)
Ramachandran outliers	78287	2252 (2.80-2.80)
Sidechain outliers	78261	2254 (2.80-2.80)
RSRZ outliers	66119	1802 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	764	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	SO4	A	803	-	X

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6241 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kojibiose phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	751	6148	3960	1013	1154	21	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	757	GLY	-	EXPRESSION TAG	UNP A4XGP2
A	758	SER	-	EXPRESSION TAG	UNP A4XGP2
A	759	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	760	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	761	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	762	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	763	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	764	HIS	-	EXPRESSION TAG	UNP A4XGP2


- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	2	23	12	11	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	757	GLY	-	EXPRESSION TAG	UNP A4XGP2
A	758	SER	-	EXPRESSION TAG	UNP A4XGP2
A	759	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	760	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	761	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	762	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	763	HIS	-	EXPRESSION TAG	UNP A4XGP2
A	764	HIS	-	EXPRESSION TAG	UNP A4XGP2

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		

### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kojibiose phosphorylase

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	192.51Å 192.51Å 202.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 2.80 48.13 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.13-2.80) 98.7 (48.13-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.37 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.204 , 0.259 0.207 , 0.259	Depositor DCC
$R_{free}$ test set	2338 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	55.9	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 29.9	EDS
Estimated twinning fraction	0.000 for -l,-k,-h 0.000 for -h,l,k	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	1 of 46425 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6241	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, BGC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.85	1/6289 (0.0%)	0.96	10/8497 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	507	GLU	CD-OE1	5.23	1.31	1.25

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	116	ARG	NE-CZ-NH1	-6.64	116.98	120.30
1	A	239	ARG	NE-CZ-NH2	6.25	123.42	120.30
1	A	625	LYS	CD-CE-NZ	6.23	126.02	111.70
1	A	666	ASN	CB-CA-C	-5.67	99.07	110.40
1	A	658	THR	N-CA-CB	5.46	120.67	110.30
1	A	665	ASP	CB-CG-OD2	-5.34	113.49	118.30
1	A	625	LYS	CA-CB-CG	5.10	124.63	113.40
1	A	566	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	376	ARG	NE-CZ-NH2	-5.08	117.76	120.30
1	A	433	MET	CB-CG-SD	5.07	127.61	112.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	551	ASP	Peptide
1	A	749	ILE	Peptide

## 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6148	0	6079	89	0
2	A	23	0	21	1	0
3	A	5	0	0	0	0
4	A	65	0	0	1	0
All	All	6241	0	6100	89	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (89) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:5:GLU:O	1:A:6:ARG:HB2	1.62	0.97
1:A:352:PHE:O	1:A:356:THR:HB	1.67	0.94
1:A:733:ASN:HA	1:A:750:LYS:HE2	1.65	0.78
1:A:5:GLU:O	1:A:6:ARG:CB	2.30	0.77
1:A:750:LYS:CB	1:A:751:GLY:HA3	2.22	0.70
1:A:636:SER:HB3	1:A:658:THR:HG21	1.76	0.68
1:A:750:LYS:HB3	1:A:751:GLY:HA3	1.75	0.68
1:A:248:LEU:HD12	1:A:248:LEU:N	2.11	0.65
1:A:247:VAL:C	1:A:248:LEU:HD12	2.18	0.64
1:A:430:TYR:HA	1:A:500:TRP:CH2	2.33	0.63
1:A:21:PHE:O	1:A:25:PHE:HB2	1.97	0.63
1:A:709:GLU:O	1:A:710:LYS:CB	2.47	0.63
1:A:286:TYR:OH	1:A:316:SER:OG	2.14	0.62
1:A:750:LYS:CB	1:A:751:GLY:CA	2.78	0.61
1:A:239:ARG:HG3	1:A:239:ARG:HH21	1.66	0.61
1:A:747:LEU:HG	1:A:749:ILE:HD11	1.82	0.60
1:A:704:ASN:N	1:A:705:PRO:HD3	2.16	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:286:TYR:HH	1:A:316:SER:HG	1.47	0.59
1:A:29:ASN:HA	1:A:128:MET:HE2	1.85	0.58
1:A:337:TYR:C	1:A:338:LYS:HG2	2.23	0.58
1:A:709:GLU:O	1:A:710:LYS:HB2	2.04	0.57
1:A:128:MET:HE3	1:A:316:SER:HA	1.85	0.57
1:A:703:VAL:HG12	1:A:705:PRO:HD3	1.85	0.57
1:A:472:ARG:HD3	1:A:545:TYR:HA	1.86	0.57
1:A:117:ILE:HG22	1:A:143:GLU:HG3	1.86	0.57
1:A:39:GLU:OE2	1:A:138:TYR:OH	2.22	0.57
1:A:257:GLY:N	4:A:924:HOH:O	2.38	0.55
1:A:204:THR:HG22	1:A:205:LYS:N	2.22	0.55
1:A:551:ASP:O	1:A:552:LYS:HB2	2.07	0.55
1:A:450:SER:O	1:A:454:ILE:HG13	2.07	0.54
1:A:239:ARG:HH21	1:A:239:ARG:CG	2.20	0.54
1:A:63:THR:HG22	1:A:400:GLU:CD	2.30	0.52
1:A:749:ILE:HG22	1:A:750:LYS:H	1.73	0.52
1:A:352:PHE:CZ	1:A:356:THR:HG21	2.45	0.52
1:A:624:GLU:HG3	1:A:625:LYS:N	2.24	0.51
1:A:310:ASN:HA	1:A:679:LEU:HD22	1.92	0.51
1:A:248:LEU:N	1:A:248:LEU:CD1	2.72	0.51
1:A:6:ARG:O	1:A:305:LYS:NZ	2.43	0.51
1:A:634:SER:O	1:A:635:PRO:C	2.50	0.50
1:A:344:ASP:OD1	2:A:801:GLC:O6	2.25	0.49
1:A:187:LEU:C	1:A:187:LEU:HD23	2.33	0.49
1:A:29:ASN:HA	1:A:128:MET:CE	2.43	0.49
1:A:352:PHE:CE2	1:A:356:THR:HG21	2.49	0.48
1:A:350:LEU:HD23	1:A:431:ALA:HB1	1.95	0.48
1:A:68:LEU:HB3	1:A:69:PRO:HD2	1.96	0.48
1:A:116:ARG:HB3	1:A:145:TYR:CD2	2.49	0.47
1:A:193:ASP:O	1:A:194:LYS:HB2	2.16	0.47
1:A:720:TRP:CD1	1:A:745:VAL:HG21	2.49	0.46
1:A:747:LEU:HD12	1:A:748:LYS:H	1.80	0.46
1:A:350:LEU:HB3	1:A:351:PRO:HD3	1.98	0.46
1:A:22:GLU:HB3	1:A:35:ARG:NH2	2.31	0.46
1:A:570:LYS:O	1:A:571:GLU:HB2	2.16	0.46
1:A:129:ASN:HB3	1:A:285:GLU:HG3	1.97	0.45
1:A:187:LEU:HD23	1:A:188:GLY:N	2.32	0.45
1:A:621:ASP:OD1	1:A:647:GLU:OE2	2.34	0.45
1:A:569:ILE:O	1:A:570:LYS:O	2.35	0.45
1:A:734:VAL:O	1:A:734:VAL:CG2	2.64	0.44
1:A:533:GLU:HB2	1:A:534:PRO:HD3	1.98	0.44
1:A:749:ILE:HG22	1:A:750:LYS:N	2.32	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:750:LYS:HB2	1:A:751:GLY:CA	2.47	0.44
1:A:432:VAL:HB	1:A:453:ILE:HD11	1.98	0.44
1:A:548:TYR:HB3	1:A:549:HIS:CA	2.48	0.44
1:A:496:TYR:CD1	1:A:603:LEU:HD21	2.52	0.44
1:A:204:THR:CG2	1:A:205:LYS:N	2.82	0.43
1:A:166:PRO:HA	1:A:169:ARG:NH1	2.33	0.43
1:A:433:MET:HG2	1:A:433:MET:O	2.16	0.43
1:A:435:TYR:CE1	1:A:439:THR:HG21	2.54	0.43
1:A:180:ASP:HB2	1:A:219:THR:HG21	2.01	0.43
1:A:533:GLU:N	1:A:534:PRO:CD	2.82	0.42
1:A:750:LYS:HB2	1:A:751:GLY:C	2.40	0.42
1:A:343:TRP:O	1:A:344:ASP:C	2.56	0.42
1:A:156:ASP:OD1	1:A:156:ASP:C	2.58	0.42
1:A:137:LYS:HA	1:A:242:ILE:O	2.20	0.42
1:A:749:ILE:O	1:A:750:LYS:C	2.59	0.41
1:A:445:LEU:O	1:A:450:SER:OG	2.38	0.41
1:A:613:ASP:N	1:A:613:ASP:OD1	2.50	0.41
1:A:654:ASN:OD1	1:A:657:ARG:NH2	2.54	0.41
1:A:630:LYS:HD3	1:A:630:LYS:HA	1.78	0.41
1:A:108:LEU:HD23	1:A:108:LEU:HA	1.89	0.41
1:A:77:TYR:O	1:A:149:LEU:HA	2.21	0.41
1:A:280:PHE:O	1:A:281:GLU:C	2.59	0.41
1:A:330:LYS:HA	1:A:674:ILE:O	2.21	0.41
1:A:446:LEU:HD23	1:A:512:LEU:HD21	2.03	0.40
1:A:128:MET:CE	1:A:316:SER:HA	2.50	0.40
1:A:510:ASN:O	1:A:514:GLU:HG2	2.21	0.40
1:A:717:SER:O	1:A:718:ILE:HG23	2.20	0.40
1:A:565:LYS:HG3	1:A:566:ASP:N	2.36	0.40
1:A:319:ASN:HA	1:A:320:PRO:HD2	1.88	0.40
1:A:658:THR:HB	1:A:682:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	749/764 (98%)	688 (92%)	46 (6%)	15 (2%)	11 35

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	569	ILE
1	A	570	LYS
1	A	609	GLU
1	A	750	LYS
1	A	6	ARG
1	A	545	TYR
1	A	552	LYS
1	A	710	LYS
1	A	469	GLU
1	A	706	TRP
1	A	625	LYS
1	A	721	LYS
1	A	291	ASP
1	A	571	GLU
1	A	41	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	665/676 (98%)	623 (94%)	42 (6%)	25 59

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LYS
1	A	127	SER
1	A	219	THR
1	A	239	ARG
1	A	261	GLU
1	A	266	LYS
1	A	306	SER
1	A	338	LYS

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	343	TRP
1	A	356	THR
1	A	433	MET
1	A	440	ASP
1	A	450	SER
1	A	472	ARG
1	A	478	VAL
1	A	483	GLU
1	A	488	CYS
1	A	507	GLU
1	A	508	LEU
1	A	513	LEU
1	A	514	GLU
1	A	530	SER
1	A	539	GLU
1	A	552	LYS
1	A	565	LYS
1	A	568	VAL
1	A	570	LYS
1	A	574	GLN
1	A	582	GLU
1	A	609	GLU
1	A	613	ASP
1	A	615	THR
1	A	624	GLU
1	A	658	THR
1	A	664	GLU
1	A	678	SER
1	A	693	ILE
1	A	697	LYS
1	A	702	SER
1	A	734	VAL
1	A	742	LYS
1	A	746	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	667	GLN

### 5.3.3 RNA (i)

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	A	801	2	10,11,12	1.12	1 (10%)	11,15,17	1.92	4 (36%)
2	BGC	A	802	2	12,12,12	0.47	0	17,17,17	1.72	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	801	2	-	0/2/19/22	0/1/1/1
2	BGC	A	802	2	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	GLC	O5-C5	-2.87	1.40	1.45

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	802	BGC	C1-C2-C3	-4.63	103.20	110.53

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	GLC	C4-C3-C2	-3.62	105.65	110.50
2	A	801	GLC	O3-C3-C4	3.30	117.75	110.35
2	A	802	BGC	O2-C2-C1	2.95	116.17	109.89
2	A	801	GLC	O4-C4-C3	2.70	116.41	110.35
2	A	802	BGC	O5-C1-C2	-2.47	106.04	109.86
2	A	801	GLC	O5-C5-C4	-2.44	107.55	110.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	803	-	4,4,4	1.43	1 (25%)	6,6,6	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	803	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	803	SO4	O2-S	2.49	1.55	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	751/764 (98%)	-0.36	8 (1%) 77 78	35, 52, 82, 150	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	548	TYR	7.8
1	A	709	GLU	3.2
1	A	552	LYS	2.9
1	A	549	HIS	2.7
1	A	751	GLY	2.6
1	A	568	VAL	2.4
1	A	224	TYR	2.3
1	A	550	PRO	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	A	802	12/12	0.13	0.10	47,55,59,71	0
2	GLC	A	801	11/12	0.13	-0.68	40,45,54,54	0

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	803	5/5	0.20	2.38	55,55,74,83	0

## 6.5 Other polymers

There are no such residues in this entry.