



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 12:15 AM GMT

PDB ID : 2WNU  
Title : COMPLEX BETWEEN C1Q GLOBULAR HEADS AND HEPARAN SULFATE  
Authors : Garlatti, V.; Chouquet, A.; Lunardi, T.; Thielens, N.M.; Arlaud, G.J.; Gaboriaud, C.  
Deposited on : 2009-07-20  
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	134	
1	D	134	
2	B	136	
2	E	136	
3	C	131	
3	F	131	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	NAG	A	1223	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6514 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called COMPLEMENT C1Q SUBCOMPONENT SUBUNIT A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	S	0	3	0
			1056	676	181	194	5			
1	D	133	Total	C	N	O	S	0	3	0
			1056	676	181	194	5			

- Molecule 2 is a protein called COMPLEMENT C1Q SUBCOMPONENT SUBUNIT B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	0	2	1
			1062	670	189	196	7			
2	E	133	Total	C	N	O	S	3	2	1
			1060	669	187	197	7			

- Molecule 3 is a protein called COMPLEMENT C1Q SUBCOMPONENT SUBUNIT C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	129	Total	C	N	O	S	0	5	0
			1027	659	173	191	4			
3	F	129	Total	C	N	O	S	0	5	0
			1027	659	173	191	4			

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	F	2	Total	C	N	O	S	0	0
			35	12	1	19	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	31	Total	O	0	0
			31	31		
6	B	19	Total	O	0	0
			19	19		
6	C	39	Total	O	0	0
			39	39		
6	D	33	Total	O	0	0
			33	33		
6	E	19	Total	O	0	0
			19	19		
6	F	35	Total	O	0	0
			35	35		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT A




- Molecule 1: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT A




- Molecule 2: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT B



- Molecule 2: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT B



- Molecule 3: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT C



- Molecule 3: COMPLEMENT C1Q SUBCOMPONENT SUBUNIT C





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.09Å 48.16Å 87.73Å 92.86° 92.20° 113.64°	Depositor
Resolution (Å)	19.65 – 2.30 19.65 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.65-2.30) 91.1 (19.65-2.30)	Depositor EDS
$R_{merge}$	0.01	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.39 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.198 , 0.256 0.198 , 0.255	Depositor DCC
$R_{free}$ test set	1451 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.3	Xtriage
Anisotropy	0.607	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 14.4	EDS
Estimated twinning fraction	0.023 for -k,-h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 29003 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6514	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IDU, NAG, SGN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/1102	0.59	0/1494
1	D	0.46	0/1102	0.60	0/1494
2	B	0.49	0/1087	0.59	0/1468
2	E	0.92	3/1085 (0.3%)	0.69	1/1466 (0.1%)
3	C	0.50	0/1074	0.60	0/1462
3	F	0.50	0/1074	0.61	0/1462
All	All	0.58	3/6524 (0.0%)	0.61	1/8846 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
5	F	1	0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	92	THR	CB-OG1	17.59	1.78	1.43
2	E	149	SER	CB-OG	-14.71	1.23	1.42
2	E	92	THR	CB-CG2	11.01	1.88	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	92	THR	OG1-CB-CG2	-12.17	82.01	110.00

All (1) chirality outliers are listed below:



Mol	Chain	Res	Type	Atom
5	F	1218	IDU	C1

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1056	0	1020	19	0
1	D	1056	0	1020	17	0
2	B	1062	0	1037	19	0
2	E	1060	0	1032	11	0
3	C	1027	0	1003	23	0
3	F	1027	0	1003	25	0
4	A	15	0	15	0	0
5	F	35	0	15	0	0
6	A	31	0	0	6	0
6	B	19	0	0	4	0
6	C	39	0	0	1	0
6	D	33	0	0	4	0
6	E	19	0	0	3	0
6	F	35	0	0	8	0
All	All	6514	0	6145	97	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (97) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:E:180:VAL:HG11	6:F:2028:HOH:O	1.61	1.00
2:B:159[B]:ARG:NH2	2:B:188:LYS:O	2.12	0.83
1:A:111:PHE:CD2	6:A:2001:HOH:O	2.37	0.77
2:B:179:GLN:OE1	6:B:2014:HOH:O	2.04	0.73
3:F:175:ASN:OD1	6:F:2028:HOH:O	2.05	0.73
2:B:155:VAL:HB	6:B:2010:HOH:O	1.88	0.72
1:D:218:LEU:CD2	6:D:2033:HOH:O	2.37	0.71

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:218:LEU:HD21	6:D:2033:HOH:O	1.90	0.71
1:A:98:ILE:HA	6:A:2001:HOH:O	1.92	0.69
1:A:111:PHE:HD2	6:A:2001:HOH:O	1.71	0.69
1:D:206:GLN:HG2	6:D:2030:HOH:O	1.93	0.69
3:C:146:HIS:HD2	3:C:148:ALA:O	1.77	0.66
2:B:93:GLN:HG2	2:B:137:VAL:HG21	1.80	0.64
1:A:97:ALA:O	6:A:2001:HOH:O	2.16	0.63
3:C:99:GLN:OE1	3:F:97:THR:HG21	1.99	0.63
3:C:198:ASP:OD2	3:C:200[B]:VAL:HG22	2.00	0.62
3:F:93:VAL:HG23	3:F:212:PHE:HB3	1.81	0.61
3:C:147:THR:OG1	3:C:200[B]:VAL:HG23	2.01	0.60
3:F:147:THR:HG1	3:F:200[B]:VAL:HG23	1.68	0.59
2:B:94:LYS:NZ	3:C:217:ASP:HB3	2.18	0.59
3:C:89:LYS:HG2	6:C:2012:HOH:O	2.01	0.59
1:D:98:ILE:HG22	1:D:213[A]:VAL:HG22	1.85	0.59
3:C:93:VAL:HG23	3:C:212:PHE:HB3	1.84	0.59
1:A:144:LEU:HD23	1:A:205:TYR:CD2	2.40	0.57
3:F:147:THR:OG1	3:F:200[B]:VAL:HG23	2.06	0.56
1:D:165[B]:LEU:HD13	3:F:206:ASP:CB	2.36	0.55
2:E:125:ASN:ND2	6:E:2007:HOH:O	2.39	0.55
2:B:219:LEU:HD21	2:B:222:PRO:HG3	1.89	0.55
2:E:144:THR:HG22	6:E:2012:HOH:O	2.07	0.54
3:F:169:SER:HA	6:F:2025:HOH:O	2.08	0.53
3:C:95:THR:CG2	3:C:116:LEU:HD12	2.40	0.52
3:F:95:THR:HG22	3:F:116:LEU:HD12	1.92	0.52
1:A:114:VAL:HG11	1:A:117:ASN:HB2	1.91	0.52
1:A:175:LEU:HD13	6:B:2012:HOH:O	2.09	0.52
3:F:146:HIS:HD2	3:F:148:ALA:O	1.93	0.52
3:F:111[A]:ARG:NE	3:F:125:THR:O	2.44	0.51
1:D:92:ARG:N	1:D:93:PRO:HD3	2.25	0.51
3:C:95:THR:HG22	3:C:116:LEU:HD12	1.92	0.51
2:E:161:ARG:NH1	2:E:191:GLN:O	2.43	0.51
2:B:97:PHE:O	2:B:216:SER:HA	2.10	0.51
3:C:147:THR:HG1	3:C:200[B]:VAL:HG23	1.76	0.50
3:F:137:LEU:HD21	3:F:180:LEU:HD22	1.94	0.50
1:A:98:ILE:HG22	1:A:213[A]:VAL:HG22	1.94	0.49
2:B:159[B]:ARG:HG2	2:B:193:GLU:OE1	2.12	0.49
3:F:99:GLN:HG2	6:F:2002:HOH:O	2.12	0.49
1:A:92:ARG:N	1:A:93:PRO:HD3	2.28	0.49
2:E:94:LYS:NZ	3:F:217:ASP:HB3	2.29	0.48
1:A:118:GLN:HG2	2:B:140:LEU:HD11	1.95	0.47
2:E:173:TYR:OH	2:E:202:LYS:HE3	2.14	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:94:LYS:HZ3	3:C:217:ASP:HB3	1.79	0.47
3:F:97:THR:HG23	3:F:114:ALA:HB3	1.96	0.47
1:A:165[A]:LEU:HD11	3:C:116:LEU:HD21	1.96	0.47
3:C:153:LEU:N	3:C:153:LEU:HD12	2.30	0.47
3:C:146:HIS:CD2	3:C:148:ALA:O	2.64	0.47
2:B:133:PHE:O	2:B:194:ASN:HA	2.15	0.47
1:D:142:GLN:OE1	1:D:213[B]:VAL:HG13	2.15	0.47
1:A:177:GLN:HA	2:B:172:ASP:OD2	2.15	0.47
2:B:96:ALA:HB1	3:C:180:LEU:HD13	1.97	0.46
1:A:162:ARG:HB3	3:F:111[A]:ARG:NH1	2.30	0.46
2:B:162:GLU:OE2	6:B:2011:HOH:O	2.21	0.46
1:A:194:TRP:HE1	1:A:196:GLU:HG3	1.80	0.46
2:E:94:LYS:HZ1	3:F:217:ASP:HB3	1.80	0.46
2:E:109:ARG:HD2	6:E:2003:HOH:O	2.16	0.45
1:D:118:GLN:HG2	2:E:140:LEU:HD11	1.98	0.45
3:F:99:GLN:CG	6:F:2002:HOH:O	2.64	0.45
1:A:200:LYS:HD3	1:A:200:LYS:HA	1.80	0.44
1:A:194:TRP:NE1	1:A:196:GLU:HG3	2.33	0.44
2:B:162:GLU:HG2	2:B:162:GLU:H	1.50	0.43
1:D:130:VAL:HG22	1:D:192:GLN:HG2	1.99	0.43
2:E:97:PHE:O	2:E:216:SER:HA	2.19	0.43
2:E:161:ARG:HH12	2:E:190:GLU:HB3	1.83	0.43
1:A:125:HIS:HD2	6:A:2011:HOH:O	2.01	0.43
1:D:165[B]:LEU:HD13	3:F:206:ASP:HB3	2.01	0.43
2:B:120:THR:HG21	3:C:137:LEU:HD11	2.01	0.43
1:D:183[A]:MET:HE3	3:F:116:LEU:HD13	2.02	0.42
3:C:135:PRO:HD3	3:C:185:VAL:HG22	2.02	0.42
3:F:102:GLN:NE2	6:F:2004:HOH:O	2.52	0.42
1:D:144:LEU:HD11	1:D:176:PHE:HB3	2.02	0.42
1:D:128:ARG:HD3	1:D:192:GLN:OE1	2.19	0.42
1:D:138:TYR:O	1:D:216:GLY:HA2	2.20	0.42
3:C:110:ILE:HD11	3:C:193:VAL:HG13	2.02	0.42
1:D:114:VAL:HG11	1:D:117:ASN:HB2	2.01	0.42
3:F:170:LYS:HE3	6:F:2024:HOH:O	2.19	0.42
1:D:165[B]:LEU:HD12	3:F:208[B]:VAL:HG23	2.01	0.42
2:B:143:PHE:O	2:B:184:GLY:HA2	2.20	0.41
2:B:94:LYS:HZ1	3:C:217:ASP:HB3	1.85	0.41
3:C:99:GLN:OE1	3:F:97:THR:CG2	2.65	0.41
3:C:89:LYS:HD2	3:C:120:GLN:NE2	2.36	0.41
3:F:194:ASN:HB2	6:F:2031:HOH:O	2.20	0.41
1:A:158:ARG:HH11	1:A:158:ARG:HG2	1.85	0.41
2:B:201:ASP:HB3	2:B:202:LYS:HD2	2.03	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:C:139:TYR:O	3:C:211:GLY:HA2	2.20	0.41
1:D:218:LEU:HD22	6:D:2033:HOH:O	2.12	0.40
1:A:138:TYR:O	1:A:216:GLY:HA2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/134 (100%)	127 (95%)	7 (5%)	0	100	100
1	D	134/134 (100%)	126 (94%)	8 (6%)	0	100	100
2	B	133/136 (98%)	124 (93%)	9 (7%)	0	100	100
2	E	133/136 (98%)	124 (93%)	9 (7%)	0	100	100
3	C	132/131 (101%)	127 (96%)	5 (4%)	0	100	100
3	F	132/131 (101%)	127 (96%)	5 (4%)	0	100	100
All	All	798/802 (100%)	755 (95%)	43 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/117 (103%)	119 (99%)	1 (1%)	89	96
1	D	120/117 (103%)	120 (100%)	0	100	100
2	B	116/117 (99%)	111 (96%)	5 (4%)	40	52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	116/117 (99%)	111 (96%)	5 (4%)	40	52
3	C	118/115 (103%)	118 (100%)	0	100	100
3	F	118/115 (103%)	118 (100%)	0	100	100
All	All	708/698 (101%)	697 (98%)	11 (2%)	75	88

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	158	ARG
2	B	92	THR
2	B	118	VAL
2	B	123	ASN
2	B	162	GLU
2	B	189	LEU
2	E	118	VAL
2	E	123	ASN
2	E	149	SER
2	E	189	LEU
2	E	194	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	125	HIS
1	A	172	ASN
1	A	186	GLN
3	C	102	GLN
3	C	146	HIS
1	D	172	ASN
1	D	186	GLN
1	D	189	GLN
3	F	101	HIS
3	F	102	GLN
3	F	146	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	IDU	F	1218	5	14,15,17	1.20	2 (14%)	14,22,26	1.37	2 (14%)
5	SGN	F	1219	5	20,20,20	1.43	2 (10%)	30,31,31	1.20	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	IDU	F	1218	5	1/1/5/7	0/9/22/29	0/1/1/1
5	SGN	F	1219	5	-	0/11/31/31	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	1219	SGN	O2S-S1	4.25	1.46	1.42
5	F	1219	SGN	O1S-S1	4.02	1.46	1.42
5	F	1218	IDU	O2-S	2.61	1.64	1.60
5	F	1218	IDU	C5-C6	-2.04	1.50	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	1218	IDU	O62-C6-C5	3.94	118.63	113.04
5	F	1219	SGN	C6-O6-S2	2.57	120.07	116.76
5	F	1219	SGN	O2S-S1-O1S	-2.49	112.07	119.84

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	1218	IDU	O61-C6-C5	-2.32	118.00	121.92
5	F	1219	SGN	C4-C3-C2	2.21	113.58	110.44
5	F	1219	SGN	O3S-S1-O1S	-2.03	111.61	115.28

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	F	1218	IDU	C1

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	1223	-	15,15,15	0.42	0	21,21,21	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1223	-	-	2/6/26/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1223	NAG	O7-C7-N2-C2
4	A	1223	NAG	C8-C7-N2-C2

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/134 (99%)	0.07	4 (3%)	48	58	4, 12, 26, 34	0
1	D	133/134 (99%)	0.01	4 (3%)	48	58	5, 13, 27, 32	0
2	B	133/136 (97%)	0.38	12 (9%)	10	16	8, 18, 32, 35	0
2	E	133/136 (97%)	0.41	13 (9%)	8	13	8, 18, 32, 35	2 (1%)
3	C	129/131 (98%)	-0.23	0	100	100	3, 9, 18, 25	0
3	F	129/131 (98%)	-0.31	0	100	100	3, 8, 17, 25	0
All	All	790/802 (98%)	0.06	33 (4%)	35	45	3, 13, 28, 35	2 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	224	MET	5.6
2	B	92	THR	4.4
2	E	201	ASP	4.2
1	D	104	MET	3.7
1	A	90	GLN	3.6
2	E	104	ASN	3.6
2	E	162	GLU	3.6
2	E	92	THR	3.5
2	B	224	MET	3.4
2	B	175	TYR	3.2
2	E	106	PRO	3.2
2	B	162	GLU	3.1
2	B	109	ARG	3.1
1	A	104	MET	3.0
2	E	175	TYR	2.8
1	D	92	ARG	2.7
1	A	91	PRO	2.7
2	B	103	ILE	2.6
2	E	109	ARG	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	163	ARG	2.6
2	B	129	ARG	2.6
2	B	201	ASP	2.6
2	E	129	ARG	2.6
1	A	92	ARG	2.5
1	D	101	ASN	2.5
2	B	105	VAL	2.3
2	B	106	PRO	2.2
2	E	176	ASN	2.2
1	D	90	GLN	2.1
2	E	223	ASP	2.1
2	B	223	ASP	2.0
2	E	209	GLU	2.0
2	E	93	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	IDU	F	1218	15/17	0.75	14.35	80,81,81,81	15
5	SGN	F	1219	20/20	0.39	9.93	80,81,81,81	6

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	1223	15/15	0.30	3.10	62,63,64,64	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.