



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 01:41 AM GMT

PDB ID : 2WPW
Title : TANDEM GNAT PROTEIN FROM THE CLAVULANIC ACID BIOSYN-
THESIS PATHWAY (WITHOUT ACCOA)
Authors : Iqbal, A.; Arunlanantham, H.; Mcdonough, M.A.; Chowdhury, R.; Clifton, I.J.
Deposited on : 2009-08-11
Resolution : 2.38 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2963 (2.40-2.36)
Clashscore	79885	3668 (2.40-2.36)
Ramachandran outliers	78287	3600 (2.40-2.36)
Sidechain outliers	78261	3602 (2.40-2.36)
RSRZ outliers	66119	2966 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	339	
1	B	339	
1	C	339	
1	D	339	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11259 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ORF14.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	328	Total	C	N	O	S	Se	0	0	0
			2486	1554	462	463	1	6			
1	B	326	Total	C	N	O	S	Se	0	0	0
			2474	1547	460	460	1	6			
1	C	325	Total	C	N	O	S	Se	0	0	0
			2470	1545	459	459	1	6			
1	D	325	Total	C	N	O	S	Se	0	0	0
			2470	1545	459	459	1	6			

- Molecule 2 is ACETYL COENZYME *A (three-letter code: ACO) (formula: C₂₃H₃₈N₇O₁₇P₃S).

Image for chem-comp ACO is not available.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		
2	C	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		
2	D	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	285	Total	O	0	0
			285	285		
3	B	277	Total	O	0	0
			277	277		
3	C	283	Total	O	0	0
			283	283		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	310	Total	O	0	0
			310	310		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ORF14

Chain A:




• Molecule 1: ORF14

Chain B:



• Molecule 1: ORF14

Chain C:



• Molecule 1: ORF14

[illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.45Å 93.50Å 126.43Å 90.00° 97.86° 90.00°	Depositor
Resolution (Å)	32.40 – 2.38 32.40 – 2.38	Depositor EDS
% Data completeness (in resolution range)	92.2 (32.40-2.38) 92.3 (32.40-2.38)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.209 , 0.266 0.214 , 0.265	Depositor DCC
R_{free} test set	3146 reflections (5.11%)	DCC
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 22.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	4 of 68369 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11259	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.35 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3381e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.88	1/2543 (0.0%)	0.89	6/3468 (0.2%)
1	B	0.85	0/2531	0.89	5/3451 (0.1%)
1	C	0.86	2/2527 (0.1%)	0.88	5/3446 (0.1%)
1	D	0.83	0/2527	0.88	2/3446 (0.1%)
All	All	0.86	3/10128 (0.0%)	0.88	18/13811 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	44	VAL	CB-CG2	6.42	1.66	1.52
1	C	64	VAL	CB-CG2	-5.29	1.41	1.52
1	C	63	VAL	CB-CG1	-5.06	1.42	1.52

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	118	ARG	NE-CZ-NH1	10.21	125.41	120.30
1	A	339	ASP	CB-CG-OD2	-8.50	110.65	118.30
1	A	118	ARG	NE-CZ-NH1	8.23	124.42	120.30
1	B	118	ARG	NE-CZ-NH1	7.50	124.05	120.30
1	B	150	ARG	NE-CZ-NH1	7.49	124.04	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens

added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2486	0	2420	89	0
1	B	2474	0	2408	90	0
1	C	2470	0	2405	77	0
1	D	2470	0	2405	100	0
2	A	51	0	34	5	0
2	B	51	0	34	5	0
2	C	51	0	34	4	1
2	D	51	0	34	8	1
3	A	285	0	0	30	0
3	B	277	0	0	21	0
3	C	283	0	0	31	0
3	D	310	0	0	51	0
All	All	11259	0	9774	373	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 19.

The worst 5 of 373 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:103:ARG:HA	1:B:146:MSE:CE	1.59	1.33
1:D:313:MSE:HG2	3:D:2048:HOH:O	1.30	1.24
1:A:132:GLY:HA2	3:A:2127:HOH:O	1.39	1.21
1:D:150:ARG:NH1	3:D:2154:HOH:O	1.76	1.17
2:A:401:ACO:C8A	2:A:401:ACO:H51A	1.76	1.15

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:C:401:ACO:O7A	2:D:401:ACO:O2B[2_555]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/339 (96%)	315 (97%)	8 (2%)	1 (0%)	50	66
1	B	322/339 (95%)	312 (97%)	8 (2%)	2 (1%)	33	46
1	C	321/339 (95%)	312 (97%)	8 (2%)	1 (0%)	50	66
1	D	321/339 (95%)	305 (95%)	13 (4%)	3 (1%)	25	33
All	All	1288/1356 (95%)	1244 (97%)	37 (3%)	7 (0%)	38	51

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201	GLU
1	B	213	ALA
1	D	201	GLU
1	B	130	PRO
1	D	131	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/249 (100%)	224 (90%)	24 (10%)	12	16
1	B	247/249 (99%)	226 (92%)	21 (8%)	15	21
1	C	247/249 (99%)	225 (91%)	22 (9%)	14	19
1	D	247/249 (99%)	231 (94%)	16 (6%)	24	34
All	All	989/996 (99%)	906 (92%)	83 (8%)	16	22

5 of 83 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	268	MSE
1	C	37	ARG
1	D	256	LYS
1	B	280	LEU
1	B	310	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	306	ASN
1	C	159	GLN
1	D	310	ASN
1	B	317	ASN
1	C	89	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACO	A	401	-	53,53,53	1.57	10 (18%)	79,79,79	4.19	29 (36%)
2	ACO	B	401	-	53,53,53	1.69	10 (18%)	79,79,79	4.14	29 (36%)
2	ACO	C	401	-	53,53,53	1.49	7 (13%)	79,79,79	4.24	27 (34%)
2	ACO	D	401	-	53,53,53	1.82	9 (16%)	79,79,79	4.06	27 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACO	A	401	-	-	0/51/67/67	0/1/3/3
2	ACO	B	401	-	-	0/51/67/67	0/1/3/3
2	ACO	C	401	-	-	0/51/67/67	0/1/3/3
2	ACO	D	401	-	-	0/51/67/67	0/1/3/3

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	401	ACO	O4B-C1B	6.15	1.50	1.41
2	B	401	ACO	O4B-C1B	5.50	1.49	1.41
2	D	401	ACO	P3B-O7A	5.35	1.69	1.51
2	B	401	ACO	CDP-CBP	-4.87	1.43	1.53
2	C	401	ACO	O4B-C1B	4.75	1.48	1.41

The worst 5 of 112 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	ACO	CEP-CBP-CCP	-17.76	83.17	108.76
2	D	401	ACO	CDP-CBP-CCP	-16.86	84.46	108.76
2	C	401	ACO	CEP-CBP-CCP	-16.74	84.63	108.76
2	B	401	ACO	CCP-CBP-CAP	-15.14	86.57	108.70
2	C	401	ACO	CDP-CBP-CCP	-15.07	87.04	108.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	328/339 (96%)	-0.15	3 (0%) 81 83	15, 26, 54, 86	0
1	B	326/339 (96%)	-0.14	8 (2%) 54 56	13, 27, 55, 76	0
1	C	325/339 (95%)	-0.17	3 (0%) 81 83	16, 27, 51, 76	0
1	D	325/339 (95%)	-0.16	3 (0%) 81 83	14, 27, 52, 73	0
All	All	1304/1356 (96%)	-0.15	17 (1%) 74 77	13, 27, 53, 86	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	6	GLY	4.8
1	A	5	ALA	4.0
1	A	4	THR	3.8
1	B	202	LEU	3.7
1	A	6	GLY	3.5

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ACO	C	401	51/51	0.12	-0.05	15,29,39,43	0
2	ACO	A	401	51/51	0.11	-0.12	13,26,37,43	0
2	ACO	D	401	51/51	0.12	-0.19	26,33,47,49	0
2	ACO	B	401	51/51	0.11	-0.41	17,28,40,44	0

6.5 Other polymers ⓘ

There are no such residues in this entry.