



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 12:35 AM GMT

PDB ID : 2WTK
Title : STRUCTURE OF THE HETEROTRIMERIC LKB1-STRADALPHA-MO25
ALPHACOMPLEX
Authors : Zeqiraj, E.; Van Aalten, D.M.F.
Deposited on : 2009-09-16
Resolution : 2.65 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2232 (2.70-2.62)
Clashscore	79885	2700 (2.70-2.62)
Ramachandran outliers	78287	2657 (2.70-2.62)
Sidechain outliers	78261	2657 (2.70-2.62)
RSRZ outliers	66119	2234 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	341	
1	D	341	
2	B	373	
2	E	373	
3	C	305	
3	F	305	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 14850 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CALCIUM-BINDING PROTEIN 39.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	321	Total	C	N	O	S	0	0	0
			2655	1708	448	488	11			
1	D	325	Total	C	N	O	S	0	0	0
			2681	1721	450	499	11			

- Molecule 2 is a protein called STE20-RELATED KINASE ADAPTER PROTEIN ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	311	Total	C	N	O	S	0	0	0
			2473	1588	422	448	15			
2	E	308	Total	C	N	O	S	0	0	0
			2454	1576	419	444	15			


- Molecule 3 is a protein called SERINE/THREONINE-PROTEINKINASE 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	285	Total	C	N	O	S	0	0	0
			2254	1448	386	406	14			
3	F	268	Total	C	N	O	S	0	0	0
			2110	1360	363	374	13			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	194	ALA	ASP	ENGINEERED MUTATION	UNP Q15831
F	194	ALA	ASP	ENGINEERED MUTATION	UNP Q15831

- Molecule 4 is PHOSPHOAMINOPHOSPHONICACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
4	C	1	Total	C	N	O	P	0	0
			27	10	6	9	2		
4	E	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
4	F	1	Total	C	N	O	P	0	0
			27	10	6	9	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	24	Total	O	0	0
			24	24		
6	B	8	Total	O	0	0
			8	8		
6	C	18	Total	O	0	0
			18	18		
6	D	26	Total	O	0	0
			26	26		
6	E	18	Total	O	0	0
			18	18		
6	F	8	Total	O	0	0
			8	8		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CALCIUM-BINDING PROTEIN 39

Chain A:




• Molecule 1: CALCIUM-BINDING PROTEIN 39

Chain D:



• Molecule 2: STE20-RELATED KINASE ADAPTER PROTEIN ALPHA

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.36Å 118.36Å 390.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.90 – 2.65 19.96 – 2.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.90-2.65) 99.2 (19.96-2.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.67Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.240 , 0.291 0.287 , 0.322	Depositor DCC
R_{free} test set	917 reflections (1.00%)	DCC
Wilson B-factor (Å ²)	60.4	Xtriage
Anisotropy	0.505	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 33.4	EDS
Estimated twinning fraction	0.015 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 91758 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	14850	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.54	0/2700	0.62	0/3627
1	D	0.56	0/2726	0.62	0/3664
2	B	0.55	0/2534	0.69	0/3432
2	E	0.57	0/2514	0.67	0/3404
3	C	0.48	0/2305	0.62	1/3110 (0.0%)
3	F	0.68	5/2153 (0.2%)	0.62	0/2899
All	All	0.56	5/14932 (0.0%)	0.64	1/20136 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	273	ALA	CA-CB	15.55	1.85	1.52
3	F	273	ALA	N-CA	8.80	1.64	1.46
3	F	273	ALA	C-N	8.24	1.52	1.34
3	F	273	ALA	C-O	7.15	1.36	1.23
3	F	257	GLY	CA-C	6.12	1.61	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	105	LEU	CA-CB-CG	5.13	127.10	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens

added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2655	0	2710	30	0
1	D	2681	0	2728	31	0
2	B	2473	0	2439	52	0
2	E	2454	0	2420	53	0
3	C	2254	0	2280	54	0
3	F	2110	0	2162	37	0
4	B	31	0	13	2	0
4	C	27	0	12	3	0
4	E	31	0	13	1	0
4	F	27	0	12	0	0
5	D	5	0	0	1	0
6	A	24	0	0	0	0
6	B	8	0	0	1	0
6	C	18	0	0	1	0
6	D	26	0	0	0	0
6	E	18	0	0	0	0
6	F	8	0	0	1	0
All	All	14850	0	14789	252	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

The worst 5 of 252 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:F:273:ALA:CB	3:F:273:ALA:CA	1.85	1.50
3:C:104:ARG:HH11	3:C:104:ARG:HG3	1.11	1.10
2:B:161:THR:HG21	2:B:416:GLY:HA3	1.07	1.05
2:B:384:ARG:HG2	2:B:384:ARG:HH11	1.26	1.01
2:B:161:THR:CG2	2:B:416:GLY:HA3	1.95	0.97

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/341 (92%)	300 (96%)	11 (4%)	2 (1%)	33	63
1	D	319/341 (94%)	301 (94%)	16 (5%)	2 (1%)	33	63
2	B	301/373 (81%)	277 (92%)	20 (7%)	4 (1%)	18	39
2	E	296/373 (79%)	270 (91%)	22 (7%)	4 (1%)	16	37
3	C	275/305 (90%)	258 (94%)	15 (6%)	2 (1%)	30	59
3	F	254/305 (83%)	235 (92%)	15 (6%)	4 (2%)	14	32
All	All	1758/2038 (86%)	1641 (93%)	99 (6%)	18 (1%)	22	48

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	222	ASN
2	B	62	PHE
1	D	317	GLU
2	E	387	SER
3	F	214	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/312 (95%)	278 (94%)	18 (6%)	26	52
1	D	299/312 (96%)	277 (93%)	22 (7%)	20	40
2	B	274/331 (83%)	245 (89%)	29 (11%)	10	20
2	E	273/331 (82%)	243 (89%)	30 (11%)	9	19
3	C	246/262 (94%)	225 (92%)	21 (8%)	15	31
3	F	230/262 (88%)	211 (92%)	19 (8%)	16	33
All	All	1618/1810 (89%)	1479 (91%)	139 (9%)	15	31

5 of 139 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	245	LEU
1	D	132	LYS
3	F	115	ASP
3	C	278	CYS
1	D	58	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	63	GLN
1	D	228	GLN
1	D	100	GLN
2	B	254	GLN
1	D	205	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ANP	B	432	-	33,33,33	3.12	6 (18%)	51,52,52	2.60	13 (25%)
4	ANP	C	2	-	28,29,33	1.27	3 (10%)	41,45,52	1.99	10 (24%)
5	SO4	D	342	-	4,4,4	0.10	0	6,6,6	0.17	0
4	ANP	E	432	-	33,33,33	3.23	6 (18%)	51,52,52	2.14	10 (19%)
4	ANP	F	4	-	28,29,33	1.32	3 (10%)	41,45,52	1.87	7 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	B	432	-	-	0/18/38/38	0/1/3/3
4	ANP	C	2	-	-	0/15/32/38	0/1/3/3
5	SO4	D	342	-	-	0/0/0/0	0/0/0/0
4	ANP	E	432	-	-	1/18/38/38	0/1/3/3
4	ANP	F	4	-	-	0/15/32/38	0/1/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	432	ANP	PG-N3B	12.09	1.74	1.64
4	E	432	ANP	PG-N3B	11.93	1.74	1.64
4	E	432	ANP	PB-N3B	10.98	1.73	1.64
4	B	432	ANP	PB-N3B	9.66	1.72	1.64
4	E	432	ANP	PG-O1G	4.91	1.52	1.46

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	432	ANP	PB-N3B-PG	-12.59	108.89	130.07
4	E	432	ANP	PB-N3B-PG	-7.48	117.50	130.07
4	E	432	ANP	N3-C2-N1	-7.15	122.73	128.71
4	B	432	ANP	N3-C2-N1	-6.66	123.14	128.71
4	C	2	ANP	N3-C2-N1	-5.97	123.71	128.71

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	432	ANP	O1B-PB-N3B-PG

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/341 (94%)	0.14	8 (2%) 54 58	28, 57, 107, 145	0
1	D	325/341 (95%)	0.15	13 (4%) 36 39	30, 61, 105, 155	0
2	B	311/373 (83%)	0.67	19 (6%) 21 21	51, 72, 106, 139	0
2	E	308/373 (82%)	0.56	25 (8%) 12 12	51, 74, 114, 146	0
3	C	285/305 (93%)	0.42	20 (7%) 16 16	39, 75, 128, 194	0
3	F	268/305 (87%)	0.89	42 (15%) 3 3	47, 95, 137, 171	0
All	All	1818/2038 (89%)	0.46	127 (6%) 16 16	28, 71, 121, 194	0

The worst 5 of 127 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	348	PHE	5.0
3	F	206	ALA	4.9
2	E	223	HIS	4.6
3	C	332	TRP	4.5
3	F	294	PRO	4.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	SO4	D	342	5/5	0.24	1.31	69,75,78,79	0
4	ANP	F	4	27/31	0.23	-0.16	89,98,102,103	0
4	ANP	C	2	27/31	0.17	-0.46	82,102,121,123	0
4	ANP	B	432	31/31	0.18	-0.55	44,56,69,73	0
4	ANP	E	432	31/31	0.17	-0.56	48,59,81,83	0

6.5 Other polymers

There are no such residues in this entry.