



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 06:40 PM GMT

PDB ID : 2WUI  
Title : CRYSTAL STRUCTURE OF MEXZ, A KEY REPRESSOR RESPONSIBLE  
FOR ANTIBIOTIC RESISTANCE IN PSEUDOMONAS AERUGINOSA.  
Authors : Alguel, Y.; Lu, D.; Quade, N.; Zhang, X.  
Deposited on : 2009-10-05  
Resolution : 2.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	210	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1539 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTIONAL REGULATOR.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	201	1539	960	288	281	10	0	0	1

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: TRANSCRIPTIONAL REGULATOR

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	177.06Å 177.06Å 54.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	31.38 – 2.90 39.76 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (31.38-2.90) 99.6 (39.76-2.90)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.49 (at 2.90Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.227 , 0.277 0.231 , 0.275	Depositor DCC
$R_{free}$ test set	345 reflections (4.72%)	DCC
Wilson B-factor (Å <sup>2</sup> )	74.5	Xtriage
Anisotropy	0.608	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 134.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 7321 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	1539	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	138.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/1562	0.86	7/2111 (0.3%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	33	ALA	CB-CA-C	-14.29	88.66	110.10
1	A	99	PRO	CB-CA-C	-12.66	80.36	112.00
1	A	94	ARG	NE-CZ-NH1	-7.83	116.39	120.30
1	A	98	GLU	CB-CA-C	-6.19	98.03	110.40
1	A	26	GLU	CB-CA-C	-6.01	98.37	110.40
1	A	94	ARG	NE-CZ-NH2	5.41	123.00	120.30
1	A	26	GLU	N-CA-C	5.29	125.27	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1539	0	1519	122	0
All	All	1539	0	1519	122	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 40.

All (122) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:138:LEU:HA	1:A:141:ILE:HD11	1.18	1.08
1:A:147:ARG:HG3	1:A:147:ARG:HH11	1.16	1.02
1:A:183:ASP:O	1:A:187:ARG:HD3	1.61	0.99
1:A:94:ARG:NH1	1:A:185:TRP:CD2	2.36	0.93
1:A:138:LEU:CA	1:A:141:ILE:HD11	2.01	0.90
1:A:93:LEU:HD23	1:A:93:LEU:O	1.73	0.88
1:A:102:VAL:O	1:A:106:LEU:HD22	1.74	0.87
1:A:26:GLU:HG2	1:A:27:LYS:HD2	1.57	0.87
1:A:38:ALA:HB2	1:A:48:VAL:HG21	1.57	0.86
1:A:25:LEU:HA	1:A:108:ILE:HD11	1.57	0.84
1:A:28:GLY:O	1:A:32:THR:HG22	1.78	0.83
1:A:147:ARG:CG	1:A:147:ARG:HH11	1.91	0.83
1:A:25:LEU:CA	1:A:108:ILE:HD11	2.09	0.82
1:A:16:ILE:HG21	1:A:48:VAL:HG13	1.62	0.81
1:A:91:GLY:O	1:A:94:ARG:HB3	1.83	0.78
1:A:10:GLN:O	1:A:13:ARG:HB2	1.87	0.75
1:A:94:ARG:NH1	1:A:185:TRP:CE3	2.54	0.74
1:A:207:LEU:HD12	1:A:207:LEU:H	1.51	0.74
1:A:52:TYR:CD2	1:A:58:VAL:HG23	2.24	0.71
1:A:15:GLY:O	1:A:19:ALA:HB2	1.92	0.69
1:A:153:ARG:O	1:A:207:LEU:HD12	1.93	0.69
1:A:147:ARG:HG3	1:A:147:ARG:NH1	1.96	0.69
1:A:19:ALA:O	1:A:23:VAL:HG23	1.93	0.69
1:A:125:ARG:O	1:A:128:LEU:HB3	1.93	0.68
1:A:10:GLN:HB2	1:A:13:ARG:HD2	1.75	0.68
1:A:93:LEU:HD23	1:A:93:LEU:C	2.14	0.67
1:A:149:GLU:O	1:A:150:LEU:HD23	1.95	0.66
1:A:90:MET:HE3	1:A:189:GLU:N	2.09	0.66
1:A:138:LEU:HA	1:A:141:ILE:CD1	2.12	0.66
1:A:21:GLU:OE1	1:A:65:ARG:NH2	2.29	0.65
1:A:43:VAL:CG1	1:A:48:VAL:HG22	2.27	0.65
1:A:207:LEU:O	1:A:209:ASP:N	2.29	0.64
1:A:170:ILE:HD11	1:A:192:PHE:CZ	2.33	0.63
1:A:94:ARG:NH1	1:A:185:TRP:CE2	2.67	0.62
1:A:13:ARG:HA	1:A:16:ILE:HD11	1.83	0.61
1:A:16:ILE:O	1:A:19:ALA:HB3	2.01	0.61
1:A:25:LEU:N	1:A:108:ILE:HD11	2.17	0.59
1:A:25:LEU:HA	1:A:108:ILE:CD1	2.29	0.59
1:A:24:PHE:CD1	1:A:32:THR:HG21	2.37	0.59
1:A:55:LYS:HG3	1:A:56:ILE:N	2.18	0.57
1:A:141:ILE:HD12	1:A:142:ARG:H	1.71	0.56
1:A:181:ARG:HG2	1:A:181:ARG:O	2.05	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:90:MET:HE2	1:A:192:PHE:HD2	1.71	0.55
1:A:93:LEU:CD2	1:A:93:LEU:C	2.75	0.55
1:A:98:GLU:CB	1:A:99:PRO:CD	2.84	0.55
1:A:152:ALA:O	1:A:153:ARG:HB2	2.08	0.54
1:A:90:MET:CE	1:A:192:PHE:CD2	2.90	0.54
1:A:29:VAL:HA	1:A:32:THR:CG2	2.38	0.54
1:A:153:ARG:O	1:A:206:LEU:HD12	2.08	0.53
1:A:16:ILE:O	1:A:20:ALA:N	2.42	0.53
1:A:90:MET:HE2	1:A:192:PHE:CD2	2.44	0.53
1:A:169:GLY:O	1:A:173:THR:HG23	2.09	0.53
1:A:11:LYS:HD3	1:A:12:THR:N	2.24	0.53
1:A:17:LEU:O	1:A:21:GLU:HG3	2.10	0.53
1:A:94:ARG:CA	1:A:94:ARG:HE	2.23	0.52
1:A:145:VAL:HG22	1:A:146:GLU:N	2.25	0.52
1:A:155:ASP:OD1	1:A:158:LEU:HB2	2.09	0.52
1:A:120:GLU:HB3	1:A:121:PRO:HD3	1.92	0.52
1:A:94:ARG:HA	1:A:94:ARG:HE	1.74	0.52
1:A:120:GLU:N	1:A:121:PRO:CD	2.72	0.51
1:A:43:VAL:HG13	1:A:44:SER:N	2.26	0.51
1:A:38:ALA:HB2	1:A:48:VAL:CG2	2.33	0.51
1:A:55:LYS:O	1:A:58:VAL:HG12	2.10	0.51
1:A:138:LEU:O	1:A:141:ILE:HD12	2.11	0.51
1:A:90:MET:HE1	1:A:192:PHE:CE2	2.46	0.50
1:A:90:MET:CE	1:A:192:PHE:HD2	2.22	0.50
1:A:110:TYR:HE1	1:A:126:ARG:HG3	1.76	0.50
1:A:173:THR:HA	1:A:177:THR:HG23	1.93	0.50
1:A:25:LEU:HD12	1:A:104:ARG:HG2	1.93	0.50
1:A:23:VAL:HG21	1:A:37:LEU:HD12	1.94	0.50
1:A:98:GLU:CB	1:A:99:PRO:HD3	2.43	0.49
1:A:13:ARG:O	1:A:16:ILE:HG12	2.13	0.49
1:A:147:ARG:NH1	1:A:147:ARG:CG	2.60	0.49
1:A:21:GLU:O	1:A:25:LEU:HB2	2.14	0.48
1:A:167:TRP:HA	1:A:170:ILE:CD1	2.44	0.48
1:A:178:GLU:OE2	1:A:178:GLU:HA	2.13	0.48
1:A:11:LYS:HB2	1:A:11:LYS:HE2	1.58	0.48
1:A:138:LEU:O	1:A:142:ARG:HG3	2.14	0.47
1:A:180:LEU:O	1:A:184:PRO:HB3	2.15	0.47
1:A:16:ILE:HG12	1:A:17:LEU:HD23	1.96	0.47
1:A:10:GLN:CD	1:A:10:GLN:C	2.73	0.46
1:A:17:LEU:HD13	1:A:62:MET:HA	1.98	0.46
1:A:43:VAL:HG13	1:A:44:SER:O	2.16	0.46
1:A:54:ASN:N	1:A:54:ASN:ND2	2.63	0.46

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:167:TRP:HA	1:A:170:ILE:HD11	1.96	0.46
1:A:158:LEU:HD23	1:A:158:LEU:HA	1.67	0.45
1:A:12:THR:O	1:A:16:ILE:HD13	2.16	0.45
1:A:166:LEU:HD23	1:A:166:LEU:O	2.17	0.45
1:A:103:GLN:HG3	1:A:104:ARG:N	2.31	0.45
1:A:49:TYR:O	1:A:52:TYR:O	2.35	0.44
1:A:202:SER:HA	1:A:203:PRO:HD2	1.57	0.44
1:A:172:GLY:HA2	1:A:175:ALA:O	2.17	0.44
1:A:175:ALA:HA	1:A:176:TRP:HA	1.74	0.44
1:A:166:LEU:O	1:A:170:ILE:HD13	2.17	0.44
1:A:160:SER:O	1:A:161:ILE:C	2.54	0.44
1:A:138:LEU:O	1:A:141:ILE:CD1	2.66	0.43
1:A:18:ASP:OD2	1:A:19:ALA:N	2.52	0.43
1:A:90:MET:HE1	1:A:192:PHE:CD2	2.53	0.43
1:A:64:ASP:O	1:A:65:ARG:C	2.57	0.43
1:A:150:LEU:HB3	1:A:151:PRO:HD2	2.00	0.43
1:A:21:GLU:OE2	1:A:104:ARG:NH1	2.47	0.43
1:A:145:VAL:CG2	1:A:146:GLU:N	2.81	0.43
1:A:43:VAL:HG11	1:A:48:VAL:HG22	1.98	0.43
1:A:17:LEU:N	1:A:17:LEU:HD23	2.34	0.42
1:A:137:GLY:O	1:A:141:ILE:HG13	2.18	0.42
1:A:90:MET:HE2	1:A:189:GLU:HA	2.02	0.42
1:A:29:VAL:HA	1:A:32:THR:HG22	2.01	0.42
1:A:105:VAL:O	1:A:108:ILE:HB	2.19	0.42
1:A:159:ALA:O	1:A:162:TYR:HB3	2.19	0.42
1:A:143:ARG:O	1:A:147:ARG:N	2.49	0.41
1:A:152:ALA:O	1:A:153:ARG:CB	2.67	0.41
1:A:182:ASP:O	1:A:183:ASP:C	2.58	0.41
1:A:16:ILE:HA	1:A:19:ALA:HB3	2.03	0.41
1:A:37:LEU:HD21	1:A:58:VAL:HG11	2.03	0.41
1:A:153:ARG:HH12	1:A:208:ALA:HB3	1.84	0.41
1:A:20:ALA:O	1:A:21:GLU:C	2.59	0.41
1:A:110:TYR:CE1	1:A:126:ARG:HD2	2.56	0.41
1:A:29:VAL:HG12	1:A:112:LYS:O	2.21	0.40
1:A:123:LEU:O	1:A:127:GLU:HG3	2.21	0.40
1:A:75:GLU:OE1	1:A:139:ARG:NH2	2.54	0.40
1:A:156:VAL:O	1:A:157:GLU:C	2.57	0.40
1:A:72:VAL:HA	1:A:73:PRO:HD3	1.97	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	199/210 (95%)	171 (86%)	24 (12%)	4 (2%)	11 40

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	208	ALA
1	A	99	PRO
1	A	184	PRO
1	A	28	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	153/172 (89%)	110 (72%)	43 (28%)	0 2

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	GLN
1	A	11	LYS
1	A	16	ILE
1	A	17	LEU
1	A	22	ARG
1	A	25	LEU
1	A	27	LYS
1	A	29	VAL
1	A	32	THR
1	A	37	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	43	VAL
1	A	48	VAL
1	A	54	ASN
1	A	58	VAL
1	A	70	ILE
1	A	86	LEU
1	A	94	ARG
1	A	97	CYS
1	A	103	GLN
1	A	106	LEU
1	A	108	ILE
1	A	109	LEU
1	A	122	LEU
1	A	129	LEU
1	A	140	GLN
1	A	141	ILE
1	A	147	ARG
1	A	158	LEU
1	A	163	LEU
1	A	164	GLN
1	A	170	ILE
1	A	171	CYS
1	A	176	TRP
1	A	177	THR
1	A	178	GLU
1	A	179	ARG
1	A	181	ARG
1	A	182	ASP
1	A	197	ASP
1	A	202	SER
1	A	205	LEU
1	A	206	LEU
1	A	207	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	ASN
1	A	69	GLN
1	A	76	ASN
1	A	103	GLN
1	A	140	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

### 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	201/210 (95%)	0.19	4 (1%) 62 71	112, 138, 158, 176	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	76	ASN	3.0
1	A	99	PRO	2.3
1	A	12	THR	2.3
1	A	35	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

### 6.5 Other polymers ⓘ

There are no such residues in this entry.