



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 06:45 PM GMT

PDB ID : 2X9M
Title : HENDRA VIRUS ATTACHMENT GLYCOPROTEIN
Authors : Bowden, T.A.; Crispin, M.; Harvey, D.; Jones, E.Y.; Stuart, D.I.
Deposited on : 2010-03-23
Resolution : 2.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	420	
1	B	420	
1	C	420	
1	D	420	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NAG	A	1607	-	X
2	NAG	B	1607	-	X

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 13464 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYCOPROTEIN G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	S	0	0	0
			3311	2109	556	627	19			
1	B	418	Total	C	N	O	S	0	0	0
			3304	2104	555	626	19			
1	C	417	Total	C	N	O	S	0	0	0
			3296	2100	553	624	19			
1	D	417	Total	C	N	O	S	0	0	0
			3296	2100	553	624	19			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	30	Total	O	0	0
			30	30		
3	B	30	Total	O	0	0
			30	30		
3	C	34	Total	O	0	0
			34	34		
3	D	23	Total	O	0	0
			23	23		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GLYCOPROTEIN G

Chain A: 




- Molecule 1: GLYCOPROTEIN G

Chain B: 




- Molecule 1: GLYCOPROTEIN G

Chain C: 



- Molecule 1: GLYCOPROTEIN G

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	65.88Å 84.78Å 91.55Å 103.39° 99.63° 108.62°	Depositor
Resolution (Å)	44.20 – 2.90 43.57 – 2.90	Depositor EDS
% Data completeness (in resolution range)	98.9 (44.20-2.90) 97.4 (43.57-2.90)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.233 , 0.273 0.236 , 0.272	Depositor DCC
R_{free} test set	1891 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	24.0	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , -6.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 38059 reflections	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	13464	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.53 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.8148e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/3392	0.59	1/4619 (0.0%)
1	B	0.41	0/3384	0.57	0/4608
1	C	0.42	0/3376	0.59	0/4597
1	D	0.44	2/3376 (0.1%)	0.58	0/4597
All	All	0.42	2/13528 (0.0%)	0.58	1/18421 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	446	ASN	CG-ND2	-6.23	1.17	1.32
1	D	446	ASN	CG-OD1	-5.81	1.11	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	571	VAL	CG1-CB-CG2	8.18	123.99	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3311	0	0	6	0
1	B	3304	0	10	10	2
1	C	3296	0	0	8	1
1	D	3296	0	0	7	1
2	A	56	0	52	1	0
2	B	56	0	52	0	0
2	C	14	0	13	0	0
2	D	14	0	13	0	0
3	A	30	0	0	1	0
3	B	30	0	0	3	0
3	C	34	0	0	1	0
3	D	23	0	0	0	0
All	All	13464	0	140	30	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (30) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:386:HIS:N	3:A:2014:HOH:O	2.32	0.63
1:C:288:GLU:N	1:C:288:GLU:OE1	2.33	0.61
1:A:443:LYS:NZ	1:A:510:ASP:OD2	2.35	0.59
1:A:415:LYS:CE	2:A:1604:NAG:H81	2.36	0.56
1:B:212:ARG:CZ	1:B:246:LYS:NZ	2.71	0.54
1:B:443:LYS:NZ	1:B:510:ASP:OD2	2.42	0.53
1:C:288:GLU:OE1	1:C:288:GLU:CA	2.58	0.52
1:D:443:LYS:NZ	1:D:563:THR:O	2.43	0.52
1:C:443:LYS:NZ	1:C:563:THR:O	2.45	0.49
1:B:201:ARG:NH1	3:B:2001:HOH:O	2.46	0.49
1:B:443:LYS:NZ	1:B:563:THR:O	2.46	0.49
1:D:443:LYS:NZ	1:D:510:ASP:OD2	2.47	0.47
1:A:443:LYS:NZ	1:A:563:THR:O	2.48	0.47
1:C:443:LYS:NZ	1:C:510:ASP:OD2	2.48	0.47
1:A:582:ASP:OD2	1:D:357:LYS:NZ	2.49	0.46
1:B:293:THR:CG2	1:B:317:ILE:O	2.64	0.45
1:B:557:ASN:ND2	1:B:582:ASP:OD2	2.48	0.45
1:B:248:ARG:NE	3:B:2006:HOH:O	2.51	0.44
1:B:212:ARG:NH2	1:B:246:LYS:NZ	2.66	0.43
1:D:583:THR:O	1:D:584:GLY:C	2.56	0.43
1:C:205:TYR:CE1	1:C:207:LEU:CD1	3.01	0.43
1:D:205:TYR:CE1	1:D:207:LEU:CD1	3.01	0.43
1:B:205:TYR:CE1	1:B:207:LEU:CD1	3.02	0.43
1:A:293:THR:CG2	1:A:317:ILE:O	2.67	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:581:TYR:N	1:D:581:TYR:CD2	2.87	0.42
1:C:424:ILE:CG2	1:C:424:ILE:O	2.68	0.41
1:C:246:LYS:NZ	3:C:2006:HOH:O	2.53	0.41
1:D:216:CYS:SG	1:D:218:THR:CG2	3.09	0.41
1:C:286:TYR:CE2	1:C:288:GLU:OE1	2.74	0.40
1:B:425:ILE:CG2	3:B:2021:HOH:O	2.69	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:341:GLU:OE2	1:D:373:THR:OG1[1_556]	1.92	0.28
1:B:582:ASP:OD2	1:C:288:GLU:OE2[1_565]	2.13	0.07

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/420 (99%)	395 (95%)	21 (5%)	1 (0%)	56	89
1	B	416/420 (99%)	393 (94%)	23 (6%)	0	100	100
1	C	415/420 (99%)	392 (94%)	23 (6%)	0	100	100
1	D	415/420 (99%)	386 (93%)	28 (7%)	1 (0%)	56	89
All	All	1663/1680 (99%)	1566 (94%)	95 (6%)	2 (0%)	59	91

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	569	GLU
1	D	288	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/377 (100%)	361 (96%)	15 (4%)	42	81
1	B	375/377 (100%)	361 (96%)	14 (4%)	45	84
1	C	374/377 (99%)	360 (96%)	14 (4%)	45	84
1	D	374/377 (99%)	358 (96%)	16 (4%)	40	78
All	All	1499/1508 (99%)	1440 (96%)	59 (4%)	43	82

All (59) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	202	LEU
1	A	207	LEU
1	A	212	ARG
1	A	215	VAL
1	A	221	LEU
1	A	291	TYR
1	A	316	LEU
1	A	324	LYS
1	A	351	TYR
1	A	376	GLN
1	A	423	ASP
1	A	487	ARG
1	A	519	TRP
1	A	571	VAL
1	A	582	ASP
1	B	202	LEU
1	B	210	ASN
1	B	215	VAL
1	B	221	LEU
1	B	241	THR
1	B	288	GLU
1	B	291	TYR
1	B	316	LEU
1	B	324	LYS
1	B	351	TYR
1	B	376	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	487	ARG
1	B	519	TRP
1	B	582	ASP
1	C	202	LEU
1	C	215	VAL
1	C	221	LEU
1	C	233	HIS
1	C	288	GLU
1	C	291	TYR
1	C	293	THR
1	C	316	LEU
1	C	324	LYS
1	C	351	TYR
1	C	376	GLN
1	C	487	ARG
1	C	519	TRP
1	C	582	ASP
1	D	202	LEU
1	D	215	VAL
1	D	221	LEU
1	D	233	HIS
1	D	242	ARG
1	D	291	TYR
1	D	293	THR
1	D	316	LEU
1	D	324	LYS
1	D	351	TYR
1	D	376	GLN
1	D	423	ASP
1	D	424	ILE
1	D	487	ARG
1	D	519	TRP
1	D	569	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	1604	1	12,14,15	0.78	1 (8%)	15,19,21	1.27	1 (6%)
2	NAG	A	1605	1	12,14,15	0.68	0	15,19,21	0.97	1 (6%)
2	NAG	A	1606	1	12,14,15	0.73	0	15,19,21	1.10	1 (6%)
2	NAG	A	1607	1	12,14,15	0.55	0	15,19,21	1.69	3 (20%)
2	NAG	B	1604	1	12,14,15	0.50	0	15,19,21	1.51	2 (13%)
2	NAG	B	1605	1	12,14,15	0.72	0	15,19,21	1.13	2 (13%)
2	NAG	B	1606	1	12,14,15	0.53	0	15,19,21	1.18	2 (13%)
2	NAG	B	1607	1	12,14,15	0.69	0	15,19,21	0.82	1 (6%)
2	NAG	C	1604	1	12,14,15	0.55	0	15,19,21	1.39	2 (13%)
2	NAG	D	1604	1	12,14,15	0.60	0	15,19,21	1.21	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1604	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1605	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1606	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1607	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1604	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1605	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1606	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1607	1	-	0/6/23/26	0/1/1/1
2	NAG	C	1604	1	-	0/6/23/26	0/1/1/1
2	NAG	D	1604	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1604	NAG	O5-C5	-2.42	1.40	1.45

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1604	NAG	O5-C5-C6	4.19	111.38	106.98
2	A	1607	NAG	O5-C5-C6	4.17	111.36	106.98
2	C	1604	NAG	O5-C5-C4	3.63	115.26	110.65
2	B	1604	NAG	C3-C2-N2	-3.48	106.47	111.76
2	D	1604	NAG	C3-C2-N2	-3.44	106.52	111.76
2	B	1606	NAG	O5-C5-C6	3.04	110.17	106.98
2	A	1607	NAG	C3-C2-N2	-2.87	107.40	111.76
2	B	1605	NAG	O5-C5-C6	2.86	109.98	106.98
2	A	1604	NAG	O4-C4-C5	-2.61	102.40	109.28
2	A	1606	NAG	O3-C3-C4	2.51	115.98	110.35
2	A	1605	NAG	O5-C5-C6	2.44	109.55	106.98
2	C	1604	NAG	C3-C4-C5	2.44	114.56	110.20
2	A	1607	NAG	O5-C5-C4	2.42	113.72	110.65
2	B	1607	NAG	C3-C2-N2	-2.34	108.20	111.76
2	B	1605	NAG	C3-C2-N2	-2.23	108.36	111.76
2	B	1606	NAG	C3-C2-N2	-2.17	108.46	111.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/420 (99%)	-0.14	8 (1%) 64 72	7, 25, 49, 91	0
1	B	418/420 (99%)	-0.15	5 (1%) 75 83	8, 26, 52, 90	0
1	C	417/420 (99%)	-0.06	9 (2%) 59 67	8, 27, 64, 94	0
1	D	417/420 (99%)	-0.07	8 (1%) 64 72	8, 27, 55, 85	0
All	All	1671/1680 (99%)	-0.10	30 (1%) 65 74	7, 26, 55, 94	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	583	THR	4.6
1	A	420	LEU	4.1
1	C	422	GLY	3.3
1	B	420	LEU	3.2
1	C	583	THR	3.1
1	C	194	THR	2.9
1	D	554	ASP	2.9
1	D	600	GLN	2.9
1	B	553	GLU	2.9
1	D	422	GLY	2.8
1	D	601	CYS	2.7
1	C	491	SER	2.7
1	C	581	TYR	2.7
1	D	196	THR	2.5
1	A	518	ASN	2.4
1	A	376	GLN	2.2
1	B	385	ILE	2.2
1	C	385	ILE	2.2
1	C	600	GLN	2.1
1	C	584	GLY	2.1
1	B	196	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	581	TYR	2.1
1	D	581	TYR	2.1
1	A	191	GLN	2.1
1	B	193	THR	2.1
1	A	553	GLU	2.1
1	A	185	PRO	2.0
1	D	191	GLN	2.0
1	A	602	SER	2.0
1	C	196	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q < 0.9
2	NAG	B	1607	14/15	0.23	2.73	17,20,22,23	0
2	NAG	A	1607	14/15	0.29	2.17	19,20,23,24	0
2	NAG	B	1606	14/15	0.20	1.97	20,23,24,25	0
2	NAG	A	1606	14/15	0.20	1.54	18,20,21,22	0
2	NAG	D	1604	14/15	0.24	0.84	14,18,19,20	0
2	NAG	C	1604	14/15	0.23	0.74	16,18,20,21	0
2	NAG	A	1604	14/15	0.15	-0.44	3,8,10,10	0
2	NAG	B	1605	14/15	0.15	-0.54	7,14,15,17	0
2	NAG	A	1605	14/15	0.14	-0.97	6,9,11,12	0
2	NAG	B	1604	14/15	0.14	-1.68	9,13,14,14	0

6.5 Other polymers ⓘ

There are no such residues in this entry.