



# Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 11:43 PM GMT

PDB ID : 2X9P  
Title : X-RAY STRUCTURE OF THE SUBSTRATE-FREE CYTOCHROME P450  
PIMD -A POLYENE MACROLIDE ANTIBIOTIC PIMARICIN EPOXI-  
DASE  
Authors : Kells, P.M.; Ouellet, H.; Santos-Aberturas, J.; Aparicio, J.F.; Podust, L.M.  
Deposited on : 2010-03-23  
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	404	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	SO4	A	1399	-	X
3	SO4	A	1400	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3190 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PIMD PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	385	2955	1877	518	550	10	0	3	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	EXPRESSION TAG	UNP Q9EW92
A	-5	ARG	-	EXPRESSION TAG	UNP Q9EW92
A	-4	GLY	-	EXPRESSION TAG	UNP Q9EW92
A	-3	SER	-	EXPRESSION TAG	UNP Q9EW92
A	-2	HIS	-	EXPRESSION TAG	UNP Q9EW92
A	-1	HIS	-	EXPRESSION TAG	UNP Q9EW92
A	0	HIS	-	EXPRESSION TAG	UNP Q9EW92
A	1	HIS	-	EXPRESSION TAG	UNP Q9EW92
A	2	HIS	-	EXPRESSION TAG	UNP Q9EW92
A	3	HIS	-	EXPRESSION TAG	UNP Q9EW92
A	4	GLY	-	EXPRESSION TAG	UNP Q9EW92

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	177	Total 177	O 177	0	0

### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: PIMD PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.60Å 139.38Å 70.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	90.17 – 2.10 16.98 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (90.17-2.10) 99.9 (16.98-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.76 (at 2.11Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.186 , 0.239 0.185 , 0.234	Depositor DCC
$R_{free}$ test set	1725 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.1	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 36.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 34146 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3190	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	5/3021 (0.2%)	1.00	10/4119 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	256	GLN	CG-CD	-7.28	1.34	1.51
1	A	295	GLU	CG-CD	-5.74	1.43	1.51
1	A	157	VAL	CB-CG1	-5.65	1.41	1.52
1	A	146	THR	CB-CG2	-5.10	1.35	1.52
1	A	358	THR	CB-CG2	-5.01	1.35	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	95	ARG	NE-CZ-NH2	-9.71	115.44	120.30
1	A	319	ARG	NE-CZ-NH1	7.44	124.02	120.30
1	A	146	THR	CA-CB-CG2	7.43	122.81	112.40
1	A	256	GLN	CA-CB-CG	-7.34	97.25	113.40
1	A	257	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	A	95	ARG	NE-CZ-NH1	7.18	123.89	120.30
1	A	257	ARG	NE-CZ-NH1	6.19	123.40	120.30
1	A	319	ARG	NE-CZ-NH2	-6.08	117.26	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	50	ARG	NE-CZ-NH1	-5.32	117.64	120.30
1	A	95	ARG	CG-CD-NE	-5.09	101.12	111.80

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	146	THR	CB

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	5	SER	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2955	0	0	4	0
2	A	43	0	0	2	0
3	A	15	0	0	0	0
4	A	177	0	0	2	0
All	All	3190	0	0	5	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (5) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:354:LEU:O	1:A:358:THR:CG2	2.59	0.51
1:A:38:ARG:NH1	4:A:2020:HOH:O	2.44	0.49
1:A:102:ARG:NH1	2:A:1398:HEM:O1D	2.46	0.48
2:A:1398:HEM:O2D	4:A:2175:HOH:O	2.20	0.47
1:A:116:GLU:OE2	1:A:213:ARG:NH2	2.48	0.47

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/404 (95%)	375 (98%)	8 (2%)	1 (0%)	50	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	179	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/328 (92%)	284 (95%)	16 (5%)	32	28

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ARG
1	A	93	SER
1	A	115	LEU
1	A	141	LEU
1	A	146	THR
1	A	168	LEU
1	A	175	ILE
1	A	179	ASP
1	A	182	GLU
1	A	288	ARG
1	A	295	GLU
1	A	302	ARG
1	A	358	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	371	ARG
1	A	381	LEU
1	A	394	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	1398	1	49,50,50	2.45	14 (28%)	46,82,82	2.41	10 (21%)
3	SO4	A	1399	-	4,4,4	0.53	0	6,6,6	0.38	0
3	SO4	A	1400	-	4,4,4	0.37	0	6,6,6	0.50	0
3	SO4	A	1401	-	4,4,4	1.03	0	6,6,6	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1398	1	-	0/14/114/114	0/0/8/8
3	SO4	A	1399	-	-	0/0/0/0	0/0/0/0
3	SO4	A	1400	-	-	0/0/0/0	0/0/0/0
3	SO4	A	1401	-	-	0/0/0/0	0/0/0/0

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1398	HEM	C3B-C2B	-6.83	1.31	1.43
2	A	1398	HEM	C3D-C4D	6.62	1.46	1.44
2	A	1398	HEM	C3C-C2C	-6.42	1.32	1.43
2	A	1398	HEM	C3B-CAB	4.72	1.55	1.40
2	A	1398	HEM	C3C-CAC	4.50	1.54	1.40
2	A	1398	HEM	C3D-C2D	3.79	1.50	1.43
2	A	1398	HEM	C4A-C3A	3.64	1.44	1.40
2	A	1398	HEM	FE-NA	3.29	2.06	1.92
2	A	1398	HEM	CHB-C1B	3.07	1.40	1.35
2	A	1398	HEM	CMC-C2C	2.88	1.56	1.47
2	A	1398	HEM	C2D-C1D	2.60	1.45	1.44
2	A	1398	HEM	C4C-NC	-2.29	1.34	1.38
2	A	1398	HEM	C3B-C4B	2.16	1.47	1.44
2	A	1398	HEM	C2B-C1B	2.15	1.45	1.44

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1398	HEM	CBA-CAA-C2A	-9.25	96.40	112.69
2	A	1398	HEM	C3B-C4B-NB	-7.30	108.78	114.00
2	A	1398	HEM	CAD-C3D-C4D	4.29	132.25	124.53
2	A	1398	HEM	C4D-ND-C1D	3.70	108.94	105.16
2	A	1398	HEM	C4A-CHB-C1B	-3.61	122.72	127.47
2	A	1398	HEM	CMA-C3A-C4A	-3.55	123.15	128.62
2	A	1398	HEM	CHD-C1D-ND	3.19	127.24	124.58
2	A	1398	HEM	CMA-C3A-C2A	2.90	130.42	124.94
2	A	1398	HEM	C2D-C1D-ND	-2.67	109.78	112.93
2	A	1398	HEM	O2A-CGA-CBA	2.22	122.05	114.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	385/404 (95%)	-0.25	10 (2%) 53 58	16, 29, 55, 68	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	177	LYS	3.9
1	A	82	LEU	3.7
1	A	179	ASP	3.6
1	A	182	GLU	2.8
1	A	180	ASP	2.7
1	A	391	ALA	2.6
1	A	178	LEU	2.3
1	A	5	SER	2.2
1	A	85	LEU	2.2
1	A	376	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	1399	5/5	0.22	5.61	62,64,65,67	0
3	SO4	A	1400	5/5	0.19	2.07	63,63,64,64	0
2	HEM	A	1398	43/43	0.10	-0.01	11,18,35,42	0
3	SO4	A	1401	5/5	0.06	-1.17	24,24,27,27	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.