



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 04:16 AM GMT

PDB ID : 1XAD  
Title : CHIMERA ISOPROPYLMALATE DEHYDROGENASE BETWEEN  
BACILLUS SUBTILIS (M) AND THERMUS THERMOPHILUS (T) FROM  
N-TERMINAL: 20% T MIDDLE 20% M RESIDUAL 60% T, MUTATED AT  
S82R. LOW TEMPERATURE (150K) STRUCTURE.  
Authors : Nagata, C.; Moriyama, H.; Tanaka, N.  
Deposited on : 1995-11-09  
Resolution : 2.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : NOT EXECUTED  
EDS : NOT EXECUTED  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	345	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3818 atoms, of which 987 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-ISOPROPYLMALATE DEHYDROGENASE 2T2M6T S82R.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	345	Total	C	H	N	O	S	0	0	0
			3161	1665	549	450	491	6			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	GLN	GLY	CONFLICT	UNP P00351
A	80	ASN	LEU	CONFLICT	UNP P00351
A	83	GLU	LYS	CONFLICT	UNP P00351
A	84	LEU	ILE	CONFLICT	UNP P00351
A	85	ARG	SER	CONFLICT	UNP P00351
A	88	LYS	THR	CONFLICT	UNP P00351
A	93	ILE	LEU	CONFLICT	UNP P00351
A	96	GLN	SER	CONFLICT	UNP P00351
A	97	LEU	GLN	CONFLICT	UNP P00351
A	106	VAL	ALA	CONFLICT	UNP P00351
A	110	GLU	PRO	CONFLICT	UNP P00351
A	111	SER	GLY	CONFLICT	UNP P00351
A	113	SER	GLU	CONFLICT	UNP P00351
A	114	ASP	ARG	CONFLICT	UNP P00351
A	115	ALA	LEU	CONFLICT	UNP P00351
A	120	LYS	GLU	CONFLICT	UNP P00351
A	122	TYR	ILE	CONFLICT	UNP P00351
A	123	ILE	ALA	CONFLICT	UNP P00351
A	124	ASP	ARG	CONFLICT	UNP P00351
A	125	ASN	GLY	CONFLICT	UNP P00351
A	128	PHE	VAL	CONFLICT	UNP P00351
A	129	VAL	LEU	CONFLICT	UNP P00351

- Molecule 2 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	219	Total	H	O	0	0
			657	438	219		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 3-ISOPROPYLMALATE DEHYDROGENASE 2T2M6T S82R



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.40Å 77.40Å 156.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.10)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.194 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3818	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.12	1/2668 (0.0%)	1.90	65/3621 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	275	SER	CA-CB	-6.02	1.44	1.52

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	ARG	NE-CZ-NH1	12.70	126.65	120.30
1	A	204	ARG	NE-CZ-NH1	11.16	125.88	120.30
1	A	77	TRP	CG-CD2-CE3	8.99	141.99	133.90
1	A	85	ARG	NE-CZ-NH1	8.95	124.78	120.30
1	A	156	ARG	NE-CZ-NH2	-8.93	115.83	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	157	TYR	Sidechain
1	A	80	ASN	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2612	549	2081	73	0
2	A	219	438	0	24	0
All	All	2831	987	2081	73	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:94:ARG:HD2	1:A:134:LEU:HD13	1.58	0.85
1:A:30:GLU:HB2	1:A:32:LEU:HD12	1.64	0.80
1:A:325:PRO:HA	1:A:329:GLY:O	1.90	0.72
1:A:90:LEU:HD22	2:A:468:HOH:H2	1.51	0.70
1:A:83:GLU:HA	1:A:88:LYS:HG3	1.75	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	343/345 (99%)	312 (91%)	26 (8%)	5 (2%)	15   8

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	81	PRO

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	273	HIS
1	A	228	ALA
1	A	28	GLU
1	A	207	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/270 (100%)	242 (90%)	28 (10%)	10 6

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	152	TRP
1	A	176	ARG
1	A	324	PRO
1	A	157	TYR
1	A	165	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	222	HIS
1	A	300	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.