



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 04:39 AM GMT

PDB ID : 2XO4
Title : RIBONUCLEOTIDE REDUCTASE Y730NH2Y MODIFIED R1 SUBUNIT OF E. COLI
Authors : Minnihan, E.C.; Seyedsayamdost, M.R.; Uhlin, U.; Stubbe, J.
Deposited on : 2010-08-09
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	761	
1	B	761	
1	C	761	
2	D	20	
2	E	20	
2	F	20	
2	P	20	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18959 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEOSIDE-DIPHOSPHATEREDUCTASE 1 SUBUNIT ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	734	Total	C	N	O	S	0	0	0
			5846	3712	1005	1105	24			
1	B	734	Total	C	N	O	S	0	0	0
			5846	3712	1005	1105	24			
1	C	734	Total	C	N	O	S	0	0	0
			5846	3712	1005	1105	24			

- Molecule 2 is a protein called RIBONUCLEOSIDE-DIPHOSPHATEREDUCTASE 1 SUBUNIT BETA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	16	Total	C	N	O	0	0	0
			129	77	19	33			
2	E	16	Total	C	N	O	0	0	0
			129	77	19	33			
2	F	16	Total	C	N	O	0	0	0
			129	77	19	33			
2	P	3	Total	C	N	O	0	0	0
			27	20	3	4			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	287	Total	O	0	0
			287	287		
3	B	306	Total	O	0	0
			306	306		
3	C	400	Total	O	0	0
			400	400		
3	D	4	Total	O	0	0
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	3	Total 3	O 3	0	0
3	F	4	Total 4	O 4	0	0
3	P	3	Total 3	O 3	0	0




• Molecule 2: RIBONUCLEOSIDE-DIPHOSPHATEREDUCTASE 1 SUBUNIT BETA

Chain D:




• Molecule 2: RIBONUCLEOSIDE-DIPHOSPHATEREDUCTASE 1 SUBUNIT BETA

Chain E:




• Molecule 2: RIBONUCLEOSIDE-DIPHOSPHATEREDUCTASE 1 SUBUNIT BETA

Chain F:



• Molecule 2: RIBONUCLEOSIDE-DIPHOSPHATEREDUCTASE 1 SUBUNIT BETA

Chain P:



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	224.59Å 224.59Å 336.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	169.03 – 2.50 84.21 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (169.03-2.50) 98.7 (84.21-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.188 , 0.233 0.187 , 0.231	Depositor DCC
R_{free} test set	5539 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 27.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 110807 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18959	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TY2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/5959	0.57	0/8069
1	B	0.43	0/5959	0.56	0/8069
1	C	0.49	0/5959	0.60	2/8069 (0.0%)
2	D	0.42	0/129	0.56	0/173
2	E	0.41	0/129	0.61	0/173
2	F	0.44	0/129	0.61	0/173
2	P	0.58	0/27	0.63	0/36
All	All	0.45	0/18291	0.58	2/24762 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	216	ARG	NE-CZ-NH2	-5.91	117.35	120.30
1	C	639	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5846	0	5769	76	0
1	B	5846	0	5769	63	0
1	C	5846	0	5769	60	0
2	D	129	0	111	3	0
2	E	129	0	111	3	0
2	F	129	0	111	1	0
2	P	27	0	31	2	0
3	A	287	0	0	17	0
3	B	306	0	0	17	0
3	C	400	0	0	21	0
3	D	4	0	0	0	0
3	E	3	0	0	0	0
3	F	4	0	0	1	0
3	P	3	0	0	2	0
All	All	18959	0	17671	202	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

The worst 5 of 202 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:623:GLU:HG2	3:B:2088:HOH:O	1.53	1.07
1:A:10:ARG:HG3	1:A:10:ARG:HH11	1.28	0.98
1:C:639:ARG:HG3	1:C:639:ARG:HH11	1.34	0.92
1:A:294:GLN:HG3	1:A:295:GLY:N	1.85	0.90
1:B:450:ASN:HB2	3:B:2187:HOH:O	1.74	0.86

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	731/761 (96%)	698 (96%)	25 (3%)	8 (1%)	21 34
1	B	731/761 (96%)	699 (96%)	23 (3%)	9 (1%)	19 32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	731/761 (96%)	701 (96%)	24 (3%)	6 (1%)	27	46
2	D	14/20 (70%)	13 (93%)	1 (7%)	0	100	100
2	E	14/20 (70%)	11 (79%)	3 (21%)	0	100	100
2	F	14/20 (70%)	14 (100%)	0	0	100	100
2	P	1/20 (5%)	1 (100%)	0	0	100	100
All	All	2236/2363 (95%)	2137 (96%)	76 (3%)	23 (1%)	22	38

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	294	GLN
1	B	216	ARG
1	B	268	ILE
1	A	216	ARG
1	A	269	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	628/650 (97%)	586 (93%)	42 (7%)	23	40
1	B	628/650 (97%)	591 (94%)	37 (6%)	28	48
1	C	628/650 (97%)	586 (93%)	42 (7%)	23	40
2	D	16/19 (84%)	13 (81%)	3 (19%)	2	3
2	E	16/19 (84%)	13 (81%)	3 (19%)	2	3
2	F	16/19 (84%)	13 (81%)	3 (19%)	2	3
2	P	3/19 (16%)	3 (100%)	0	100	100
All	All	1935/2026 (96%)	1805 (93%)	130 (7%)	23	40

5 of 130 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	268	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	455	GLU
1	C	708	LYS
1	B	297	VAL
1	B	364	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	221	GLN
1	B	630	ASN
1	C	663	HIS
1	B	328	ASN
1	A	527	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TY2	A	730	1	13,13,14	5.90	5 (38%)	15,17,19	1.85	3 (20%)
1	TY2	B	730	1	13,13,14	5.94	5 (38%)	15,17,19	1.67	4 (26%)
1	TY2	C	730	1	13,13,14	5.62	5 (38%)	15,17,19	1.90	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TY2	A	730	1	-	0/4/6/8	0/1/1/1
1	TY2	B	730	1	-	0/4/6/8	0/1/1/1
1	TY2	C	730	1	-	0/4/6/8	0/1/1/1

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	730	TY2	O-C	18.68	1.24	1.11
1	A	730	TY2	O-C	18.55	1.24	1.11
1	C	730	TY2	O-C	17.53	1.23	1.11
1	C	730	TY2	CD2-CE2	-7.10	1.29	1.40
1	A	730	TY2	CD2-CE2	-7.04	1.29	1.40

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	730	TY2	C-CA-N	-5.46	108.38	113.83
1	A	730	TY2	C-CA-N	-5.22	108.61	113.83
1	B	730	TY2	C-CA-N	-3.82	110.01	113.83
1	C	730	TY2	CD1-CE1-CZ	-3.24	116.97	120.50
1	A	730	TY2	CD1-CE1-CZ	-2.93	117.31	120.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	734/761 (96%)	0.05	21 (2%) 49 51	22, 38, 64, 84	0
1	B	734/761 (96%)	0.06	17 (2%) 57 60	27, 38, 60, 82	0
1	C	734/761 (96%)	-0.08	11 (1%) 70 72	17, 29, 52, 75	0
2	D	16/20 (80%)	1.34	2 (12%) 5 4	71, 83, 90, 90	0
2	E	16/20 (80%)	1.03	1 (6%) 19 19	71, 82, 87, 87	0
2	F	16/20 (80%)	1.13	3 (18%) 2 2	62, 76, 81, 82	0
2	P	3/20 (15%)	0.67	0 100 100	29, 29, 35, 40	0
All	All	2253/2363 (95%)	0.04	55 (2%) 56 58	17, 36, 66, 90	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	296	GLY	8.0
1	A	273	ALA	7.2
1	B	296	GLY	6.6
1	A	274	PHE	5.9
1	C	271	GLY	5.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TY2	C	730	13/14	0.15	0.68	21,22,23,24	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	TY2	A	730	13/14	0.13	-0.55	30,31,32,32	0
1	TY2	B	730	13/14	0.13	-0.92	31,32,32,32	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.