



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 09:47 PM GMT

PDB ID : 1XS2
Title : Structural Basis for Catalytic Racemization and Substrate Specificity of an N-Acylamino Acid Racemase Homologue from *Deinococcus radiodurans*
Authors : Wang, W.-C.; Chiu, W.-C.; Hsu, S.-K.; Wu, C.-L.; Chen, C.-Y.; Liu, J.-S.; Hsu, W.-H.
Deposited on : 2004-10-18
Resolution : 2.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	375	
1	B	375	
1	C	375	
1	D	375	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11768 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-Acylamino Acid Racemase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	0	0
			2766	1733	507	516	10			
1	B	360	Total	C	N	O	S	0	0	0
			2766	1733	507	516	10			
1	C	360	Total	C	N	O	S	0	0	0
			2766	1733	507	516	10			
1	D	360	Total	C	N	O	S	0	0	0
			2766	1733	507	516	10			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	SER	ALA	SEE REMARK 999	UNP Q9RYA6
A	148	ASP	GLY	SEE REMARK 999	UNP Q9RYA6
A	158	ARG	LYS	SEE REMARK 999	UNP Q9RYA6
A	252	SER	ALA	SEE REMARK 999	UNP Q9RYA6
A	315	SER	PRO	SEE REMARK 999	UNP Q9RYA6
B	94	SER	ALA	SEE REMARK 999	UNP Q9RYA6
B	148	ASP	GLY	SEE REMARK 999	UNP Q9RYA6
B	158	ARG	LYS	SEE REMARK 999	UNP Q9RYA6
B	252	SER	ALA	SEE REMARK 999	UNP Q9RYA6
B	315	SER	PRO	SEE REMARK 999	UNP Q9RYA6
C	94	SER	ALA	SEE REMARK 999	UNP Q9RYA6
C	148	ASP	GLY	SEE REMARK 999	UNP Q9RYA6
C	158	ARG	LYS	SEE REMARK 999	UNP Q9RYA6
C	252	SER	ALA	SEE REMARK 999	UNP Q9RYA6
C	315	SER	PRO	SEE REMARK 999	UNP Q9RYA6
D	94	SER	ALA	SEE REMARK 999	UNP Q9RYA6
D	148	ASP	GLY	SEE REMARK 999	UNP Q9RYA6
D	158	ARG	LYS	SEE REMARK 999	UNP Q9RYA6
D	252	SER	ALA	SEE REMARK 999	UNP Q9RYA6
D	315	SER	PRO	SEE REMARK 999	UNP Q9RYA6

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Mg 1	0	0
2	A	1	Total 1	Mg 1	0	0
2	D	1	Total 1	Mg 1	0	0
2	C	1	Total 1	Mg 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	202	Total 202	O 202	0	0
3	B	207	Total 207	O 207	0	0
3	C	143	Total 143	O 143	0	0
3	D	148	Total 148	O 148	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: N-Acylamino Acid Racemase




• Molecule 1: N-Acylamino Acid Racemase



• Molecule 1: N-Acylamino Acid Racemase



• Molecule 1: N-Acylamino Acid Racemase





4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 4	Depositor
Cell constants a, b, c, α , β , γ	116.67Å 116.67Å 120.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30	Depositor
% Data completeness (in resolution range)	98.8 (30.00-2.30)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.173 , 0.238	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11768	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2817	0.86	11/3822 (0.3%)
1	B	0.66	0/2817	0.83	10/3822 (0.3%)
1	C	0.62	0/2817	0.80	6/3822 (0.2%)
1	D	0.62	0/2817	0.80	9/3822 (0.2%)
All	All	0.64	0/11268	0.82	36/15288 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	195	ASP	CB-CG-OD2	6.81	124.43	118.30
1	B	189	ASP	CB-CG-OD2	6.38	124.04	118.30
1	C	148	ASP	CB-CG-OD2	6.37	124.04	118.30
1	A	148	ASP	CB-CG-OD2	6.36	124.03	118.30
1	B	343	ASP	CB-CG-OD2	6.18	123.86	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	97	LEU	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2766	0	2769	47	0
1	B	2766	0	2769	43	0
1	C	2766	0	2769	50	0
1	D	2766	0	2769	34	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	202	0	0	7	0
3	B	207	0	0	12	1
3	C	143	0	0	14	0
3	D	148	0	0	6	1
All	All	11768	0	11076	171	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

The worst 5 of 171 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:283:HIS:HE1	1:A:316:ASN:H	1.08	1.00
1:C:287:GLN:NE2	1:C:318:ARG:HH11	1.60	0.98
1:A:178:GLN:HE21	1:A:178:GLN:H	1.07	0.97
1:C:283:HIS:HE1	1:C:316:ASN:H	1.17	0.88
1:A:221:GLN:NE2	1:A:245:ASP:H	1.72	0.88

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:B:2462:HOH:O	3:D:4522:HOH:O[3_555]	2.05	0.15

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	356/375 (95%)	343 (96%)	13 (4%)	0	100	100
1	B	356/375 (95%)	342 (96%)	14 (4%)	0	100	100
1	C	356/375 (95%)	342 (96%)	14 (4%)	0	100	100
1	D	356/375 (95%)	344 (97%)	12 (3%)	0	100	100
All	All	1424/1500 (95%)	1371 (96%)	53 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/296 (96%)	268 (94%)	16 (6%)	30	38
1	B	284/296 (96%)	268 (94%)	16 (6%)	30	38
1	C	284/296 (96%)	266 (94%)	18 (6%)	25	32
1	D	284/296 (96%)	273 (96%)	11 (4%)	43	57
All	All	1136/1184 (96%)	1075 (95%)	61 (5%)	31	40

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	298	MET
1	C	36	LYS
1	D	172	LYS
1	B	299	LEU
1	B	334	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	267	ASN
1	B	373	HIS
1	D	231	HIS
1	B	316	ASN
1	C	35	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.