



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 09:58 PM GMT

PDB ID : 1XSS  
Title : Semi-rational engineering of a green-emitting coral fluorescent protein into an efficient highlighter.  
Authors : Tsutsui, H.; Karasawa, S.; Shimizu, H.; Nukina, N.; Miyawaki, A.  
Deposited on : 2004-10-20  
Resolution : 1.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2199 (1.60-1.60)
Ramachandran outliers	78287	2126 (1.60-1.60)
Sidechain outliers	78261	2125 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	225	
1	B	225	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4212 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	0	0
			1808	1156	306	334	12			
1	B	225	Total	C	N	O	S	0	0	0
			1817	1162	308	335	12			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	VAL	MET	CONFLICT	UNP Q53UG7
A	64	DYG	ASP	CHROMOPHORE	UNP Q53UG7
A	64	DYG	TYR	CHROMOPHORE	UNP Q53UG7
A	64	DYG	GLY	CHROMOPHORE	UNP Q53UG7
B	1	VAL	MET	CONFLICT	UNP Q53UG7
B	64	DYG	ASP	CHROMOPHORE	UNP Q53UG7
B	64	DYG	TYR	CHROMOPHORE	UNP Q53UG7
B	64	DYG	GLY	CHROMOPHORE	UNP Q53UG7

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	298	Total 298	O 298	0	0
4	B	285	Total 285	O 285	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: fluorescent protein

Chain A: 



- Molecule 1: fluorescent protein

Chain B: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.74Å 119.08Å 49.29Å 90.00° 120.72° 90.00°	Depositor
Resolution (Å)	14.50 – 1.60	Depositor
% Data completeness (in resolution range)	(Not available) (14.50-1.60)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.197 , 0.229	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4212	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, DYG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/1828	0.67	0/2463
1	B	0.32	0/1837	0.67	0/2474
All	All	0.33	0/3665	0.67	0/4937

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1808	0	1765	28	0
1	B	1817	0	1778	30	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	298	0	0	1	1
4	B	285	0	0	1	2
All	All	4212	0	3543	54	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (54) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:81:GLN:HE22	1:B:187:GLN:H	1.46	0.62
1:A:111:LYS:N	1:A:111:LYS:HD2	2.16	0.60
1:A:83:PHE:HB3	1:A:84:PRO:HA	1.85	0.59
1:B:83:PHE:HB3	1:B:84:PRO:HA	1.85	0.58
1:A:105:ASN:HD21	1:A:119:TYR:HB3	1.70	0.56
1:A:111:LYS:HD2	1:A:111:LYS:H	1.69	0.56
1:B:73:GLU:CD	1:B:73:GLU:H	2.09	0.54
1:B:85:GLU:CD	1:B:184:LYS:HD3	2.28	0.54
1:B:81:GLN:NE2	1:B:187:GLN:H	2.08	0.52
1:A:12:LEU:C	1:A:12:LEU:HD12	2.32	0.50
1:A:32:GLN:NE2	1:A:35:GLU:HG3	2.27	0.49
1:A:126:VAL:HB	1:B:90:GLU:HB3	1.94	0.49
1:A:90:GLU:HB3	1:B:126:VAL:HB	1.94	0.49
1:B:12:LEU:HD12	1:B:12:LEU:C	2.33	0.49
1:B:70:LYS:HD3	1:B:217:LYS:HE2	1.92	0.49
1:A:32:GLN:HE21	1:A:35:GLU:HG3	1.79	0.47
1:B:105:ASN:HD21	1:B:119:TYR:HB3	1.78	0.47
1:B:32:GLN:HE21	1:B:35:GLU:HG3	1.79	0.46
1:B:85:GLU:OE2	1:B:184:LYS:HD3	2.16	0.46
1:A:70:LYS:HD3	1:A:217:LYS:HE2	1.97	0.46
1:B:17:ALA:HA	1:B:21:HIS:O	2.15	0.46
1:B:177:ARG:NH2	4:B:2425:HOH:O	2.45	0.46
1:A:100:ILE:HD11	1:B:100:ILE:HD13	1.98	0.46
1:B:32:GLN:NE2	1:B:35:GLU:HG3	2.31	0.45
1:A:17:ALA:HA	1:A:21:HIS:O	2.16	0.45
1:A:77:ASP:O	1:A:81:GLN:HG3	2.16	0.45
1:A:160:VAL:CG1	1:A:176:PHE:HB2	2.47	0.45
1:A:105:ASN:HD22	1:A:106:ASN:N	2.15	0.45
1:A:137:ARG:HG3	1:A:137:ARG:HH21	1.82	0.45
1:A:138:LYS:HD2	4:A:1441:HOH:O	2.16	0.45
1:A:85:GLU:CD	1:A:184:LYS:HD3	2.37	0.44
1:A:100:ILE:HD13	1:B:100:ILE:HD11	1.99	0.44
1:A:138:LYS:HA	1:A:167:GLN:NE2	2.32	0.44
1:B:60:VAL:HG21	1:B:123:PHE:CD2	2.53	0.44
1:B:9:LYS:HA	1:B:30:SER:HA	1.99	0.43
1:A:85:GLU:OE2	1:A:184:LYS:HD3	2.18	0.43
1:B:108:THR:HG22	1:B:183:LYS:NZ	2.34	0.42
1:B:105:ASN:HD22	1:B:106:ASN:N	2.17	0.42
1:B:24:VAL:CG1	1:B:45:ILE:HD11	2.50	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:136:GLN:HB3	1:B:138:LYS:HZ3	1.85	0.42
1:B:150:TYR:HB3	1:B:191:TYR:CD1	2.54	0.42
1:A:108:THR:HG22	1:A:183:LYS:NZ	2.34	0.42
1:B:160:VAL:CG1	1:B:176:PHE:HB2	2.50	0.42
1:A:150:TYR:HB3	1:A:191:TYR:CD1	2.54	0.42
1:A:131:ASN:N	1:A:131:ASN:HD22	2.18	0.42
1:A:136:GLN:HB3	1:A:138:LYS:HE3	2.02	0.41
1:B:39:ASN:HD22	1:B:39:ASN:H	1.68	0.41
1:A:60:VAL:HG21	1:A:123:PHE:CD2	2.55	0.41
1:B:160:VAL:HG13	1:B:176:PHE:HB2	2.02	0.41
1:B:39:ASN:HD22	1:B:39:ASN:N	2.19	0.41
1:A:24:VAL:CG1	1:A:45:ILE:HD11	2.50	0.41
1:B:19:ASN:HD21	1:B:128:PHE:H	1.69	0.41
1:B:39:ASN:HA	1:B:212:LEU:O	2.21	0.40
1:A:148:LYS:N	1:A:148:LYS:HD2	2.36	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:B:2442:HOH:O	4:B:2442:HOH:O[2_656]	1.12	1.08
4:B:2363:HOH:O	4:B:2363:HOH:O[2_656]	1.73	0.47
4:A:1407:HOH:O	4:A:1407:HOH:O[2_656]	1.85	0.35

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/225 (98%)	219 (99%)	2 (1%)	0	100	100
1	B	222/225 (99%)	218 (98%)	3 (1%)	1 (0%)	38	13
All	All	443/450 (98%)	437 (99%)	5 (1%)	1 (0%)	56	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	112	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/195 (100%)	190 (98%)	4 (2%)	66	37
1	B	195/195 (100%)	192 (98%)	3 (2%)	76	53
All	All	389/390 (100%)	382 (98%)	7 (2%)	71	44

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
1	A	111	LYS
1	A	131	ASN
1	A	148	LYS
1	B	39	ASN
1	B	105	ASN
1	B	148	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	105	ASN
1	A	127	ASN
1	A	131	ASN
1	A	161	ASN
1	A	167	GLN
1	B	19	ASN
1	B	32	GLN
1	B	39	ASN
1	B	81	GLN
1	B	105	ASN
1	B	115	ASN
1	B	127	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	161	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	DYG	A	64	1	24,24,25	4.06	6 (25%)	31,33,35	2.00	6 (19%)
1	DYG	B	64	1	24,24,25	4.17	6 (25%)	31,33,35	1.99	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DYG	A	64	1	-	0/12/31/32	0/2/2/2
1	DYG	B	64	1	-	0/12/31/32	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	DYG	O-C	17.64	1.23	1.11
1	A	64	DYG	O-C	17.04	1.23	1.11
1	B	64	DYG	CB2-CA2	6.84	1.39	1.35
1	A	64	DYG	CB2-CA2	6.68	1.39	1.35
1	A	64	DYG	CA2-C2	-4.15	1.44	1.48
1	B	64	DYG	CA2-C2	-4.14	1.44	1.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	DYG	CG2-CB2	-3.68	1.39	1.46
1	A	64	DYG	CG2-CB2	-3.64	1.39	1.46
1	A	64	DYG	CA3-C	2.33	1.51	1.48
1	A	64	DYG	CD2-CG2	2.31	1.43	1.39
1	B	64	DYG	CA3-C	2.24	1.51	1.48
1	B	64	DYG	CD2-CG2	2.21	1.43	1.39

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	64	DYG	O2-C2-CA2	7.17	135.10	130.96
1	A	64	DYG	O2-C2-CA2	7.15	135.09	130.96
1	A	64	DYG	CA3-N3-C2	-5.32	120.16	123.46
1	B	64	DYG	CA3-N3-C2	-5.11	120.29	123.46
1	B	64	DYG	CA2-N2-C1	3.76	108.81	105.79
1	A	64	DYG	CA2-N2-C1	3.68	108.75	105.79
1	A	64	DYG	N3-C1-N2	-3.30	109.09	111.51
1	B	64	DYG	N3-C1-N2	-3.25	109.12	111.51
1	A	64	DYG	CA3-N3-C1	2.21	129.33	124.64
1	B	64	DYG	CA3-N3-C1	2.13	129.15	124.64
1	B	64	DYG	CA2-C2-N3	-2.11	102.23	103.44
1	A	64	DYG	CA2-C2-N3	-2.02	102.29	103.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.