



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 02:00 AM GMT

PDB ID : 2Y1C
Title : X-RAY STRUCTURE OF 1-DEOXY-D-XYLULOSE 5-PHOSPHATE REDUCTOISOMERASE, DXR, RV2870C, FROM MYCOBACTERIUM TUBERCULOSIS, IN COMPLEX WITH MANGANESE.
Authors : Henriksson, L.M.; Larsson, A.M.S.; Bergfors, T.; Bjorkelid, C.; Unge, T.; Mowbray, S.L.; Jones, T.A.
Deposited on : 2010-12-08
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	398	
1	B	398	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5998 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-DEOXY-D-XYLULOSE 5-PHOSPHATE REDUCTOISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	0	0
			2724	1710	489	516	9			
1	B	379	Total	C	N	O	S	0	2	0
			2799	1753	507	530	9			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	THR	-	EXPRESSION TAG	UNP A2VLK3
A	-7	MET	-	EXPRESSION TAG	UNP A2VLK3
A	-6	ALA	-	EXPRESSION TAG	UNP A2VLK3
A	-5	HIS	-	EXPRESSION TAG	UNP A2VLK3
A	-4	HIS	-	EXPRESSION TAG	UNP A2VLK3
A	-3	HIS	-	EXPRESSION TAG	UNP A2VLK3
A	-2	HIS	-	EXPRESSION TAG	UNP A2VLK3
A	-1	HIS	-	EXPRESSION TAG	UNP A2VLK3
A	0	HIS	-	EXPRESSION TAG	UNP A2VLK3
B	-8	THR	-	EXPRESSION TAG	UNP A2VLK3
B	-7	MET	-	EXPRESSION TAG	UNP A2VLK3
B	-6	ALA	-	EXPRESSION TAG	UNP A2VLK3
B	-5	HIS	-	EXPRESSION TAG	UNP A2VLK3
B	-4	HIS	-	EXPRESSION TAG	UNP A2VLK3
B	-3	HIS	-	EXPRESSION TAG	UNP A2VLK3
B	-2	HIS	-	EXPRESSION TAG	UNP A2VLK3
B	-1	HIS	-	EXPRESSION TAG	UNP A2VLK3
B	0	HIS	-	EXPRESSION TAG	UNP A2VLK3

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mn 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	168	Total 168	O 168	0	0
3	B	305	Total 305	O 305	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 1-DEOXY-D-XYLULOSE 5-PHOSPHATE REDUCTOISOMERASE

Chain A: 



• Molecule 1: 1-DEOXY-D-XYLULOSE 5-PHOSPHATE REDUCTOISOMERASE

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.83Å 67.48Å 84.62Å 90.00° 107.48° 90.00°	Depositor
Resolution (Å)	40.00 – 1.90 37.52 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.5 (40.00-1.90) 98.5 (37.52-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.195 , 0.242 0.195 , 0.240	Depositor DCC
R_{free} test set	2943 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 58370 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5998	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2776	0.52	0/3792
1	B	0.50	0/2853	0.58	0/3897
All	All	0.44	0/5629	0.55	0/7689

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2724	0	2719	17	0
1	B	2799	0	2788	31	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	168	0	0	2	0
3	B	305	0	0	4	0
All	All	5998	0	5507	48	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (48) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:38[B]:ARG:NH2	3:B:2026:HOH:O	2.18	0.77
1:B:376:ARG:O	1:B:380:GLU:HG2	1.85	0.76
1:B:199:ALA:HB1	1:B:207:PRO:HB3	1.74	0.68
1:B:203:TRP:HB3	1:B:205:MET:HB3	1.77	0.66
1:B:205:MET:H	1:B:207:PRO:HD2	1.62	0.65
1:B:37:ASP:HB3	3:B:2025:HOH:O	1.98	0.61
1:A:46:ALA:HB3	1:A:54:LEU:HD22	1.81	0.60
1:A:143:ARG:HG2	1:A:146:GLN:HB2	1.85	0.59
1:B:136:SER:O	1:B:140:ARG:HB2	2.03	0.58
1:A:44:LEU:HB2	1:A:66:ILE:HG12	1.84	0.58
1:A:193:THR:HG22	1:A:194:PRO:HD2	1.86	0.57
1:B:195:GLU:HA	1:B:199:ALA:O	2.07	0.55
1:B:143:ARG:HG3	1:B:146:GLN:HB2	1.90	0.53
1:B:72:HIS:HE1	3:B:2009:HOH:O	1.92	0.52
1:A:180:PRO:HD2	1:A:214:ALA:HA	1.95	0.47
1:B:68:VAL:HG23	1:B:80:ILE:HG21	1.97	0.47
1:A:68:VAL:HG23	1:A:80:ILE:HG21	1.96	0.47
1:B:150:VAL:O	1:B:270:PRO:HB2	2.15	0.46
1:B:379:ARG:NH2	3:B:2300:HOH:O	2.48	0.46
1:B:122:ARG:HD2	1:B:148:VAL:HG21	1.97	0.46
1:B:181:PHE:HB3	1:B:189:LEU:HD21	1.96	0.46
1:B:211:LEU:HD23	1:B:334:ALA:HB1	1.98	0.46
1:A:129:GLU:OE1	1:A:133:ALA:HB3	2.16	0.46
1:B:250:MET:HG2	1:B:260:ALA:CB	2.46	0.45
1:B:247:ILE:HD13	1:B:262:ALA:HB2	1.99	0.45
1:B:206:GLY:N	1:B:207:PRO:CD	2.80	0.45
1:A:218:ASN:ND2	3:A:2088:HOH:O	2.46	0.45
1:B:202:THR:HG22	1:B:202:THR:O	2.17	0.44
1:A:32:ILE:HG23	1:A:39:PHE:HB2	1.98	0.44
1:A:227:HIS:NE2	3:A:2093:HOH:O	2.37	0.43
1:A:360:GLN:HA	1:A:360:GLN:HE21	1.82	0.43
1:A:170:ALA:O	1:A:236:ARG:HD2	2.19	0.42
1:A:115:ALA:O	1:A:119:THR:HG23	2.20	0.42
1:B:170:ALA:O	1:B:236:ARG:HD2	2.19	0.42
1:B:149:PRO:HB2	1:B:154:HIS:HB2	2.01	0.42
1:B:75:GLN:HB3	1:B:75:GLN:HE21	1.65	0.42
1:B:122:ARG:HD2	1:B:148:VAL:CG2	2.50	0.41
1:B:200:HIS:HA	1:B:201:PRO:HD2	1.89	0.41
1:A:122:ARG:NH2	1:A:277:TRP:O	2.52	0.41
1:A:211:LEU:HD23	1:A:334:ALA:HB1	2.02	0.41
1:A:269:LEU:HB3	1:A:270:PRO:CD	2.50	0.41
1:B:250:MET:HG2	1:B:260:ALA:HB2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:200:HIS:O	1:B:202:THR:N	2.54	0.41
1:B:58:ARG:HG3	1:B:63:VAL:O	2.21	0.41
1:B:289:ASP:OD1	1:B:291:HIS:HB2	2.21	0.41
1:B:341:ARG:HH12	1:B:389:MET:C	2.25	0.41
1:A:32:ILE:HG21	1:A:41:VAL:HG23	2.03	0.40
1:B:132:VAL:HG11	1:B:221:LEU:HD22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/398 (92%)	355 (96%)	12 (3%)	1 (0%)	50	37
1	B	379/398 (95%)	372 (98%)	5 (1%)	2 (0%)	38	23
All	All	747/796 (94%)	727 (97%)	17 (2%)	3 (0%)	43	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	201	PRO
1	A	249	SER
1	B	78	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/298 (93%)	269 (97%)	8 (3%)	55	44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	284/298 (95%)	277 (98%)	7 (2%)	60	50
All	All	561/596 (94%)	546 (97%)	15 (3%)	57	47

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	ASP
1	A	129	GLU
1	A	167	ASP
1	A	192	VAL
1	A	193	THR
1	A	209	ASN
1	A	359	ASP
1	A	360	GLN
1	B	71	GLU
1	B	75	GLN
1	B	82	TYR
1	B	205	MET
1	B	311	LEU
1	B	380	GLU
1	B	381	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	60	GLN
1	A	360	GLN
1	B	75	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	372/398 (93%)	0.47	35 (9%) 9 8	17, 33, 61, 71	0
1	B	379/398 (95%)	-0.03	11 (2%) 49 50	12, 21, 37, 61	0
All	All	751/796 (94%)	0.22	46 (6%) 21 20	12, 25, 52, 71	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	203	TRP	6.0
1	A	388	GLY	5.8
1	A	389	MET	5.6
1	A	11	GLY	5.2
1	A	51	LEU	5.0
1	A	78	GLY	4.9
1	A	55	LEU	4.5
1	A	77	VAL	4.5
1	A	205	MET	4.3
1	B	202	THR	4.1
1	A	201	PRO	3.9
1	B	207	PRO	3.8
1	A	384	ARG	3.7
1	A	202	THR	3.7
1	A	291	HIS	3.5
1	A	80	ILE	3.2
1	A	207	PRO	3.2
1	A	191	HIS	3.1
1	B	201	PRO	3.1
1	A	359	ASP	3.0
1	A	60	GLN	2.9
1	A	85	SER	2.9
1	B	206	GLY	2.9
1	A	357	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	83	HIS	2.8
1	A	206	GLY	2.8
1	A	79	ASP	2.7
1	B	11	GLY	2.7
1	A	203	TRP	2.7
1	B	75	GLN	2.7
1	B	291	HIS	2.6
1	A	199	ALA	2.5
1	A	360	GLN	2.5
1	A	342	ILE	2.5
1	A	335	ALA	2.5
1	A	387	SER	2.5
1	B	205	MET	2.4
1	A	69	ALA	2.4
1	A	208	MET	2.4
1	B	77	VAL	2.2
1	A	339	ALA	2.2
1	A	62	GLY	2.2
1	A	356	HIS	2.2
1	B	204	SER	2.1
1	A	204	SER	2.0
1	A	37	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MN	B	1390	1/1	0.09	-0.71	22,22,22,22	0
2	MN	A	1390	1/1	0.10	-1.20	22,22,22,22	0

6.5 Other polymers ⓘ

There are no such residues in this entry.