



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2014 – 04:44 PM GMT

PDB ID : 2Y2G
Title : PENICILLIN-BINDING PROTEIN 1B (PBP-1B) IN COMPLEX WITH AN
ALKYL BORONATE (A01)
Authors : Contreras-Martel, C.; Amoroso, A.; Woon, E.C.; Zervosen, A.; Inglis, S.; Mar-
tins, A.; Verlaine, O.; Rydzik, A.; Job, V.; Luxen, A.; Joris, B.; Schofield,
C.J.; Dessen, A.
Deposited on : 2010-12-15
Resolution : 2.05 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1380 (2.06-2.02)
Clashscore	79885	1577 (2.06-2.02)
Ramachandran outliers	78287	1565 (2.06-2.02)
Sidechain outliers	78261	1565 (2.06-2.02)
RSRZ outliers	66119	1381 (2.06-2.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	494	
1	B	494	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	CL	A	1202	-	X
4	CL	A	1204	-	X
4	CL	B	1235	-	X
4	CL	B	1792	-	X
5	NA	A	1300	-	X
6	EDO	B	1401	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 7930 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PENICILLIN-BINDING PROTEIN 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	466	Total	C	N	O	S	0	2	0
			3610	2260	614	721	15			
1	B	466	Total	C	N	O	S	0	1	0
			3603	2255	613	720	15			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	656	GLY	ASN	ENGINEERED MUTATION	UNP Q7CRA4
A	686	GLN	ARG	ENGINEERED MUTATION	UNP Q7CRA4
A	687	GLN	ARG	ENGINEERED MUTATION	UNP Q7CRA4
B	656	GLY	ASN	ENGINEERED MUTATION	UNP Q7CRA4
B	686	GLN	ARG	ENGINEERED MUTATION	UNP Q7CRA4
B	687	GLN	ARG	ENGINEERED MUTATION	UNP Q7CRA4

- Molecule 2 is [(2,6-DIMETHOXYPHENYL)CARBONYLAMINO]METHYL-TRIHYDROXY-BORON (three-letter code: A01) (formula: C₁₀H₁₅BNO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	B	C	N	O	0	0
			17	1	10	1	5		
2	B	1	Total	B	C	N	O	0	0
			17	1	10	1	5		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	17	Total	Cl	0	0
			17	17		
4	A	13	Total	Cl	0	0
			13	13		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	318	Total	O	0	0
			318	318		
7	B	307	Total	O	0	0
			307	307		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PENICILLIN-BINDING PROTEIN 1B

Chain A: 



• Molecule 1: PENICILLIN-BINDING PROTEIN 1B

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.53Å 97.53Å 149.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.74 – 2.05 48.76 – 2.05	Depositor EDS
% Data completeness (in resolution range)	90.3 (48.74-2.05) 90.3 (48.76-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.6.0095	Depositor
R, R_{free}	0.222 , 0.260 0.223 , 0.260	Depositor DCC
R_{free} test set	4088 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.2	EDS
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	3 of 81301 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7930	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 96.58 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.6408e-10. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4, A01, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/3686	0.51	0/5001
1	B	0.47	0/3676	0.51	0/4989
All	All	0.47	0/7362	0.51	0/9990

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3610	0	0	21	0
1	B	3603	0	0	25	0
2	A	17	0	13	4	0
2	B	17	0	13	4	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	13	0	0	1	0
4	B	17	0	0	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	8	0	12	1	0
6	B	8	0	12	1	0
7	A	318	0	0	5	0
7	B	307	0	0	11	0
All	All	7930	0	50	48	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (48) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:645:ASN:O	1:A:716:ARG:NH2	2.09	0.85
1:A:442:ASN:ND2	1:A:445:GLU:CG	2.47	0.76
1:B:525:TYR:CD2	1:B:526[B]:ARG:NH1	2.59	0.69
1:B:645:ASN:O	1:B:716:ARG:NH2	2.26	0.67
1:B:741:VAL:O	1:B:742:GLU:CG	2.44	0.65
1:A:654:THR:N	2:A:1000:A01:H1	1.96	0.63
1:A:526[B]:ARG:CD	7:A:2176:HOH:O	2.47	0.61
1:B:654:THR:N	2:B:1000:A01:H1	1.99	0.61
1:B:107:GLU:CA	7:B:2001:HOH:O	2.50	0.60
1:B:403:GLN:NE2	7:B:2038:HOH:O	2.34	0.59
2:B:1000:A01:H12	7:B:2244:HOH:O	2.01	0.58
1:B:759:ASN:ND2	7:B:2294:HOH:O	2.36	0.57
1:B:526[B]:ARG:CD	7:B:2139:HOH:O	2.53	0.56
1:A:656:GLY:O	2:A:1000:A01:H151	2.07	0.55
1:B:494:ASN:ND2	4:B:1792:CL:CL	2.78	0.54
1:A:599:GLU:N	7:A:2205:HOH:O	2.40	0.53
1:A:785:SER:O	1:A:788:GLY:N	2.41	0.53
1:A:705:GLN:OE1	6:A:1401:EDO:H11	2.09	0.53
1:A:403:GLN:NE2	7:A:2037:HOH:O	2.42	0.53
1:B:687:GLN:CB	7:B:2242:HOH:O	2.57	0.51
1:B:742:GLU:CG	1:B:742:GLU:O	2.59	0.51
1:A:716:ARG:NH1	7:A:2281:HOH:O	2.46	0.49
1:B:119:ILE:CG2	1:B:340:TYR:OH	2.62	0.48
1:B:525:TYR:CE2	1:B:526[B]:ARG:NH1	2.82	0.48
1:B:656:GLY:O	2:B:1000:A01:H151	2.13	0.48
1:A:554:LEU:N	1:A:555:PRO:CD	2.75	0.48
1:A:498:TYR:OH	2:A:1000:A01:H103	2.14	0.47
1:B:548:GLU:OE2	1:B:550:GLY:N	2.48	0.46
1:A:498:TYR:CZ	2:A:1000:A01:H103	2.52	0.45
1:B:779:TYR:O	1:B:783:TRP:CD1	2.70	0.44
1:A:739:VAL:CG1	1:A:740:SER:N	2.80	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:107:GLU:CG	1:B:115:VAL:CG1	2.95	0.44
1:A:519:ILE:N	1:A:520:PRO:CD	2.81	0.44
1:B:500:ASN:CB	7:B:2111:HOH:O	2.66	0.43
1:B:669:ARG:NH2	1:B:715:GLU:OE2	2.51	0.43
1:B:415:ASP:OD2	1:B:685:SER:OG	2.37	0.42
1:B:602:ASP:N	7:B:2197:HOH:O	2.52	0.42
2:B:1000:A01:C12	7:B:2244:HOH:O	2.66	0.42
1:B:365:GLU:CB	7:B:2020:HOH:O	2.67	0.41
1:B:705:GLN:OE1	6:B:1401:EDO:H11	2.21	0.41
1:B:749:THR:CG2	1:B:750:GLY:N	2.82	0.41
1:A:446:ASN:ND2	1:A:679:ASP:OD2	2.54	0.41
1:A:526[A]:ARG:NE	7:A:2149:HOH:O	2.53	0.41
1:A:749:THR:CG2	1:A:750:GLY:N	2.83	0.41
1:A:787:VAL:O	1:A:787:VAL:CG1	2.69	0.41
1:B:419:ARG:NH1	7:B:2057:HOH:O	2.55	0.40
1:A:760:LYS:N	4:A:1203:CL:CL	2.91	0.40
1:A:443:TYR:O	1:A:447:GLN:NE2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	464/494 (94%)	450 (97%)	14 (3%)	0	100	100
1	B	463/494 (94%)	442 (96%)	21 (4%)	0	100	100
All	All	927/988 (94%)	892 (96%)	35 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of

similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/408 (94%)	369 (96%)	15 (4%)	43	35
1	B	382/408 (94%)	363 (95%)	19 (5%)	34	24
All	All	766/816 (94%)	732 (96%)	34 (4%)	40	29

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	GLU
1	A	349	GLU
1	A	364	LYS
1	A	444	GLN
1	A	516	SER
1	A	526[A]	ARG
1	A	526[B]	ARG
1	A	569	ASN
1	A	681	ASN
1	A	726	SER
1	A	738	LYS
1	A	742	GLU
1	A	744	LYS
1	A	749	THR
1	A	789	SER
1	B	118	SER
1	B	119	ILE
1	B	364	LYS
1	B	366	LEU
1	B	367	LYS
1	B	383	ILE
1	B	385	ASN
1	B	516	SER
1	B	569	ASN
1	B	630	THR
1	B	716	ARG
1	B	726	SER
1	B	744	LYS
1	B	745	GLU
1	B	749	THR
1	B	753	VAL
1	B	785	SER
1	B	787	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	789	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 40 ligands modelled in this entry, 32 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A01	A	1000	1,5	17,17,18	1.48	1 (5%)	21,22,25	2.65	7 (33%)
3	SO4	A	1100	-	4,4,4	0.67	0	6,6,6	0.32	0
6	EDO	A	1400	-	3,3,3	0.55	0	2,2,2	0.48	0
6	EDO	A	1401	-	3,3,3	0.57	0	2,2,2	0.45	0
2	A01	B	1000	1,5	17,17,18	1.46	1 (5%)	21,22,25	2.48	7 (33%)
3	SO4	B	1100	-	4,4,4	0.73	0	6,6,6	0.52	0
6	EDO	B	1400	-	3,3,3	0.51	0	2,2,2	0.51	0
6	EDO	B	1401	-	3,3,3	0.60	0	2,2,2	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A01	A	1000	1,5	-	0/10/13/14	0/1/1/1
3	SO4	A	1100	-	-	0/0/0/0	0/0/0/0
6	EDO	A	1400	-	-	0/1/1/1	0/0/0/0
6	EDO	A	1401	-	-	0/1/1/1	0/0/0/0
2	A01	B	1000	1,5	-	0/10/13/14	0/1/1/1
3	SO4	B	1100	-	-	0/0/0/0	0/0/0/0
6	EDO	B	1400	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1401	-	-	0/1/1/1	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1000	A01	C3-N4	-5.05	1.36	1.44
2	A	1000	A01	C3-N4	-4.94	1.36	1.44

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	A01	O2-B-C3	-7.02	108.35	121.37
2	A	1000	A01	O1-B-C3	-6.41	109.48	121.37
2	B	1000	A01	O2-B-C3	-6.35	109.59	121.37
2	B	1000	A01	O1-B-C3	-6.04	110.17	121.37
2	A	1000	A01	O15-C14-C13	-3.86	117.99	124.37
2	A	1000	A01	O15-C14-C7	3.63	121.50	115.82
2	B	1000	A01	O15-C14-C7	3.57	121.40	115.82
2	B	1000	A01	O15-C14-C13	-3.41	118.74	124.37
2	B	1000	A01	C10-O9-C8	3.30	122.46	117.59
2	A	1000	A01	C10-O9-C8	3.10	122.17	117.59
2	A	1000	A01	O9-C8-C7	2.70	120.05	115.82
2	A	1000	A01	O9-C8-C11	-2.69	119.92	124.37
2	B	1000	A01	O9-C8-C7	2.25	119.34	115.82
2	B	1000	A01	O9-C8-C11	-2.07	120.95	124.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	466/494 (94%)	0.25	33 (7%) 16 14	25, 42, 85, 134	0
1	B	466/494 (94%)	0.27	33 (7%) 16 14	24, 42, 86, 145	0
All	All	932/988 (94%)	0.26	66 (7%) 16 14	24, 42, 86, 145	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	787	VAL	9.0
1	A	787	VAL	7.3
1	B	746	VAL	6.8
1	A	744	LYS	6.4
1	B	117	ALA	6.2
1	A	117	ALA	6.1
1	A	738	LYS	5.7
1	A	739	VAL	5.7
1	A	746	VAL	5.7
1	B	744	LYS	5.6
1	B	739	VAL	5.5
1	B	738	LYS	4.9
1	B	748	VAL	4.7
1	B	118	SER	4.6
1	B	786	ILE	4.4
1	B	737	GLY	4.3
1	A	786	ILE	4.2
1	B	116	ILE	4.1
1	A	788	GLY	4.0
1	A	119	ILE	4.0
1	A	748	VAL	3.9
1	B	789	SER	3.9
1	A	737	GLY	3.9
1	B	743	GLY	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	115	VAL	3.6
1	A	789	SER	3.5
1	A	741	VAL	3.5
1	B	741	VAL	3.5
1	A	745	GLU	3.4
1	B	119	ILE	3.3
1	B	736	PRO	3.3
1	A	780	GLN	3.3
1	B	115	VAL	3.2
1	A	363	ALA	3.2
1	A	736	PRO	3.2
1	A	743	GLY	3.2
1	A	116	ILE	3.1
1	B	750	GLY	3.1
1	B	788	GLY	3.1
1	B	108	ILE	3.0
1	B	114	THR	3.0
1	B	747	GLU	3.0
1	B	780	GLN	2.9
1	A	742	GLU	2.9
1	A	740	SER	2.8
1	B	364	LYS	2.8
1	B	742	GLU	2.8
1	B	745	GLU	2.8
1	A	118	SER	2.7
1	B	740	SER	2.7
1	A	747	GLU	2.6
1	A	108	ILE	2.5
1	A	735	LYS	2.5
1	B	729	LEU	2.5
1	B	444	GLN	2.4
1	A	760	LYS	2.4
1	B	110	TYR	2.4
1	A	749	THR	2.4
1	A	444	GLN	2.4
1	A	364	LYS	2.3
1	A	107	GLU	2.3
1	B	760	LYS	2.2
1	A	782	ALA	2.1
1	A	729	LEU	2.0
1	B	749	THR	2.0
1	B	363	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CL	A	1204	1/1	0.29	13.29	77,77,77,77	0
4	CL	B	1235	1/1	0.17	9.60	60,60,60,60	0
4	CL	B	1792	1/1	0.19	5.25	40,40,40,40	1
4	CL	A	1202	1/1	0.22	5.19	64,64,64,64	0
6	EDO	B	1401	4/4	0.14	3.46	39,55,59,61	0
5	NA	A	1300	1/1	0.21	2.15	40,40,40,40	0
4	CL	B	1214	1/1	0.11	1.85	57,57,57,57	0
4	CL	A	1214	1/1	0.11	1.64	64,64,64,64	0
4	CL	B	1203	1/1	0.29	1.04	76,76,76,76	0
4	CL	B	1229	1/1	0.15	0.86	55,55,55,55	0
4	CL	B	1200	1/1	0.10	0.78	47,47,47,47	0
4	CL	A	1206	1/1	0.14	0.60	56,56,56,56	0
4	CL	B	1791	1/1	0.17	0.55	29,29,29,29	0
4	CL	B	1206	1/1	0.22	0.53	78,78,78,78	0
6	EDO	A	1401	4/4	0.12	0.40	53,57,59,70	0
4	CL	A	1229	1/1	0.13	0.19	59,59,59,59	0
4	CL	A	1208	1/1	0.25	0.14	73,73,73,73	0
5	NA	B	1300	1/1	0.15	0.10	37,37,37,37	0
2	A01	B	1000	17/18	0.12	0.10	43,65,90,101	0
4	CL	A	1203	1/1	0.20	0.07	75,75,75,75	0
6	EDO	A	1400	4/4	0.10	-0.17	42,45,49,52	0
2	A01	A	1000	17/18	0.11	-0.21	35,59,84,105	0
4	CL	B	1213	1/1	0.11	-0.31	53,53,53,53	0
4	CL	B	1201	1/1	0.08	-0.59	48,48,48,48	0
4	CL	A	1216	1/1	0.13	-0.61	67,67,67,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	CL	A	1201	1/1	0.09	-0.62	47,47,47,47	0
4	CL	B	1208	1/1	0.17	-0.81	71,71,71,71	0
4	CL	B	1234	1/1	0.09	-0.97	56,56,56,56	0
4	CL	B	1216	1/1	0.11	-1.06	69,69,69,69	0
6	EDO	B	1400	4/4	0.10	-1.06	38,49,54,60	0
4	CL	B	1224	1/1	0.07	-1.13	59,59,59,59	0
4	CL	A	1213	1/1	0.08	-1.25	60,60,60,60	0
4	CL	B	1202	1/1	0.16	-1.68	62,62,62,62	0
3	SO4	A	1100	5/5	0.08	-1.70	38,41,45,57	0
4	CL	B	1212	1/1	0.07	-2.02	43,43,43,43	0
3	SO4	B	1100	5/5	0.07	-2.23	42,44,52,54	0
4	CL	A	1212	1/1	0.06	-2.35	48,48,48,48	0
4	CL	B	1211	1/1	0.09	-2.47	44,44,44,44	0
4	CL	A	1211	1/1	0.08	-2.64	40,40,40,40	0
4	CL	A	1200	1/1	0.07	-3.04	51,51,51,51	0

6.5 Other polymers ⓘ

There are no such residues in this entry.