



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 10:48 PM GMT

PDB ID : 2Y9X
Title : CRYSTAL STRUCTURE OF PPO3, A TYROSINASE FROM AGARICUS BISPORUS, IN DEOXY-FORM THAT CONTAINS ADDITIONAL UNKNOWN LECTIN-LIKE SUBUNIT, WITH INHIBITOR TROPOLONE
Authors : Ismaya, W.T.; Rozeboom, H.J.; Weijn, A.; Mes, J.J.; Fusetti, F.; Wichers, H.J.; Dijkstra, B.W.
Deposited on : 2011-02-17
Resolution : 2.78 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2193 (2.80-2.76)
Clashscore	79885	2751 (2.80-2.76)
Ramachandran outliers	78287	2699 (2.80-2.76)
Sidechain outliers	78261	2701 (2.80-2.76)
RSRZ outliers	66119	2196 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	391	
1	B	391	
1	C	391	
1	D	391	
2	E	150	
2	F	150	
2	G	150	
2	H	150	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	OTR	A	410	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
5	OTR	B	410	-	X
5	OTR	C	410	-	X
5	OTR	D	410	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 17125 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLYPHENOL OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	391	Total	C	N	O	S	0	0	0
			3207	2069	534	592	12			
1	B	391	Total	C	N	O	S	0	0	0
			3207	2069	534	592	12			
1	C	391	Total	C	N	O	S	0	0	0
			3207	2069	534	592	12			
1	D	391	Total	C	N	O	S	0	0	0
			3207	2069	534	592	12			

- Molecule 2 is a protein called LECTIN-LIKE FOLD PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	136	Total	C	N	O	S	0	0	0
			1056	666	183	205	2			
2	F	136	Total	C	N	O	S	0	0	0
			1056	666	183	205	2			
2	G	136	Total	C	N	O	S	0	0	0
			1056	666	183	205	2			
2	H	136	Total	C	N	O	S	0	0	0
			1056	666	183	205	2			


- Molecule 3 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Cu	0	0
			2	2		
3	A	2	Total	Cu	0	0
			2	2		
3	D	2	Total	Cu	0	0
			2	2		
3	C	2	Total	Cu	0	0
			2	2		

- Molecule 4 is HOLMIUM ATOM (three-letter code: HO) (formula: Ho).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ho	0	0
			1	1		
4	A	1	Total	Ho	0	0
			1	1		
4	D	1	Total	Ho	0	0
			1	1		
4	C	1	Total	Ho	0	0
			1	1		

- Molecule 5 is 2-HYDROXYCYCLOHEPTA-2,4,6-TRIEN-1-ONE (three-letter code: OTR) (formula: C₇H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			9	7	2		
5	B	1	Total	C	O	0	0
			9	7	2		
5	C	1	Total	C	O	0	0
			9	7	2		
5	D	1	Total	C	O	0	0
			9	7	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	4	Total 4	O 4	0	0
6	B	6	Total 6	O 6	0	0
6	C	4	Total 4	O 4	0	0
6	D	5	Total 5	O 5	0	0
6	E	1	Total 1	O 1	0	0
6	F	1	Total 1	O 1	0	0
6	G	3	Total 3	O 3	0	0
6	H	1	Total 1	O 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: POLYPHENOL OXIDASE




• Molecule 1: POLYPHENOL OXIDASE




• Molecule 1: POLYPHENOL OXIDASE




• Molecule 1: POLYPHENOL OXIDASE




• Molecule 2: LECTIN-LIKE FOLD PROTEIN




• Molecule 2: LECTIN-LIKE FOLD PROTEIN



● Molecule 2: LECTIN-LIKE FOLD PROTEIN



● Molecule 2: LECTIN-LIKE FOLD PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	103.84Å 104.82Å 119.36Å 90.00° 110.45° 90.00°	Depositor
Resolution (Å)	58.03 – 2.78 58.03 – 2.78	Depositor EDS
% Data completeness (in resolution range)	89.1 (58.03-2.78) 89.1 (58.03-2.78)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.235 , 0.289 0.243 , 0.296	Depositor DCC
R_{free} test set	2716 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.508	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 25.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 53986 reflections	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	17125	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HO, CU, OTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/3317	0.48	0/4526
1	B	0.35	0/3317	0.48	0/4526
1	C	0.36	0/3317	0.48	0/4526
1	D	0.36	0/3317	0.48	1/4526 (0.0%)
2	E	0.34	0/1082	0.50	0/1470
2	F	0.35	0/1082	0.49	0/1470
2	G	0.36	0/1082	0.49	0/1470
2	H	0.36	0/1082	0.48	0/1470
All	All	0.36	0/17596	0.48	1/23984 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	255	LEU	CA-CB-CG	5.53	128.02	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3207	0	0	1	0
1	B	3207	0	0	3	0
1	C	3207	0	0	2	0
1	D	3207	0	0	4	0
2	E	1056	0	0	1	0
2	F	1056	0	0	2	0
2	G	1056	0	0	0	0
2	H	1056	0	0	2	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	9	0	5	0	0
5	B	9	0	5	0	0
5	C	9	0	5	0	0
5	D	9	0	5	0	0
6	A	4	0	0	0	0
6	B	6	0	0	1	0
6	C	4	0	0	0	0
6	D	5	0	0	0	0
6	E	1	0	0	0	0
6	F	1	0	0	0	0
6	G	3	0	0	0	0
6	H	1	0	0	0	0
All	All	17125	0	20	13	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (13) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:353:ASP:OD2	6:B:2005:HOH:O	2.13	0.67
1:B:84:THR:OG1	1:B:87:THR:OG1	2.27	0.53
1:D:95:ARG:NE	1:D:300:ASP:OD1	2.43	0.51
2:F:52:ARG:NH2	2:F:150:VAL:O	2.45	0.50
1:C:84:THR:OG1	1:C:87:THR:OG1	2.31	0.49
1:C:321:ARG:NH2	2:E:45:THR:O	2.45	0.49
1:A:84:THR:OG1	1:A:87:THR:OG1	2.31	0.48
1:D:84:THR:OG1	1:D:87:THR:OG1	2.36	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:H:139:GLU:O	2:H:145:ARG:NH2	2.52	0.43
1:D:367:ASP:OD1	1:D:367:ASP:N	2.51	0.42
1:B:178:HIS:NE2	1:B:208:GLU:OE1	2.52	0.42
1:D:321:ARG:NH2	2:F:45:THR:O	2.53	0.41
2:H:101:ASN:ND2	2:H:101:ASN:N	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/391 (100%)	371 (95%)	16 (4%)	2 (0%)	38	75
1	B	389/391 (100%)	373 (96%)	16 (4%)	0	100	100
1	C	389/391 (100%)	369 (95%)	20 (5%)	0	100	100
1	D	389/391 (100%)	371 (95%)	17 (4%)	1 (0%)	50	84
2	E	132/150 (88%)	122 (92%)	9 (7%)	1 (1%)	27	64
2	F	132/150 (88%)	124 (94%)	8 (6%)	0	100	100
2	G	132/150 (88%)	123 (93%)	6 (4%)	3 (2%)	10	29
2	H	132/150 (88%)	123 (93%)	8 (6%)	1 (1%)	27	64
All	All	2084/2164 (96%)	1976 (95%)	100 (5%)	8 (0%)	43	80

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	248	VAL
2	G	77	ARG
2	E	112	GLU
2	G	46	PRO
2	G	127	GLN
2	H	124	ALA
1	A	315	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	366	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/342 (100%)	334 (98%)	8 (2%)	63	91
1	B	342/342 (100%)	336 (98%)	6 (2%)	71	94
1	C	342/342 (100%)	330 (96%)	12 (4%)	48	82
1	D	342/342 (100%)	333 (97%)	9 (3%)	59	89
2	E	112/123 (91%)	107 (96%)	5 (4%)	38	73
2	F	112/123 (91%)	105 (94%)	7 (6%)	25	57
2	G	112/123 (91%)	108 (96%)	4 (4%)	47	82
2	H	112/123 (91%)	108 (96%)	4 (4%)	47	82
All	All	1816/1860 (98%)	1761 (97%)	55 (3%)	53	86

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	125	ILE
1	A	146	ASP
1	A	168	THR
1	A	253	ASN
1	A	255	LEU
1	A	259	HIS
1	A	367	ASP
1	B	3	ASP
1	B	146	ASP
1	B	158	LYS
1	B	255	LEU
1	B	259	HIS
1	B	385	ASP
1	C	3	ASP
1	C	40	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	125	ILE
1	C	146	ASP
1	C	168	THR
1	C	201	TYR
1	C	207	GLN
1	C	255	LEU
1	C	259	HIS
1	C	261	THR
1	C	367	ASP
1	C	377	GLU
1	D	3	ASP
1	D	146	ASP
1	D	158	LYS
1	D	168	THR
1	D	234	LYS
1	D	259	HIS
1	D	266	ILE
1	D	367	ASP
1	D	390	HIS
2	E	24	ASN
2	E	45	THR
2	E	101	ASN
2	E	127	GLN
2	E	130	THR
2	F	52	ARG
2	F	59	LEU
2	F	95	SER
2	F	101	ASN
2	F	103	GLU
2	F	127	GLN
2	F	130	THR
2	G	10	ASP
2	G	80	THR
2	G	112	GLU
2	G	125	SER
2	H	10	ASP
2	H	85	SER
2	H	101	ASN
2	H	112	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 12 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	OTR	A	410	-	8,9,9	1.33	2 (25%)	9,11,11	2.19	1 (11%)
5	OTR	B	410	-	8,9,9	1.44	2 (25%)	9,11,11	2.77	1 (11%)
5	OTR	C	410	-	8,9,9	1.34	2 (25%)	9,11,11	2.14	1 (11%)
5	OTR	D	410	-	8,9,9	1.39	2 (25%)	9,11,11	2.70	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	OTR	A	410	-	-	0/0/0/0	0/1/1/1
5	OTR	B	410	-	-	0/0/0/0	0/1/1/1
5	OTR	C	410	-	-	0/0/0/0	0/1/1/1
5	OTR	D	410	-	-	0/0/0/0	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	410	0TR	CB3-CA3	-2.42	1.33	1.38
5	B	410	0TR	CB3-CA3	-2.42	1.33	1.38
5	B	410	0TR	OA1-CA1	2.30	1.26	1.23
5	C	410	0TR	CB3-CA3	-2.25	1.34	1.38
5	A	410	0TR	CB3-CA3	-2.24	1.34	1.38
5	D	410	0TR	OA1-CA1	2.16	1.26	1.23
5	A	410	0TR	OA1-CA1	2.02	1.26	1.23
5	C	410	0TR	OA1-CA1	2.02	1.26	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	410	0TR	CA6-CA1-CA2	-8.26	123.50	127.96
5	D	410	0TR	CA6-CA1-CA2	-8.04	123.62	127.96
5	A	410	0TR	CA6-CA1-CA2	-6.47	124.47	127.96
5	C	410	0TR	CA6-CA1-CA2	-6.29	124.57	127.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	391/391 (100%)	0.07	6 (1%) 70 73	38, 49, 62, 74	0
1	B	391/391 (100%)	0.16	10 (2%) 53 56	39, 51, 64, 73	0
1	C	391/391 (100%)	0.26	11 (2%) 50 54	40, 55, 74, 91	0
1	D	391/391 (100%)	0.07	2 (0%) 88 90	38, 52, 68, 80	0
2	E	136/150 (90%)	0.24	1 (0%) 84 86	40, 51, 67, 84	0
2	F	136/150 (90%)	0.26	3 (2%) 59 62	42, 54, 74, 84	0
2	G	136/150 (90%)	0.43	9 (6%) 18 17	40, 54, 74, 88	0
2	H	136/150 (90%)	0.44	8 (5%) 22 22	41, 54, 78, 93	0
All	All	2108/2164 (97%)	0.19	50 (2%) 56 59	38, 52, 68, 93	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	SER	5.6
1	C	117	THR	4.8
2	E	125	SER	4.7
1	C	2	SER	4.5
1	D	246	ALA	4.2
2	G	126	SER	4.2
2	G	125	SER	4.2
2	H	125	SER	4.0
2	F	125	SER	3.9
2	H	126	SER	3.6
2	F	9	LEU	3.5
2	H	114	GLY	3.5
1	B	248	VAL	3.5
2	H	28	THR	3.5
1	C	121	GLN	3.5
2	G	35	SER	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	122	ALA	3.1
1	C	116	PHE	3.0
2	H	127	GLN	2.9
1	A	2	SER	2.9
1	B	75	LEU	2.9
1	A	247	VAL	2.9
1	B	2	SER	2.8
2	G	127	GLN	2.7
2	H	124	ALA	2.7
2	G	136	LEU	2.7
1	B	73	PRO	2.7
2	H	115	GLY	2.7
1	A	120	ASP	2.6
2	F	136	LEU	2.6
1	B	392	GLY	2.6
2	G	9	LEU	2.5
1	C	167	GLY	2.5
1	A	270	PRO	2.5
1	C	43	LEU	2.4
1	A	276	VAL	2.3
2	G	114	GLY	2.3
1	A	246	ALA	2.2
1	B	373	GLY	2.2
1	B	76	HIS	2.2
2	G	124	ALA	2.2
1	B	125	ILE	2.1
1	C	111	THR	2.1
2	H	128	GLY	2.1
1	C	114	GLN	2.1
1	B	3	ASP	2.0
1	C	163	THR	2.0
2	G	128	GLY	2.0
1	C	115	ARG	2.0
1	B	271	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	OTR	C	410	9/9	0.40	13.30	102,102,102,102	0
5	OTR	A	410	9/9	0.26	3.11	82,82,83,83	0
5	OTR	D	410	9/9	0.26	2.63	73,73,73,74	0
5	OTR	B	410	9/9	0.21	2.10	71,71,71,71	0
3	CU	D	401	1/1	0.07	-2.07	42,42,42,42	0
4	HO	A	402	1/1	0.10	-2.27	63,63,63,63	0
4	HO	C	402	1/1	0.09	-2.51	72,72,72,72	0
4	HO	B	402	1/1	0.09	-2.78	59,59,59,59	0
3	CU	D	400	1/1	0.05	-3.07	44,44,44,44	0
3	CU	C	400	1/1	0.08	-4.29	42,42,42,42	0
3	CU	A	400	1/1	0.05	-4.39	43,43,43,43	0
3	CU	B	400	1/1	0.06	-4.62	42,42,42,42	0
4	HO	D	402	1/1	0.07	-4.93	64,64,64,64	0
3	CU	A	401	1/1	0.08	-5.57	40,40,40,40	0
3	CU	C	401	1/1	0.04	-7.08	42,42,42,42	0
3	CU	B	401	1/1	0.05	-11.00	42,42,42,42	0

6.5 Other polymers ⓘ

There are no such residues in this entry.