



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 06:28 PM GMT

PDB ID : 1YOK
Title : crystal structure of human LRH-1 bound with TIF-2 peptide and phosphatidylglycerol
Authors : Krylova, I.N.; Sablin, E.P.; Moore, J.; Xu, R.X.; Waitt, G.M.; MacKay, J.A.; Juzumiene, D.; Bynum, J.M.; Madauss, K.; Montana, V.; Lebedeva, L.; Suzawa, M.; Williams, J.D.; Williams, S.P.; Guy, R.K.; Thornton, J.W.; Fletterick, R.J.; Willson, T.M.; Ingraham, H.A.
Deposited on : 2005-01-27
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	256	
2	B	14	
2	C	14	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	P6L	A	100	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2253 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Orphan nuclear receptor NR5A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	S	0	0	0
			1907	1224	317	353	13			


There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	286	MET	-	CLONING ARTIFACT	UNP O00482
A	287	LYS	-	CLONING ARTIFACT	UNP O00482
A	288	LYS	-	CLONING ARTIFACT	UNP O00482
A	289	HIS	-	CLONING ARTIFACT	UNP O00482
A	290	HIS	-	CLONING ARTIFACT	UNP O00482
A	291	HIS	-	CLONING ARTIFACT	UNP O00482
A	292	HIS	-	CLONING ARTIFACT	UNP O00482
A	293	HIS	-	CLONING ARTIFACT	UNP O00482
A	294	HIS	-	CLONING ARTIFACT	UNP O00482
A	295	LEU	-	CLONING ARTIFACT	UNP O00482
A	296	VAL	-	CLONING ARTIFACT	UNP O00482
A	297	PRO	-	CLONING ARTIFACT	UNP O00482
A	298	ARG	-	CLONING ARTIFACT	UNP O00482
A	299	GLY	-	CLONING ARTIFACT	UNP O00482

- Molecule 2 is a protein called Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	11	Total	C	N	O	0	0	0
			93	60	16	17			
2	C	14	Total	C	N	O	0	0	0
			116	72	19	25			

- Molecule 3 is (2S)-3-[[[(2S)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-2-[(6E)-HEXADEC-6-ENOYLOXY]PROPYL(8E)-OCTADEC-8-ENOATE (three-letter code: P6L) (formula: C₄₀H₇₅O₁₀P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			51	40	10	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	70	Total	O	0	0
			70	70		
4	B	8	Total	O	0	0
			8	8		
4	C	8	Total	O	0	0
			8	8		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.94Å 67.20Å 79.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.50 38.99 – 2.48	Depositor EDS
% Data completeness (in resolution range)	91.9 (40.00-2.50) 91.5 (38.99-2.48)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.48Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.230 , 0.290 0.227 , 0.282	Depositor DCC
R_{free} test set	647 reflections (6.04%)	DCC
Wilson B-factor (Å ²)	58.2	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 54.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 11270 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2253	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: P6L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1941	0.60	0/2623
2	B	0.42	0/93	0.66	0/124
2	C	0.43	0/116	0.65	0/154
All	All	0.43	0/2150	0.60	0/2901

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	747	TYR	Sidechain

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1907	0	1908	91	0
2	B	93	0	97	7	0
2	C	116	0	109	11	0
3	A	51	0	74	24	0
4	A	70	0	0	5	0
4	B	8	0	0	1	0
4	C	8	0	0	0	0
All	All	2253	0	2188	114	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 26.

The worst 5 of 114 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:416:ILE:HG23	3:A:100:P6L:H292	1.46	0.96
1:A:353:LEU:HD22	1:A:531:LEU:HD21	1.48	0.95
1:A:356:ILE:HD12	1:A:386:LEU:HD23	1.49	0.92
3:A:100:P6L:H182	3:A:100:P6L:H291	1.55	0.89
2:C:740:LYS:O	2:C:741:GLU:HB3	1.72	0.87

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/256 (91%)	212 (91%)	15 (6%)	5 (2%)	10	15
2	B	9/14 (64%)	8 (89%)	0	1 (11%)	1	0
2	C	12/14 (86%)	8 (67%)	2 (17%)	2 (17%)	0	0
All	All	253/284 (89%)	228 (90%)	17 (7%)	8 (3%)	6	8

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	331	ALA
1	A	332	ASN
2	C	741	GLU
1	A	302	PRO
2	B	751	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/231 (91%)	206 (98%)	5 (2%)	61	86
2	B	10/13 (77%)	10 (100%)	0	100	100
2	C	12/13 (92%)	11 (92%)	1 (8%)	16	29
All	All	233/257 (91%)	227 (97%)	6 (3%)	59	84

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	379	GLN
2	C	741	GLU
1	A	463	ASN
1	A	361	ARG
1	A	518	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	410	GLN
2	C	742	ASN
1	A	474	GLN
1	A	327	GLN
1	A	430	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	P6L	A	100	-	50,50,50	0.94	2 (4%)	56,56,56	1.36	11 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	P6L	A	100	-	-	2/55/55/55	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	100	P6L	C23-C22	3.71	1.52	1.31
3	A	100	P6L	C34-C33	3.70	1.52	1.31

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	100	P6L	O4-C14-C18	4.25	120.86	111.56
3	A	100	P6L	O12-P11-O9	3.83	115.61	104.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	100	P6L	C7-C5-C6	2.59	117.77	111.86
3	A	100	P6L	C32-C33-C34	-2.59	110.77	125.43
3	A	100	P6L	C24-C23-C22	-2.58	110.80	125.43

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	100	P6L	C5-O4-C14-C18
3	A	100	P6L	C5-O4-C14-O15

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	236/256 (92%)	0.39	14 (5%) 22 22	36, 58, 90, 134	0
2	B	11/14 (78%)	0.02	0 100 100	43, 53, 75, 86	0
2	C	14/14 (100%)	1.49	4 (28%) 1 1	58, 69, 107, 114	0
All	All	261/284 (91%)	0.43	18 (6%) 17 16	36, 59, 91, 134	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	740	LYS	8.6
1	A	333	ARG	6.4
1	A	300	SER	5.7
1	A	334	SER	4.7
2	C	753	ASP	4.5

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	P6L	A	100	51/51	0.27	2.28	65,92,101,101	0

6.5 Other polymers ⓘ

There are no such residues in this entry.