



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 05:16 AM GMT

PDB ID : 1Z2C  
Title : Crystal structure of mDIA1 GBD-FH3 in complex with RhoC-GMPPNP  
Authors : Rose, R.; Weyand, M.; Lammers, M.; Ishizaki, T.; Ahmadian, M.R.; Wittinghofer, A.  
Deposited on : 2005-03-08  
Resolution : 3.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	193	
1	C	193	
2	B	383	
2	D	383	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	MG	B	5	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8399 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rho-related GTP-binding protein RhoC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	179	Total	C	N	O	S	0	0	0
			1419	894	242	273	10			
1	C	179	Total	C	N	O	S	0	0	0
			1416	892	242	273	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	ASN	PHE	ENGINEERED	UNP P08134
C	25	ASN	PHE	ENGINEERED	UNP P08134


- Molecule 2 is a protein called Diaphanous protein homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	346	Total	C	N	O	S	0	0	0
			2788	1756	477	531	24			
2	D	335	Total	C	N	O	S	0	0	0
			2709	1712	461	512	24			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is PHOSPHOAMINOPHOSPHONICACID-GUANYLATE ESTER (three-letter code: GNP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>13</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
4	C	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Rho-related GTP-binding protein RhoC

Chain A: 




- Molecule 1: Rho-related GTP-binding protein RhoC

Chain C: 




- Molecule 2: Diaphanous protein homolog 1

Chain B: 



- Molecule 2: Diaphanous protein homolog 1

Chain D: 



R322	Y323	H324	E328	R331	L338	Q339	E340	L341	I344	E345	D363	R375	K396	A397	E398	L402	L405	L410	V411	R412	M413	L424	I425	E426	Q431	I432	K436	ASN	GLY	THR	ASP	PRO	ASP	PHE	LYS	CYS	ARG	HIS	LEU	GLN	ILE	ASP
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	148.64Å 85.35Å 123.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.00 19.80 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-3.00) 100.0 (19.80-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.95 (at 2.98Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.211 , 0.285 0.209 , 0.280	Depositor DCC
$R_{free}$ test set	1605 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	57.5	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 23.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 31996 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8399	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.71	0/1445	0.85	5/1953 (0.3%)
1	C	0.62	0/1442	0.74	1/1950 (0.1%)
2	B	0.78	3/2825 (0.1%)	0.83	1/3801 (0.0%)
2	D	0.87	3/2745 (0.1%)	0.86	3/3694 (0.1%)
All	All	0.77	6/8457 (0.1%)	0.83	10/11398 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	107	LYS	CD-CE	20.83	2.03	1.51
2	D	107	LYS	CE-NZ	14.29	1.84	1.49
2	B	347	GLU	CG-CD	8.44	1.64	1.51
2	B	347	GLU	CB-CG	7.38	1.66	1.52
2	B	98	GLU	CG-CD	6.84	1.62	1.51
2	D	398	GLU	CG-CD	-5.06	1.44	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	107	LYS	CD-CE-NZ	-11.82	84.51	111.70
2	D	107	LYS	CG-CD-CE	-7.66	88.91	111.90
1	A	68	ARG	NE-CZ-NH1	6.46	123.53	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	329	LEU	CA-CB-CG	6.17	129.50	115.30
1	C	179	LEU	CA-CB-CG	5.86	128.79	115.30
1	A	67	ASP	CB-CG-OD1	5.63	123.37	118.30
2	D	263	PRO	C-N-CA	5.61	135.73	121.70
1	A	70	ARG	NE-CZ-NH1	5.57	123.08	120.30
1	A	8	LEU	CA-CB-CG	5.49	127.93	115.30
1	A	59	ASP	CB-CG-OD1	5.47	123.22	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	216	MET	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1419	0	0	7	0
1	C	1416	0	0	15	0
2	B	2788	0	0	13	0
2	D	2709	0	0	22	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	32	0	13	0	0
4	C	32	0	13	1	0
All	All	8399	0	26	56	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (56) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:107:LYS:NZ	2:D:107:LYS:CE	1.84	1.38

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:107:LYS:CE	2:D:107:LYS:CD	2.03	1.35
2:D:107:LYS:CE	2:D:107:LYS:CG	2.61	0.78
1:C:140:ARG:CB	1:C:140:ARG:NH1	2.47	0.78
1:A:25:ASN:OD1	1:A:171:PHE:CD1	2.44	0.70
2:D:201:ASP:OD1	2:D:203:ARG:NH1	2.24	0.70
2:D:322:ARG:NH1	2:D:363:ASP:OD1	2.26	0.68
2:B:214:ALA:O	2:B:217:ASN:CB	2.44	0.66
2:D:264:GLU:O	2:D:265:ASP:CB	2.45	0.65
2:D:107:LYS:NZ	2:D:107:LYS:CD	2.61	0.63
2:B:322:ARG:NH1	2:B:363:ASP:OD1	2.36	0.58
2:B:216:MET:O	2:B:218:ASN:N	2.37	0.58
2:B:320:ASP:OD1	2:B:370:ARG:NH2	2.39	0.56
1:A:65:ASP:N	1:A:65:ASP:OD2	2.40	0.54
2:B:121:HIS:O	2:B:123:SER:N	2.42	0.53
1:C:1:MET:O	1:C:2:ALA:CB	2.57	0.53
1:C:160:SER:O	1:C:164:LYS:N	2.43	0.52
2:D:101:GLN:OE1	2:D:105:ARG:NH2	2.43	0.51
2:D:285:ARG:NH1	2:D:328:GLU:OE1	2.44	0.51
1:C:7:LYS:N	1:C:78:ASP:OD2	2.44	0.51
2:D:322:ARG:NH1	2:D:322:ARG:CG	2.75	0.50
1:C:5:ARG:NE	1:C:54:GLU:OE1	2.46	0.49
2:D:201:ASP:OD2	2:D:201:ASP:C	2.51	0.47
1:A:85:SER:OG	1:A:87:ASP:OD1	2.32	0.47
1:A:122:ARG:NH2	1:A:158:GLU:OE1	2.48	0.47
1:C:67:ASP:O	2:D:165:ASN:ND2	2.49	0.46
2:B:317:GLU:OE1	2:B:317:GLU:N	2.48	0.46
2:B:216:MET:C	2:B:218:ASN:N	2.69	0.46
1:C:163:THR:O	1:C:164:LYS:CB	2.64	0.45
2:D:268:GLU:O	2:D:272:GLU:CG	2.64	0.45
1:A:74:TYR:N	1:A:75:PRO:CD	2.80	0.45
1:C:93:GLU:CA	1:C:93:GLU:OE1	2.65	0.45
1:A:128:ARG:NH1	1:A:128:ARG:CG	2.79	0.45
1:C:118:LYS:O	1:C:121:LEU:N	2.50	0.44
2:D:324:HIS:C	2:D:324:HIS:CD2	2.91	0.44
1:C:120:ASP:OD1	1:C:160:SER:OG	2.36	0.44
2:B:262:GLN:CG	2:B:263:PRO:CD	2.97	0.43
1:A:13:ASP:OD2	1:A:94:ASN:ND2	2.51	0.43
2:D:190:HIS:O	2:D:237:ARG:NH2	2.52	0.43
2:B:97:ASN:C	2:B:97:ASN:OD1	2.57	0.42
2:B:297:THR:O	2:B:302:LYS:NZ	2.52	0.42
1:C:124:ASP:O	1:C:127:THR:N	2.53	0.42
2:B:283:VAL:CG1	2:B:284:GLU:N	2.83	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:140:ARG:CB	1:C:140:ARG:CZ	2.98	0.42
2:D:189:LEU:O	2:D:191:ASP:N	2.53	0.42
2:D:102:GLN:N	2:D:103:PRO:CD	2.83	0.41
2:D:139:TYR:O	2:D:140:ILE:C	2.58	0.41
2:D:262:GLN:O	2:D:262:GLN:CG	2.68	0.41
2:B:177:GLU:O	2:B:178:GLY:C	2.59	0.41
1:C:18:LYS:NZ	4:C:195:GNP:O1B	2.54	0.41
1:C:122:ARG:NH2	1:C:158:GLU:OE2	2.54	0.41
2:B:167:PRO:O	2:B:169:SER:N	2.54	0.41
2:D:285:ARG:NH1	2:D:328:GLU:OE2	2.55	0.40
1:C:5:ARG:NH2	1:C:54:GLU:OE1	2.54	0.40
2:D:122:THR:O	2:D:123:SER:C	2.60	0.40
2:D:338:LEU:O	2:D:341:LEU:N	2.53	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/193 (92%)	156 (88%)	18 (10%)	3 (2%)	14	54
1	C	177/193 (92%)	160 (90%)	14 (8%)	3 (2%)	14	54
2	B	340/383 (89%)	307 (90%)	28 (8%)	5 (2%)	15	58
2	D	329/383 (86%)	292 (89%)	32 (10%)	5 (2%)	15	58
All	All	1023/1152 (89%)	915 (89%)	92 (9%)	16 (2%)	14	56

All (16) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	VAL
2	B	122	THR
2	B	217	ASN
1	C	28	ASP
2	D	190	HIS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	265	ASP
2	D	345	GLU
1	A	133	LYS
1	C	2	ALA
2	D	122	THR
1	A	28	ASP
2	B	168	VAL
2	D	231	GLY
2	B	133	SER
1	C	133	LYS
2	B	261	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/170 (92%)	135 (86%)	22 (14%)	5	23
1	C	156/170 (92%)	139 (89%)	17 (11%)	9	35
2	B	316/350 (90%)	280 (89%)	36 (11%)	8	33
2	D	306/350 (87%)	260 (85%)	46 (15%)	4	20
All	All	935/1040 (90%)	814 (87%)	121 (13%)	6	26

All (121) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	8	LEU
1	A	25	ASN
1	A	27	LYS
1	A	32	GLU
1	A	37	THR
1	A	43	ILE
1	A	53	VAL
1	A	65	ASP
1	A	75	PRO
1	A	80	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	90	ASP
1	A	95	ILE
1	A	104	LYS
1	A	113	ILE
1	A	122	ARG
1	A	129	ARG
1	A	134	MET
1	A	145	ARG
1	A	152	SER
1	A	164	LYS
1	A	179	LEU
2	B	109	ILE
2	B	110	VAL
2	B	132	SER
2	B	134	ARG
2	B	152	LEU
2	B	154	SER
2	B	158	SER
2	B	173	THR
2	B	179	LEU
2	B	201	ASP
2	B	206	HIS
2	B	228	THR
2	B	230	GLU
2	B	260	LEU
2	B	262	GLN
2	B	280	MET
2	B	295	SER
2	B	306	LEU
2	B	308	LEU
2	B	317	GLU
2	B	319	LEU
2	B	323	VAL
2	B	325	ILE
2	B	331	ARG
2	B	338	LEU
2	B	340	GLU
2	B	345	GLU
2	B	351	VAL
2	B	370	ARG
2	B	377	GLU
2	B	380	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	391	THR
2	B	411	VAL
2	B	418	ARG
2	B	436	LYS
2	B	440	ASP
1	C	16	CYS
1	C	28	ASP
1	C	29	GLN
1	C	32	GLU
1	C	43	ILE
1	C	60	THR
1	C	65	ASP
1	C	79	VAL
1	C	123	GLN
1	C	125	GLU
1	C	127	THR
1	C	137	GLU
1	C	142	GLU
1	C	150	ARG
1	C	168	ARG
1	C	176	ARG
1	C	179	LEU
2	D	83	VAL
2	D	93	ASP
2	D	98	GLU
2	D	99	GLU
2	D	117	SER
2	D	144	ARG
2	D	154	SER
2	D	172	GLN
2	D	173	THR
2	D	179	LEU
2	D	182	LEU
2	D	186	LEU
2	D	187	LYS
2	D	203	ARG
2	D	209	ILE
2	D	213	LYS
2	D	226	LEU
2	D	227	GLU
2	D	228	THR
2	D	230	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	252	LYS
2	D	253	LEU
2	D	255	SER
2	D	269	ARG
2	D	283	VAL
2	D	285	ARG
2	D	290	LEU
2	D	294	LYS
2	D	314	THR
2	D	322	ARG
2	D	331	ARG
2	D	340	GLU
2	D	344	ILE
2	D	345	GLU
2	D	375	ARG
2	D	396	LYS
2	D	402	LEU
2	D	405	LEU
2	D	410	LEU
2	D	411	VAL
2	D	413	ASN
2	D	424	LEU
2	D	426	GLU
2	D	431	GLN
2	D	432	ILE
2	D	436	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GNP	A	195	3	34,34,34	1.58	7 (20%)	50,54,54	4.97	8 (16%)
4	GNP	C	195	3	34,34,34	1.91	7 (20%)	50,54,54	5.99	13 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GNP	A	195	3	-	0/18/38/38	0/1/3/3
4	GNP	C	195	3	-	0/18/38/38	0/1/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	195	GNP	PG-N3B	-5.63	1.59	1.64
4	C	195	GNP	PB-N3B	-4.91	1.60	1.64
4	A	195	GNP	PG-O1G	4.86	1.52	1.46
4	C	195	GNP	PB-O3A	-3.65	1.54	1.59
4	C	195	GNP	PG-O1G	3.63	1.50	1.46
4	C	195	GNP	PA-O3A	-3.54	1.53	1.59
4	A	195	GNP	PB-N3B	-3.30	1.61	1.64
4	A	195	GNP	PB-O3A	-3.03	1.54	1.59
4	A	195	GNP	PA-O3A	-2.60	1.55	1.59
4	A	195	GNP	PG-N3B	-2.52	1.62	1.64
4	A	195	GNP	PB-O2B	-2.34	1.48	1.55
4	C	195	GNP	C8-N7	-2.10	1.30	1.34
4	C	195	GNP	PB-O2B	-2.06	1.49	1.55
4	A	195	GNP	C8-N7	-2.01	1.30	1.34

All (21) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	195	GNP	C6-C5-N7	-40.53	128.68	134.14
4	A	195	GNP	C6-C5-N7	-33.63	129.61	134.14
4	A	195	GNP	O3G-PG-O1G	-4.66	101.65	113.60
4	C	195	GNP	PB-N3B-PG	-4.35	122.75	130.07
4	C	195	GNP	O2B-PB-O1B	3.95	119.01	109.89
4	C	195	GNP	O3G-PG-O1G	-3.78	103.89	113.60
4	C	195	GNP	O1B-PB-N3B	-3.63	106.34	111.83
4	A	195	GNP	C6-N1-C2	3.13	124.98	119.51
4	C	195	GNP	C6-N1-C2	2.89	124.56	119.51
4	C	195	GNP	O3G-PG-O2G	2.88	115.91	107.66
4	A	195	GNP	O3G-PG-O2G	2.81	115.71	107.66
4	C	195	GNP	C4-C5-N7	2.76	111.89	109.52
4	C	195	GNP	C3'-C2'-C1'	2.33	104.56	100.91
4	A	195	GNP	O2B-PB-O1B	2.31	115.21	109.89
4	C	195	GNP	C2-N3-C4	-2.27	111.90	115.09
4	A	195	GNP	O4'-C1'-N9	2.23	110.52	108.44
4	A	195	GNP	PB-N3B-PG	-2.16	126.44	130.07
4	C	195	GNP	O3A-PB-N3B	2.15	112.56	106.59
4	C	195	GNP	C2'-C3'-C4'	2.10	106.83	102.65
4	C	195	GNP	C2'-C1'-N9	-2.08	107.92	113.27
4	A	195	GNP	N2-C2-N1	-2.05	115.60	117.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	179/193 (92%)	-0.35	0 100 100	28, 52, 93, 108	0
1	C	179/193 (92%)	-0.31	0 100 100	35, 56, 100, 119	0
2	B	346/383 (90%)	-0.39	0 100 100	26, 44, 76, 101	0
2	D	335/383 (87%)	-0.40	0 100 100	21, 49, 78, 95	0
All	All	1039/1152 (90%)	-0.37	0 100 100	21, 48, 85, 119	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	B	5	1/1	0.43	8.97	66,66,66,66	0
3	MG	A	194	1/1	0.18	0.82	31,31,31,31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	C	194	1/1	0.21	0.39	56,56,56,56	0
4	GNP	C	195	32/32	0.13	-0.77	49,57,61,61	0
4	GNP	A	195	32/32	0.12	-0.96	42,54,56,57	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.