



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 11:10 AM GMT

PDB ID : 1ZTU  
Title : Structure of the chromophore binding domain of bacterial phytochrome  
Authors : Wagner, J.R.; Brunzelle, J.S.; Forest, K.T.; Vierstra, R.D.  
Deposited on : 2005-05-27  
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	341	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2485 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bacteriophytochrome.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	Se	0	1	0
			2408	1537	429	433	4	5			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MSE	-	cloning artifact	UNP Q9RZA4
A	-12	ALA	-	cloning artifact	UNP Q9RZA4
A	-11	SER	-	cloning artifact	UNP Q9RZA4
A	-10	MSE	-	cloning artifact	UNP Q9RZA4
A	-9	THR	-	cloning artifact	UNP Q9RZA4
A	-8	GLY	-	cloning artifact	UNP Q9RZA4
A	-7	GLY	-	cloning artifact	UNP Q9RZA4
A	-6	GLN	-	cloning artifact	UNP Q9RZA4
A	-5	GLN	-	cloning artifact	UNP Q9RZA4
A	-4	MSE	-	cloning artifact	UNP Q9RZA4
A	-3	GLY	-	cloning artifact	UNP Q9RZA4
A	-2	ARG	-	cloning artifact	UNP Q9RZA4
A	-1	GLY	-	cloning artifact	UNP Q9RZA4
A	0	SER	-	cloning artifact	UNP Q9RZA4
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q9RZA4
A	144	MSE	MET	MODIFIED RESIDUE	UNP Q9RZA4
A	174	MSE	MET	MODIFIED RESIDUE	UNP Q9RZA4
A	240	THR	PRO	ENGINEERED	UNP Q9RZA4
A	259	MSE	MET	MODIFIED RESIDUE	UNP Q9RZA4
A	261	MSE	MET	MODIFIED RESIDUE	UNP Q9RZA4
A	267	MSE	MET	MODIFIED RESIDUE	UNP Q9RZA4
A	322	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	323	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	324	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	325	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	326	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	327	HIS	-	EXPRESSION TAG	UNP Q9RZA4

- Molecule 2 is BILIVERDINE IX ALPHA (three-letter code: BLA) (formula:  $C_{33}H_{34}N_4O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	43	33	4	6	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	34	34	34	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacteriophytochrome

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.88Å 133.67Å 49.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 19.84 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.1 (20.00-2.50) 97.0 (19.84-2.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.76 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.237 , 0.266 0.288 , 0.317	Depositor DCC
$R_{free}$ test set	750 reflections (4.96%)	DCC
Wilson B-factor (Å <sup>2</sup> )	57.2	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 15.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 15162 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2485	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/2472	0.73	4/3372 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	LEU	CA-CB-CG	7.19	131.84	115.30
1	A	273	LEU	CA-CB-CG	5.93	128.94	115.30
1	A	221	LEU	CA-CB-CG	5.92	128.91	115.30
1	A	24	CYS	N-CA-C	5.39	125.56	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2408	0	2403	39	3
2	A	43	0	31	4	0
3	A	34	0	0	4	0
All	All	2485	0	2434	39	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (39) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:172:ARG:HD2	1:A:174:MSE:HE2	1.56	0.87
1:A:42:LEU:HD22	1:A:54:MET:HG3	1.61	0.80
1:A:8[B]:PHE:HE1	1:A:211:GLN:HB2	1.52	0.74
1:A:242:THR:HG23	1:A:244:ALA:H	1.54	0.72
1:A:275:VAL:HG13	1:A:287:ILE:HB	1.79	0.65
1:A:170:PHE:HD1	1:A:291:HIS:HB2	1.63	0.63
1:A:9:PHE:HE1	1:A:23:ASN:HB2	1.67	0.59
1:A:287:ILE:HD11	1:A:312:LEU:HD23	1.84	0.58
1:A:176:TYR:CZ	1:A:184:GLY:HA3	2.38	0.58
1:A:51:VAL:HG23	1:A:70:ARG:HA	1.85	0.58
1:A:145:PHE:HA	1:A:148:GLU:HB2	1.85	0.58
1:A:179:ALA:HB1	1:A:180:PRO:HD2	1.87	0.57
1:A:170:PHE:CD1	1:A:291:HIS:HB2	2.43	0.54
1:A:104:ASP:HB3	1:A:110:HIS:H	1.72	0.53
1:A:174:MSE:HE1	1:A:290:HIS:CE1	2.43	0.53
1:A:9:PHE:CE1	1:A:23:ASN:HB2	2.44	0.52
1:A:216:TYR:HA	3:A:361:HOH:O	2.11	0.50
1:A:274:SER:OG	2:A:328:BLA:HBA1	2.10	0.50
1:A:290:HIS:CE1	2:A:328:BLA:OB	2.65	0.50
1:A:322:HIS:CG	1:A:322:HIS:O	2.66	0.49
1:A:142:ASN:C	1:A:144:MSE:H	2.16	0.49
1:A:23:ASN:O	1:A:26:ARG:HG2	2.13	0.48
1:A:322:HIS:C	1:A:324:HIS:H	2.17	0.48
1:A:287:ILE:HD11	1:A:312:LEU:CD2	2.44	0.48
1:A:239:ASN:HB3	1:A:242:THR:HG22	1.96	0.47
1:A:42:LEU:CD2	1:A:54:MET:HG3	2.37	0.47
1:A:140:LEU:HD21	1:A:307:TYR:CE1	2.50	0.47
1:A:43:THR:OG1	1:A:53:GLN:HB2	2.16	0.46
1:A:310:ARG:HG3	3:A:360:HOH:O	2.16	0.45
1:A:219:HIS:C	3:A:361:HOH:O	2.54	0.45
1:A:274:SER:OG	2:A:328:BLA:CBA	2.65	0.44
1:A:9:PHE:CE1	1:A:26:ARG:HG3	2.53	0.44
1:A:239:ASN:OD1	1:A:241:GLN:HB2	2.18	0.44
1:A:223:LEU:HB3	1:A:273:LEU:HD22	2.00	0.43
1:A:29:ILE:HG21	2:A:328:BLA:HMD1	2.00	0.43
1:A:322:HIS:C	1:A:324:HIS:N	2.73	0.42
1:A:68:VAL:HG13	1:A:68:VAL:O	2.20	0.41
1:A:226:ASP:HB3	1:A:229:ALA:HB2	2.03	0.41
1:A:216:TYR:C	3:A:361:HOH:O	2.60	0.40



All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:307:TYR:CD2	1:A:307:TYR:CE2[2.665]	1.86	0.34
1:A:307:TYR:CE2	1:A:307:TYR:CZ[2.665]	2.00	0.20
1:A:307:TYR:CD2	1:A:307:TYR:CZ[2.665]	2.14	0.06

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	308/341 (90%)	288 (94%)	14 (4%)	6 (2%)	12 19

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	CYS
1	A	110	HIS
1	A	66	PRO
1	A	143	ALA
1	A	119	GLY
1	A	323	HIS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	255/264 (97%)	225 (88%)	30 (12%)	8 14

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	26	ARG
1	A	56	LEU
1	A	72	GLN
1	A	74	LEU
1	A	95	ASP
1	A	97	LEU
1	A	103	LEU
1	A	104	ASP
1	A	111	LEU
1	A	121	LEU
1	A	141	ARG
1	A	144	MSE
1	A	154	ARG
1	A	167	LEU
1	A	177	LYS
1	A	199	LEU
1	A	206	SER
1	A	220	LEU
1	A	222	ARG
1	A	223	LEU
1	A	232	VAL
1	A	238	LEU
1	A	273	LEU
1	A	275	VAL
1	A	297	LEU
1	A	310	ARG
1	A	313	SER
1	A	319	LYS
1	A	322	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BLA	A	328	1	46,46,46	3.82	15 (32%)	65,67,67	2.35	23 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLA	A	328	1	-	1/26/74/74	0/4/4/4

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	328	BLA	CHA-C4D	19.86	1.48	1.35
2	A	328	BLA	CHB-C1B	7.10	1.50	1.34
2	A	328	BLA	C3C-C2C	5.37	1.47	1.36
2	A	328	BLA	C3D-C2D	5.24	1.48	1.36
2	A	328	BLA	C3B-C2B	5.24	1.46	1.36
2	A	328	BLA	CBC-CAC	4.49	1.52	1.29
2	A	328	BLA	CHD-C4C	4.35	1.48	1.37
2	A	328	BLA	C1A-NA	-4.09	1.32	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	328	BLA	C2A-C3A	3.92	1.49	1.37
2	A	328	BLA	CHD-C1D	3.49	1.48	1.40
2	A	328	BLA	C4A-C3A	3.34	1.50	1.42
2	A	328	BLA	C1B-C2B	2.93	1.50	1.45
2	A	328	BLA	OC-C1C	2.46	1.28	1.23
2	A	328	BLA	OB-C4B	2.19	1.27	1.23
2	A	328	BLA	C1A-C2A	2.06	1.47	1.42

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	328	BLA	C3B-C4B-NB	8.55	114.39	106.66
2	A	328	BLA	C4C-NC-C1C	-4.67	104.14	110.74
2	A	328	BLA	C1A-C2A-C3A	-4.32	102.45	106.92
2	A	328	BLA	C2A-C1A-NA	4.30	113.89	106.79
2	A	328	BLA	C4B-C3B-C2B	-4.26	102.51	107.97
2	A	328	BLA	C4A-C3A-C2A	-4.18	104.19	107.04
2	A	328	BLA	C3C-C4C-NC	3.63	113.89	106.59
2	A	328	BLA	C2C-C1C-NC	3.47	116.25	106.27
2	A	328	BLA	CMC-C2C-C1C	3.46	130.55	121.59
2	A	328	BLA	C4D-C3D-C2D	-3.13	103.22	106.86
2	A	328	BLA	CMB-C2B-C1B	2.97	128.29	124.23
2	A	328	BLA	OB-C4B-C3B	-2.86	122.58	130.22
2	A	328	BLA	C3D-C4D-ND	2.85	114.30	109.79
2	A	328	BLA	C1D-C2D-C3D	-2.61	103.41	106.64
2	A	328	BLA	C1A-CHA-C4D	2.58	134.97	129.92
2	A	328	BLA	CHB-C1B-C2B	-2.42	121.61	126.83
2	A	328	BLA	C1B-NB-C4B	-2.40	107.35	110.74
2	A	328	BLA	CBC-CAC-C3C	-2.30	115.68	127.09
2	A	328	BLA	O2A-CGA-CBA	2.26	122.19	114.22
2	A	328	BLA	CAA-C2A-C3A	2.18	135.22	129.00
2	A	328	BLA	CAB-C3B-C4B	2.09	130.86	123.12
2	A	328	BLA	CHD-C4C-NC	-2.07	121.70	126.19
2	A	328	BLA	C2B-C1B-NB	2.03	110.35	107.14

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	328	BLA	C1B-CHB-C4A-NA

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	313/341 (91%)	1.13	58 (18%) 2 2	37, 47, 64, 92	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	67	THR	9.1
1	A	68	VAL	8.7
1	A	6	LEU	7.3
1	A	109	GLY	6.8
1	A	105	TRP	6.2
1	A	63	GLY	5.7
1	A	325	HIS	5.3
1	A	138	HIS	5.3
1	A	65	GLU	5.1
1	A	324	HIS	5.1
1	A	71	GLY	4.8
1	A	106	PRO	4.5
1	A	288	ALA	4.1
1	A	66	PRO	4.0
1	A	289	CYS	3.9
1	A	104	ASP	3.5
1	A	124	LEU	3.5
1	A	110	HIS	3.4
1	A	175	LEU	3.4
1	A	130	GLU	3.4
1	A	308	LEU	3.3
1	A	129	THR	3.3
1	A	256	THR	3.2
1	A	47	HIS	3.0
1	A	286	LEU	3.0
1	A	102	THR	3.0
1	A	307	TYR	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	64	GLN	2.9
1	A	282	GLN	2.9
1	A	123	ILE	2.7
1	A	287	ILE	2.7
1	A	228	ARG	2.7
1	A	145	PHE	2.7
1	A	62	LEU	2.6
1	A	322	HIS	2.6
1	A	312	LEU	2.6
1	A	5	PRO	2.6
1	A	80	GLU	2.6
1	A	56	LEU	2.5
1	A	59	ALA	2.5
1	A	305	LEU	2.4
1	A	70	ARG	2.4
1	A	259	MSE	2.4
1	A	115	VAL	2.4
1	A	8[A]	PHE	2.3
1	A	193	GLU	2.3
1	A	72	GLN	2.3
1	A	301	LEU	2.2
1	A	88	ALA	2.2
1	A	42	LEU	2.2
1	A	152	ASN	2.2
1	A	323	HIS	2.2
1	A	218	ARG	2.2
1	A	300	ASP	2.2
1	A	208	ILE	2.2
1	A	144	MSE	2.1
1	A	255	ALA	2.2
1	A	41	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BLA	A	328	43/43	0.21	-0.37	19,33,34,40	0

## 6.5 Other polymers

There are no such residues in this entry.