



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 03:24 AM GMT

PDB ID : 1SQL
Title : Crystal structure of 7,8-dihydroneopterin aldolase in complex with guanine
Authors : Bauer, S.; Schott, A.K.; Illarionova, V.; Bacher, A.; Huber, R.; Fischer, M.
Deposited on : 2004-03-19
Resolution : 2.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2938 (2.20-2.20)
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	146	
1	B	146	
1	C	146	
1	D	146	
1	E	146	
1	F	146	
1	G	146	
1	H	146	
1	I	146	
1	J	146	
1	K	146	
1	L	146	
1	M	146	
1	N	146	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	O	146	
1	P	146	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	GUN	A	1001	X	-
2	GUN	B	1002	X	-
2	GUN	B	1003	X	-
2	GUN	D	1004	X	-
2	GUN	F	1006	X	-
2	GUN	G	1007	X	-
2	GUN	G	1008	X	-
2	GUN	H	1005	X	-
2	GUN	I	1009	X	-
2	GUN	J	1010	X	-
2	GUN	J	1011	X	-
2	GUN	L	1012	X	-
2	GUN	M	1014	X	-
2	GUN	O	1015	X	-
2	GUN	P	1013	X	-
2	GUN	P	1016	X	-

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 16400 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dihydroneopterin aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	120	Total	C	N	O	S	5	0	0
			947	611	158	177	1			
1	B	121	Total	C	N	O	S	0	0	0
			955	615	160	179	1			
1	C	122	Total	C	N	O	S	7	0	0
			962	619	161	181	1			
1	D	120	Total	C	N	O	S	0	0	0
			947	611	158	177	1			
1	E	120	Total	C	N	O	S	0	0	0
			947	611	158	177	1			
1	F	121	Total	C	N	O	S	0	0	0
			955	615	160	179	1			
1	G	121	Total	C	N	O	S	6	0	0
			955	615	160	179	1			
1	H	122	Total	C	N	O	S	0	0	0
			962	619	161	181	1			
1	I	121	Total	C	N	O	S	6	0	0
			955	615	160	179	1			
1	J	123	Total	C	N	O	S	0	0	0
			968	622	162	183	1			
1	K	120	Total	C	N	O	S	0	0	0
			947	611	158	177	1			
1	L	122	Total	C	N	O	S	0	0	0
			962	619	161	181	1			
1	M	123	Total	C	N	O	S	7	0	0
			968	622	162	183	1			
1	N	121	Total	C	N	O	S	0	0	0
			955	615	160	179	1			
1	O	120	Total	C	N	O	S	0	0	0
			947	611	158	177	1			
1	P	122	Total	C	N	O	S	0	0	0
			963	621	161	180	1			

- Molecule 2 is GUANINE (three-letter code: GUN) (formula: $C_5H_5N_5O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			11	5	5	1		
2	B	1	Total	C	N	O	0	0
			11	5	5	1		
2	B	1	Total	C	N	O	0	0
			11	5	5	1		
2	D	1	Total	C	N	O	0	0
			11	5	5	1		
2	H	1	Total	C	N	O	0	0
			11	5	5	1		
2	F	1	Total	C	N	O	0	0
			11	5	5	1		
2	G	1	Total	C	N	O	0	0
			11	5	5	1		
2	G	1	Total	C	N	O	0	0
			11	5	5	1		
2	I	1	Total	C	N	O	0	0
			11	5	5	1		
2	J	1	Total	C	N	O	0	0
			11	5	5	1		
2	J	1	Total	C	N	O	0	0
			11	5	5	1		
2	L	1	Total	C	N	O	0	0
			11	5	5	1		
2	P	1	Total	C	N	O	0	0
			11	5	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	M	1	Total	C	N	O	0	0
			11	5	5	1		
2	O	1	Total	C	N	O	0	0
			11	5	5	1		
2	P	1	Total	C	N	O	0	0
			11	5	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	59	Total	O	0	0
			59	59		
3	B	67	Total	O	0	0
			67	67		
3	C	51	Total	O	0	0
			51	51		
3	D	50	Total	O	0	0
			50	50		
3	E	52	Total	O	0	0
			52	52		
3	F	52	Total	O	0	0
			52	52		
3	G	69	Total	O	0	0
			69	69		
3	H	67	Total	O	0	0
			67	67		
3	I	60	Total	O	0	0
			60	60		
3	J	70	Total	O	0	0
			70	70		
3	K	56	Total	O	0	0
			56	56		
3	L	54	Total	O	0	0
			54	54		
3	M	50	Total	O	0	0
			50	50		
3	N	53	Total	O	0	0
			53	53		
3	O	60	Total	O	0	0
			60	60		
3	P	59	Total	O	0	0
			59	59		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: dihydroneopterin aldolase




- Molecule 1: dihydroneopterin aldolase



- Molecule 1: dihydroneopterin aldolase




- Molecule 1: dihydroneopterin aldolase




- Molecule 1: dihydroneopterin aldolase

Chain E:




- Molecule 1: dihydroneopterin aldolase

Chain F:




- Molecule 1: dihydroneopterin aldolase

Chain G:




- Molecule 1: dihydroneopterin aldolase

Chain H:



- Molecule 1: dihydroneopterin aldolase

Chain I:



- Molecule 1: dihydroneopterin aldolase


Chain J:





- Molecule 1: dihydroneopterin aldolase

Chain K:



- Molecule 1: dihydroneopterin aldolase

Chain L:



- Molecule 1: dihydroneopterin aldolase

Chain M:




- Molecule 1: dihydroneopterin aldolase

Chain N:




- Molecule 1: dihydroneopterin aldolase

Chain O:



- Molecule 1: dihydroneopterin aldolase

Chain P:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	63.55Å 84.22Å 89.05Å 90.14° 89.99° 76.17°	Depositor
Resolution (Å)	20.00 – 2.20 15.94 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.20) 94.9 (15.94-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.90 (at 2.20Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.204 , 0.260 0.182 , 0.212	Depositor DCC
R_{free} test set	3837 reflections (4.63%)	DCC
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.352	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 38.8	EDS
Estimated twinning fraction	0.477 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 86786 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16400	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GUN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/961	0.68	0/1294
1	B	0.39	0/969	0.66	0/1305
1	C	0.37	0/976	0.64	0/1315
1	D	0.40	0/961	0.64	0/1294
1	E	0.40	0/961	0.67	0/1294
1	F	0.39	0/969	0.64	0/1305
1	G	0.40	0/969	0.65	0/1305
1	H	0.38	0/976	0.64	0/1315
1	I	0.38	0/969	0.65	0/1305
1	J	0.38	0/982	0.65	0/1323
1	K	0.39	0/961	0.66	0/1294
1	L	0.39	0/976	0.65	0/1315
1	M	0.37	0/982	0.66	0/1323
1	N	0.39	0/969	0.63	0/1305
1	O	0.38	0/961	0.68	0/1294
1	P	0.39	0/977	0.66	0/1316
All	All	0.39	0/15519	0.65	0/20902

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	947	0	983	30	0
1	B	955	0	989	43	0
1	C	962	0	996	37	0
1	D	947	0	983	40	0
1	E	947	0	983	48	0
1	F	955	0	989	38	0
1	G	955	0	989	33	0
1	H	962	0	996	21	0
1	I	955	0	989	32	0
1	J	968	0	1001	36	0
1	K	947	0	983	26	0
1	L	962	0	996	41	0
1	M	968	0	1001	43	0
1	N	955	0	989	31	0
1	O	947	0	983	31	0
1	P	963	0	1000	36	0
2	A	11	0	5	0	0
2	B	22	0	10	1	0
2	D	11	0	5	0	0
2	F	11	0	5	1	0
2	G	22	0	10	1	0
2	H	11	0	5	0	0
2	I	11	0	5	2	0
2	J	22	0	10	0	0
2	L	11	0	5	1	0
2	M	11	0	5	0	0
2	O	11	0	5	0	0
2	P	22	0	10	2	0
3	A	59	0	0	3	0
3	B	67	0	0	2	0
3	C	51	0	0	2	0
3	D	50	0	0	1	0
3	E	52	0	0	2	0
3	F	52	0	0	1	0
3	G	69	0	0	4	0
3	H	67	0	0	1	0
3	I	60	0	0	2	0
3	J	70	0	0	3	0
3	K	56	0	0	1	0
3	L	54	0	0	0	0
3	M	50	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	N	53	0	0	6	0
3	O	60	0	0	4	0
3	P	59	0	0	1	0
All	All	16400	0	15930	533	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 17.

The worst 5 of 533 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:L:88:HIS:HA	1:L:120:ARG:NH1	1.80	0.95
1:J:88:HIS:CD2	1:J:88:HIS:H	1.64	0.95
1:L:88:HIS:HA	1:L:120:ARG:HH11	1.30	0.93
1:N:46:ASN:HB2	3:N:171:HOH:O	1.69	0.93
1:J:88:HIS:H	1:J:88:HIS:HD2	1.10	0.90

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/146 (81%)	115 (98%)	3 (2%)	0	100	100
1	B	119/146 (82%)	116 (98%)	3 (2%)	0	100	100
1	C	120/146 (82%)	116 (97%)	4 (3%)	0	100	100
1	D	118/146 (81%)	113 (96%)	5 (4%)	0	100	100
1	E	118/146 (81%)	116 (98%)	2 (2%)	0	100	100
1	F	119/146 (82%)	114 (96%)	5 (4%)	0	100	100
1	G	119/146 (82%)	115 (97%)	4 (3%)	0	100	100
1	H	120/146 (82%)	117 (98%)	3 (2%)	0	100	100
1	I	119/146 (82%)	118 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	J	121/146 (83%)	119 (98%)	2 (2%)	0	100	100
1	K	118/146 (81%)	114 (97%)	4 (3%)	0	100	100
1	L	120/146 (82%)	116 (97%)	4 (3%)	0	100	100
1	M	121/146 (83%)	115 (95%)	6 (5%)	0	100	100
1	N	119/146 (82%)	114 (96%)	5 (4%)	0	100	100
1	O	118/146 (81%)	117 (99%)	1 (1%)	0	100	100
1	P	120/146 (82%)	116 (97%)	4 (3%)	0	100	100
All	All	1907/2336 (82%)	1851 (97%)	56 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/127 (81%)	96 (93%)	7 (7%)	22	23
1	B	104/127 (82%)	99 (95%)	5 (5%)	35	41
1	C	105/127 (83%)	102 (97%)	3 (3%)	55	66
1	D	103/127 (81%)	98 (95%)	5 (5%)	35	40
1	E	103/127 (81%)	95 (92%)	8 (8%)	18	17
1	F	104/127 (82%)	99 (95%)	5 (5%)	35	41
1	G	104/127 (82%)	98 (94%)	6 (6%)	28	31
1	H	105/127 (83%)	99 (94%)	6 (6%)	29	32
1	I	104/127 (82%)	102 (98%)	2 (2%)	69	81
1	J	106/127 (84%)	99 (93%)	7 (7%)	24	25
1	K	103/127 (81%)	96 (93%)	7 (7%)	22	23
1	L	105/127 (83%)	98 (93%)	7 (7%)	23	24
1	M	106/127 (84%)	101 (95%)	5 (5%)	36	42
1	N	104/127 (82%)	97 (93%)	7 (7%)	23	24
1	O	103/127 (81%)	97 (94%)	6 (6%)	28	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	105/127 (83%)	96 (91%)	9 (9%)	15	14
All	All	1667/2032 (82%)	1572 (94%)	95 (6%)	29	32

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	88	HIS
1	J	109	ILE
1	P	9	LEU
1	H	109	ILE
1	J	47	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	121	ASN
1	J	101	ASN
1	O	91	ASN
1	J	88	HIS
1	D	89	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GUN	A	1001	-	12,12,12	3.96	5 (41%)	14,17,17	12.03	9 (64%)
2	GUN	B	1002	-	12,12,12	4.15	5 (41%)	14,17,17	12.27	9 (64%)
2	GUN	B	1003	-	12,12,12	3.93	4 (33%)	14,17,17	12.52	9 (64%)
2	GUN	D	1004	-	12,12,12	3.93	5 (41%)	14,17,17	11.90	9 (64%)
2	GUN	F	1006	-	12,12,12	3.95	5 (41%)	14,17,17	11.97	9 (64%)
2	GUN	G	1007	-	12,12,12	3.96	5 (41%)	14,17,17	11.10	9 (64%)
2	GUN	G	1008	-	12,12,12	3.76	5 (41%)	14,17,17	11.97	9 (64%)
2	GUN	H	1005	-	12,12,12	4.12	5 (41%)	14,17,17	11.91	9 (64%)
2	GUN	I	1009	-	12,12,12	3.79	5 (41%)	14,17,17	12.21	9 (64%)
2	GUN	J	1010	-	12,12,12	3.85	5 (41%)	14,17,17	11.60	9 (64%)
2	GUN	J	1011	-	12,12,12	3.96	5 (41%)	14,17,17	12.02	9 (64%)
2	GUN	L	1012	-	12,12,12	3.82	5 (41%)	14,17,17	11.95	9 (64%)
2	GUN	M	1014	-	12,12,12	4.04	5 (41%)	14,17,17	12.08	9 (64%)
2	GUN	O	1015	-	12,12,12	3.95	5 (41%)	14,17,17	11.57	9 (64%)
2	GUN	P	1013	-	12,12,12	3.88	5 (41%)	14,17,17	12.14	9 (64%)
2	GUN	P	1016	-	12,12,12	4.11	5 (41%)	14,17,17	12.32	9 (64%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GUN	A	1001	-	-	0/0/0/0	0/0/2/2
2	GUN	B	1002	-	-	0/0/0/0	0/0/2/2
2	GUN	B	1003	-	-	0/0/0/0	0/0/2/2
2	GUN	D	1004	-	-	0/0/0/0	0/0/2/2
2	GUN	F	1006	-	-	0/0/0/0	0/0/2/2
2	GUN	G	1007	-	-	0/0/0/0	0/0/2/2
2	GUN	G	1008	-	-	0/0/0/0	0/0/2/2
2	GUN	H	1005	-	-	0/0/0/0	0/0/2/2
2	GUN	I	1009	-	-	0/0/0/0	0/0/2/2
2	GUN	J	1010	-	-	0/0/0/0	0/0/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GUN	J	1011	-	-	0/0/0/0	0/0/2/2
2	GUN	L	1012	-	-	0/0/0/0	0/0/2/2
2	GUN	M	1014	-	-	0/0/0/0	0/0/2/2
2	GUN	O	1015	-	-	0/0/0/0	0/0/2/2
2	GUN	P	1013	-	-	0/0/0/0	0/0/2/2
2	GUN	P	1016	-	-	0/0/0/0	0/0/2/2

The worst 5 of 79 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1002	GUN	C4-N3	12.29	1.47	1.34
2	P	1016	GUN	C4-N3	12.04	1.47	1.34
2	H	1005	GUN	C4-N3	12.03	1.47	1.34
2	M	1014	GUN	C4-N3	11.99	1.47	1.34
2	A	1001	GUN	C4-N3	11.80	1.47	1.34

The worst 5 of 144 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1003	GUN	C6-C5-N7	43.66	140.02	134.14
2	P	1016	GUN	C6-C5-N7	42.78	139.90	134.14
2	B	1002	GUN	C6-C5-N7	42.72	139.89	134.14
2	I	1009	GUN	C6-C5-N7	42.47	139.86	134.14
2	P	1013	GUN	C6-C5-N7	42.20	139.82	134.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	120/146 (82%)	-0.45	0 100 100	11, 19, 33, 40	1 (0%)
1	B	121/146 (82%)	-0.42	0 100 100	12, 20, 33, 55	0
1	C	122/146 (83%)	-0.28	0 100 100	13, 23, 37, 59	1 (0%)
1	D	120/146 (82%)	-0.30	1 (0%) 83 85	12, 21, 37, 49	0
1	E	120/146 (82%)	-0.34	0 100 100	12, 20, 32, 42	0
1	F	121/146 (82%)	-0.39	0 100 100	9, 20, 35, 46	0
1	G	121/146 (82%)	-0.45	0 100 100	11, 19, 32, 57	1 (0%)
1	H	122/146 (83%)	-0.41	0 100 100	10, 19, 34, 40	0
1	I	121/146 (82%)	-0.45	0 100 100	11, 19, 30, 52	1 (0%)
1	J	123/146 (84%)	-0.39	0 100 100	12, 19, 35, 46	0
1	K	120/146 (82%)	-0.36	0 100 100	13, 21, 31, 45	0
1	L	122/146 (83%)	-0.30	1 (0%) 83 85	10, 21, 36, 66	0
1	M	123/146 (84%)	-0.22	3 (2%) 56 56	15, 23, 40, 68	1 (0%)
1	N	121/146 (82%)	-0.28	0 100 100	13, 21, 37, 60	0
1	O	120/146 (82%)	-0.47	0 100 100	11, 20, 33, 43	0
1	P	122/146 (83%)	-0.36	0 100 100	12, 20, 33, 56	0
All	All	1939/2336 (83%)	-0.37	5 (0%) 91 93	9, 20, 36, 68	5 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	121	ASN	2.8
1	M	35	TRP	2.4
1	D	35	TRP	2.3
1	M	123	SER	2.2
1	M	18	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GUN	J	1011	11/11	0.12	0.57	24,27,32,34	0
2	GUN	B	1002	11/11	0.11	0.47	16,17,23,26	0
2	GUN	G	1007	11/11	0.10	-0.10	10,14,17,20	0
2	GUN	I	1009	11/11	0.11	-0.19	16,20,22,26	0
2	GUN	P	1013	11/11	0.12	-0.22	18,23,26,28	0
2	GUN	G	1008	11/11	0.10	-0.58	8,10,13,14	0
2	GUN	F	1006	11/11	0.11	-0.62	12,16,18,22	0
2	GUN	B	1003	11/11	0.10	-0.68	16,23,26,28	0
2	GUN	D	1004	11/11	0.10	-0.72	16,20,26,27	0
2	GUN	A	1001	11/11	0.10	-0.75	18,21,25,25	0
2	GUN	P	1016	11/11	0.10	-0.78	15,17,22,24	0
2	GUN	M	1014	11/11	0.10	-0.82	13,19,26,27	0
2	GUN	O	1015	11/11	0.10	-0.89	13,16,18,21	0
2	GUN	H	1005	11/11	0.10	-0.99	21,23,25,25	0
2	GUN	L	1012	11/11	0.10	-1.06	15,18,22,23	0
2	GUN	J	1010	11/11	0.06	-2.74	5,14,19,20	0

6.5 Other polymers ⓘ

There are no such residues in this entry.