



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:17 AM GMT

PDB ID : 3A6J
Title : E122Q mutant creatininase complexed with creatine
Authors : Nakajima, Y.; Yamashita, K.; Ito, K.; Yoshimoto, T.
Deposited on : 2009-09-02
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	<div> <div>2%</div> <div>88%</div> <div>11%</div> <div>.</div> </div>
1	B	260	<div> <div>%</div> <div>88%</div> <div>10%</div> <div>.</div> </div>
1	C	260	<div> <div>2%</div> <div>88%</div> <div>10%</div> <div>.</div> </div>
1	D	260	<div> <div>3%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	E	260	<div> <div>3%</div> <div>82%</div> <div>16%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	260	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	E	301	-	-	-	X
3	CRN	A	303	-	-	X	X
3	CRN	B	304	-	-	-	X
3	CRN	C	305	-	-	-	X
3	CRN	E	306	-	-	X	X
3	CRN	F	307	-	-	-	X
4	SO4	A	403	-	-	-	X
4	SO4	B	416	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Creatinine amidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			1981	1269	337	364	11			
1	B	254	Total	C	N	O	S	0	0	0
			1971	1264	335	361	11			
1	C	255	Total	C	N	O	S	0	0	0
			1975	1266	336	362	11			
1	D	256	Total	C	N	O	S	0	0	0
			1980	1269	337	363	11			
1	E	255	Total	C	N	O	S	0	0	0
			1976	1267	336	362	11			
1	F	255	Total	C	N	O	S	0	0	0
			1971	1263	335	362	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	GLN	GLU	ENGINEERED	UNP P83772
B	122	GLN	GLU	ENGINEERED	UNP P83772
C	122	GLN	GLU	ENGINEERED	UNP P83772
D	122	GLN	GLU	ENGINEERED	UNP P83772
E	122	GLN	GLU	ENGINEERED	UNP P83772
F	122	GLN	GLU	ENGINEERED	UNP P83772

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Zn	0	0
			1	1		
2	E	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	F	1	Total	Zn	0	0
			1	1		

- Molecule 3 is N-[(E)-AMINO(IMINO)METHYL]-N-METHYLGLYCINE (three-letter code: CRN) (formula: C₄H₉N₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	4	3	2		
3	B	1	Total	C	N	O	0	0
			9	4	3	2		
3	C	1	Total	C	N	O	0	0
			9	4	3	2		
3	E	1	Total	C	N	O	0	0
			9	4	3	2		
3	F	1	Total	C	N	O	0	0
			9	4	3	2		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	154	Total 154	O 154	0	0
5	B	147	Total 147	O 147	0	0
5	C	137	Total 137	O 137	0	0
5	D	141	Total 141	O 141	0	0
5	E	129	Total 129	O 129	0	0
5	F	158	Total 158	O 158	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Creatinine amidohydrolase




- Molecule 1: Creatinine amidohydrolase




- Molecule 1: Creatinine amidohydrolase




- Molecule 1: Creatinine amidohydrolase



● Molecule 1: Creatinine amidohydrolase



● Molecule 1: Creatinine amidohydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	102.20 Å 152.70 Å 166.90 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 49.33 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.00) 99.8 (49.33-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.58 (at 2.00 Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.195 , 0.212 0.195 , 0.212	Depositor DCC
R_{free} test set	8777 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.494	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 175544 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12836	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CRN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/2030	0.59	0/2764
1	B	0.30	0/2019	0.58	0/2746
1	C	0.30	0/2023	0.58	0/2751
1	D	0.30	0/2028	0.57	0/2758
1	E	0.29	0/2024	0.58	0/2753
1	F	0.30	0/2019	0.58	0/2747
All	All	0.30	0/12143	0.58	0/16519

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1981	0	1958	20	0
1	B	1971	0	1955	20	0
1	C	1975	0	1958	17	0
1	D	1980	0	1963	26	0
1	E	1976	0	1963	30	0
1	F	1971	0	1947	15	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	9	0	7	5	0
3	B	9	0	7	3	0
3	C	9	0	7	2	0
3	E	9	0	7	4	0
3	F	9	0	7	1	0
4	A	15	0	0	0	0
4	B	10	0	0	2	0
4	C	10	0	0	0	0
4	D	10	0	0	0	0
4	E	10	0	0	0	0
4	F	10	0	0	1	0
5	A	154	0	0	2	0
5	B	147	0	0	0	0
5	C	137	0	0	0	0
5	D	141	0	0	2	0
5	E	129	0	0	0	0
5	F	158	0	0	0	0
All	All	12836	0	11779	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:172:LEU:HB3	1:D:176:ILE:HD11	1.55	0.89
1:E:172:LEU:HB3	1:E:176:ILE:HD11	1.57	0.86
1:B:172:LEU:HB3	1:B:176:ILE:HD11	1.62	0.79
1:E:178:HIS:HB2	3:E:306:CRN:N3	2.06	0.71
3:B:304:CRN:C7	3:B:304:CRN:HN31	2.06	0.69

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/260 (98%)	249 (98%)	6 (2%)	0	100	100
1	B	250/260 (96%)	243 (97%)	7 (3%)	0	100	100
1	C	251/260 (96%)	244 (97%)	7 (3%)	0	100	100
1	D	252/260 (97%)	245 (97%)	7 (3%)	0	100	100
1	E	251/260 (96%)	244 (97%)	6 (2%)	1 (0%)	39	33
1	F	251/260 (96%)	245 (98%)	6 (2%)	0	100	100
All	All	1510/1560 (97%)	1470 (97%)	39 (3%)	1 (0%)	56	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	79	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/215 (98%)	208 (99%)	2 (1%)	82	85
1	B	211/215 (98%)	208 (99%)	3 (1%)	74	77
1	C	211/215 (98%)	209 (99%)	2 (1%)	84	88
1	D	211/215 (98%)	208 (99%)	3 (1%)	74	77
1	E	211/215 (98%)	209 (99%)	2 (1%)	84	88
1	F	210/215 (98%)	207 (99%)	3 (1%)	74	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	1264/1290 (98%)	1249 (99%)	15 (1%)	78	81

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	122	GLN
1	D	43	ASN
1	F	43	ASN
1	C	43	ASN
1	E	122	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	75	GLN
1	D	144	GLN
1	F	122	GLN
1	D	118	ASN
1	D	122	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 6 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CRN	A	303	2	2,8,8	1.10	0	2,10,10	2.45	1 (50%)
4	SO4	A	403	-	4,4,4	0.24	0	6,6,6	0.07	0
4	SO4	A	409	-	4,4,4	0.20	0	6,6,6	0.09	0
4	SO4	A	415	-	4,4,4	0.22	0	6,6,6	0.10	0
3	CRN	B	304	2	2,8,8	1.97	1 (50%)	2,10,10	2.36	2 (100%)
4	SO4	B	410	-	4,4,4	0.19	0	6,6,6	0.13	0
4	SO4	B	416	-	4,4,4	0.22	0	6,6,6	0.08	0
3	CRN	C	305	2	2,8,8	0.60	0	2,10,10	2.41	1 (50%)
4	SO4	C	411	-	4,4,4	0.21	0	6,6,6	0.06	0
4	SO4	C	417	-	4,4,4	0.23	0	6,6,6	0.08	0
4	SO4	D	400	-	4,4,4	0.23	0	6,6,6	0.08	0
4	SO4	D	412	-	4,4,4	0.18	0	6,6,6	0.11	0
3	CRN	E	306	2	2,8,8	1.08	0	2,10,10	2.51	1 (50%)
4	SO4	E	401	-	4,4,4	0.21	0	6,6,6	0.08	0
4	SO4	E	413	-	4,4,4	0.20	0	6,6,6	0.15	0
3	CRN	F	307	2	2,8,8	1.15	0	2,10,10	2.59	1 (50%)
4	SO4	F	402	-	4,4,4	0.23	0	6,6,6	0.14	0
4	SO4	F	414	-	4,4,4	0.20	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CRN	A	303	2	-	0/2/8/8	0/0/0/0
4	SO4	A	403	-	-	0/0/0/0	0/0/0/0
4	SO4	A	409	-	-	0/0/0/0	0/0/0/0
4	SO4	A	415	-	-	0/0/0/0	0/0/0/0
3	CRN	B	304	2	-	0/2/8/8	0/0/0/0
4	SO4	B	410	-	-	0/0/0/0	0/0/0/0
4	SO4	B	416	-	-	0/0/0/0	0/0/0/0
3	CRN	C	305	2	-	0/2/8/8	0/0/0/0
4	SO4	C	411	-	-	0/0/0/0	0/0/0/0
4	SO4	C	417	-	-	0/0/0/0	0/0/0/0
4	SO4	D	400	-	-	0/0/0/0	0/0/0/0
4	SO4	D	412	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CRN	E	306	2	-	0/2/8/8	0/0/0/0
4	SO4	E	401	-	-	0/0/0/0	0/0/0/0
4	SO4	E	413	-	-	0/0/0/0	0/0/0/0
3	CRN	F	307	2	-	0/2/8/8	0/0/0/0
4	SO4	F	402	-	-	0/0/0/0	0/0/0/0
4	SO4	F	414	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	304	CRN	C5-N1	2.76	1.50	1.47

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	304	CRN	C4-N1-C5	2.10	112.87	110.04
3	B	304	CRN	C7-C5-N1	2.58	117.72	112.55
3	A	303	CRN	C4-N1-C5	2.90	113.94	110.04
3	E	306	CRN	C4-N1-C5	3.13	114.24	110.04
3	C	305	CRN	C4-N1-C5	3.17	114.30	110.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	CRN	5	0
3	B	304	CRN	3	0
4	B	416	SO4	2	0
3	C	305	CRN	2	0
3	E	306	CRN	4	0
3	F	307	CRN	1	0
4	F	402	SO4	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/260 (98%)	-0.26	4 (1%) 74 75	14, 21, 35, 48	0
1	B	254/260 (97%)	-0.07	2 (0%) 87 88	14, 23, 39, 45	0
1	C	255/260 (98%)	-0.18	5 (1%) 68 69	15, 24, 38, 45	0
1	D	256/260 (98%)	-0.01	8 (3%) 52 53	15, 25, 39, 43	0
1	E	255/260 (98%)	-0.03	7 (2%) 58 58	15, 25, 39, 49	0
1	F	255/260 (98%)	-0.28	1 (0%) 93 93	14, 22, 38, 47	0
All	All	1532/1560 (98%)	-0.14	27 (1%) 71 72	14, 24, 38, 49	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	21	ALA	6.7
1	A	22	GLY	6.2
1	E	20	ALA	5.8
1	A	20	ALA	5.0
1	E	21	ALA	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	A	403	5/5	0.84	0.32	18.48	53,54,57,59	0
3	CRN	B	304	9/9	0.69	0.34	14.31	35,39,41,41	0
3	CRN	C	305	9/9	0.76	0.29	10.78	34,39,41,42	0
3	CRN	F	307	9/9	0.80	0.23	10.12	32,37,41,44	0
3	CRN	E	306	9/9	0.56	0.46	9.31	37,44,48,48	0
3	CRN	A	303	9/9	0.78	0.38	8.01	33,39,41,42	0
2	ZN	E	301	1/1	0.99	0.10	2.73	23,23,23,23	0
2	ZN	D	301	1/1	0.99	0.09	1.57	23,23,23,23	0
2	ZN	F	301	1/1	1.00	0.10	1.11	21,21,21,21	0
2	ZN	B	301	1/1	1.00	0.09	1.09	22,22,22,22	0
2	ZN	C	301	1/1	1.00	0.10	0.72	22,22,22,22	0
2	ZN	A	301	1/1	1.00	0.07	-1.33	19,19,19,19	0
4	SO4	D	412	5/5	0.98	0.08	-	40,42,43,43	0
4	SO4	A	415	5/5	0.96	0.13	-	55,55,56,57	0
4	SO4	E	401	5/5	0.96	0.14	-	57,57,58,59	0
4	SO4	D	400	5/5	0.93	0.20	-	59,60,60,61	0
4	SO4	B	410	5/5	0.99	0.08	-	36,39,39,40	0
4	SO4	E	413	5/5	0.96	0.10	-	39,40,42,42	0
4	SO4	A	409	5/5	0.98	0.07	-	28,32,33,33	0
4	SO4	F	414	5/5	0.98	0.08	-	35,37,39,39	0
4	SO4	C	417	5/5	0.92	0.23	-	59,59,60,61	0
4	SO4	F	402	5/5	0.92	0.20	-	64,64,65,67	0
4	SO4	C	411	5/5	0.98	0.08	-	33,34,36,37	0
4	SO4	B	416	5/5	0.92	0.18	-	58,58,60,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.