



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:23 AM GMT

PDB ID : 2AAI  
Title : Crystallographic refinement of ricin to 2.5 Angstroms  
Authors : Rutenber, E.; Katzin, B.J.; Montfort, W.; Villafranca, J.E.; Ernst, S.R.;  
Collins, E.J.; Mlsna, D.; Monzingo, A.F.; Ready, M.P.; Robertus, J.D.  
Deposited on : 1993-09-07  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	267	
2	B	262	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NDG	B	270	X	-	-	-
4	NDG	B	280	X	-	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4440 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RICIN (A CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2114	1342	372	395	5			

- Molecule 2 is a protein called RICIN (B CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	262	Total	C	N	O	S	0	0	0
			2035	1273	357	393	12			

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	2	Total	C	O	0	0
			23	12	11		
3	B	2	Total	C	O	0	0
			23	12	11		

- Molecule 4 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	5	Total	C	N	O	0	0
			61	34	2	25		
4	B	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	61	Total	O	0	0
			61	61		
5	B	62	Total	O	0	0
			62	62		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: RICIN (A CHAIN)



#### • Molecule 2: RICIN (B CHAIN)



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.74Å 78.49Å 114.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.50	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.50)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.212 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4440	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NAG, NDG, GAL, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.04	0/2162	2.00	89/2941 (3.0%)
2	B	1.04	2/2080 (0.1%)	2.12	86/2842 (3.0%)
All	All	1.04	2/4242 (0.0%)	2.06	175/5783 (3.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
2	B	0	3
4	B	2	0
All	All	2	6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	131	TRP	CD1-NE1	-5.74	1.28	1.38
2	B	195	SER	CA-CB	5.14	1.60	1.52

The worst 5 of 175 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	200	THR	CA-C-N	-13.18	88.21	117.20
1	A	234	ARG	NE-CZ-NH2	-13.11	113.74	120.30
1	A	234	ARG	NE-CZ-NH1	11.23	125.91	120.30
2	B	37	TRP	CD1-CG-CD2	10.77	114.92	106.30
1	A	258	ARG	NE-CZ-NH1	10.62	125.61	120.30

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	270	NDG	C1
4	B	280	NDG	C1

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	158	GLY	Mainchain
1	A	262	PRO	Peptide
1	A	56	ARG	Sidechain
2	B	170	GLU	Mainchain
2	B	74	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	0	2083	63	0
2	B	2035	0	1974	51	0
3	B	46	0	42	0	0
4	B	122	0	104	1	0
5	A	61	0	0	2	0
5	B	62	0	0	4	0
All	All	4440	0	4203	112	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 112 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:ILE:HG23	1:A:2:PHE:H	1.51	0.76
1:A:236:ASN:HD21	1:A:238:SER:HB2	1.53	0.73
2:B:193:SER:HA	2:B:201:VAL:O	1.94	0.67
1:A:225:ALA:HB1	1:A:226:PHE:CD1	2.29	0.67
2:B:198:ARG:HB3	2:B:199:GLU:OE1	1.94	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/267 (99%)	231 (87%)	17 (6%)	17 (6%)	2	1
2	B	260/262 (99%)	222 (85%)	20 (8%)	18 (7%)	1	1
All	All	525/529 (99%)	453 (86%)	37 (7%)	35 (7%)	1	1

5 of 35 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	PRO
1	A	14	ALA
1	A	66	ALA
1	A	67	GLU
1	A	222	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/226 (100%)	184 (81%)	42 (19%)	2	3
2	B	227/227 (100%)	197 (87%)	30 (13%)	5	9
All	All	453/453 (100%)	381 (84%)	72 (16%)	3	5

5 of 72 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	232	LEU
1	A	263	PRO
2	B	225	LEU
1	A	235	ARG
1	A	239	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	233	GLN
1	A	236	ASN
2	B	196	ASN
1	A	209	ASN
2	B	189	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

14 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	GAL	B	264	3	11,11,12	1.04	1 (9%)	14,15,17	1.49	2 (14%)
3	BGC	B	265	3	12,12,12	1.27	1 (8%)	17,17,17	1.72	4 (23%)
3	GAL	B	267	3	11,11,12	1.11	1 (9%)	14,15,17	2.04	6 (42%)
3	BGC	B	268	3	12,12,12	1.21	0	17,17,17	2.15	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NDG	B	270	2,4	14,14,15	1.26	2 (14%)	15,19,21	1.53	2 (13%)
4	NAG	B	271	4	14,14,15	1.71	6 (42%)	15,19,21	3.35	5 (33%)
4	BMA	B	272	4	11,11,12	1.55	2 (18%)	14,15,17	4.11	8 (57%)
4	MAN	B	273	4	11,11,12	2.18	4 (36%)	14,15,17	2.40	8 (57%)
4	MAN	B	274	4	11,11,12	1.48	1 (9%)	14,15,17	1.08	1 (7%)
4	NDG	B	280	2,4	14,14,15	0.86	0	15,19,21	1.08	2 (13%)
4	NAG	B	281	4	14,14,15	1.06	1 (7%)	15,19,21	2.58	5 (33%)
4	BMA	B	282	4	11,11,12	1.81	2 (18%)	14,15,17	2.08	6 (42%)
4	MAN	B	283	4	11,11,12	1.38	0	14,15,17	4.00	9 (64%)
4	MAN	B	284	4	11,11,12	1.24	1 (9%)	14,15,17	3.12	12 (85%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GAL	B	264	3	-	0/2/19/22	0/1/1/1
3	BGC	B	265	3	-	0/2/22/22	0/1/1/1
3	GAL	B	267	3	-	0/2/19/22	0/1/1/1
3	BGC	B	268	3	-	0/2/22/22	0/1/1/1
4	NDG	B	270	2,4	1/1/5/7	0/6/23/26	0/1/1/1
4	NAG	B	271	4	-	0/6/23/26	0/1/1/1
4	BMA	B	272	4	-	0/2/19/22	0/1/1/1
4	MAN	B	273	4	-	0/2/19/22	0/1/1/1
4	MAN	B	274	4	-	0/2/19/22	0/1/1/1
4	NDG	B	280	2,4	1/1/5/7	0/6/23/26	0/1/1/1
4	NAG	B	281	4	-	1/6/23/26	0/1/1/1
4	BMA	B	282	4	-	0/2/19/22	0/1/1/1
4	MAN	B	283	4	-	0/2/19/22	1/1/1/1
4	MAN	B	284	4	-	0/2/19/22	1/1/1/1

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	274	MAN	C2-C3	-4.23	1.46	1.52
3	B	264	GAL	C2-C3	-2.69	1.48	1.52
4	B	270	NDG	O-C1	-2.42	1.39	1.43
4	B	281	NAG	O5-C1	-2.30	1.39	1.43
4	B	270	NDG	C1-C2	-2.17	1.49	1.52

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	B	283	MAN	O5-C1-C2	-10.85	93.25	110.86
4	B	272	BMA	C1-C2-C3	-10.63	96.96	109.54
4	B	271	NAG	C1-O5-C5	-8.51	101.44	112.25
4	B	281	NAG	C2-N2-C7	-7.88	112.92	123.04
3	B	268	BGC	C3-C4-C5	-6.74	98.45	110.20

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	280	NDG	C1
4	B	270	NDG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	281	NAG	O7-C7-N2-C2

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	283	MAN	C1-C2-C3-C4-C5-O5
4	B	284	MAN	C1-C2-C3-C4-C5-O5

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	271	NAG	1	0

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.