



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:55 PM GMT

PDB ID : 4AY2  
Title : Capturing 5' tri-phosphorylated RNA duplex by RIG-I  
Authors : Luo, D.; Pyle, A.M.  
Deposited on : 2012-06-17  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)
RNA backbone	2183	1091 (3.20-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	687	<div> <div>2%</div> <div>76%</div> <div>15%</div> <div>9%</div> </div>
2	C	20	<div> <div>65%</div> <div>25%</div> <div>10%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROBABLE ATP-DEPENDENT RNA HELICASE DDX58.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	628	Total	C	N	O	S	0	0	0
			5007	3215	845	918	29			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	306	ASN	GLN	CONFLICT	UNP O95786
A	828	ASP	GLU	CONFLICT	UNP O95786

- Molecule 2 is a RNA chain called 5'-R-PPP(GP\*GP\*CP\*GP\*CP\*GP\*GP\*CP\*UP\*UP\*CP\*GP\*GP\*CP\*CP\*GP\*CP\*GP\*CP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	20	Total	C	N	O	P	0	0	0
			436	189	76	149	22			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	
			27	10	5	10	2	

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg		
			1	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	67	Total	O		
			67	67	0	0
6	C	3	Total	O		
			3	3	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: PROBABLE ATP-DEPENDENT RNA HELICASE DDX58



#### • Molecule 2: 5'-R-PPP(GP\*GP\*CP\*GP\*CP\*GP\*GP\*CP\*UP\*UP\*CP\*GP\*GP\*CP\*CP\*GP\*CP\*GP\*CP\*C)-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.74Å 76.16Å 221.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.90 – 2.80 24.74 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.9 (24.90-2.80) 98.1 (24.74-2.80)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.59 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.218 , 0.286 0.218 , 0.288	Depositor DCC
$R_{free}$ test set	1036 reflections (5.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.9	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 36.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 20260 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5542	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, ZN, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/5111	0.56	1/6902 (0.0%)
2	C	0.38	0/449	0.66	0/698
All	All	0.41	0/5560	0.57	1/7600 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	768	LEU	CA-CB-CG	5.92	128.92	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5007	0	4986	60	0
2	C	436	0	219	6	0
3	A	1	0	0	0	0
4	A	27	0	12	2	0
5	A	1	0	0	0	0
6	A	67	0	0	1	0
6	C	3	0	0	1	0
All	All	5542	0	5217	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:575:ARG:NH2	1:A:591:GLU:OE2	2.20	0.73
1:A:269:GLY:HA2	4:A:1924:ADP:O2A	1.92	0.70
1:A:815:ALA:HB2	1:A:866:ARG:HE	1.56	0.69
1:A:451:GLN:OE1	6:A:2031:HOH:O	2.14	0.65
1:A:657:LYS:H	1:A:694:LEU:HD11	1.61	0.65
1:A:845:ARG:HB3	1:A:863:PHE:HE1	1.62	0.64
1:A:267:GLY:N	4:A:1924:ADP:O2B	2.27	0.62
1:A:461:ARG:HB2	1:A:741:LEU:HD22	1.82	0.61
1:A:888:LYS:HB3	2:C:1:GTP:H5'	1.84	0.59
1:A:478:LEU:O	1:A:482:THR:HG23	2.02	0.59
1:A:656:LEU:HA	1:A:694:LEU:HD11	1.87	0.56
1:A:657:LYS:O	1:A:693:ILE:HA	2.06	0.55
1:A:899:THR:OG1	1:A:900:GLY:N	2.37	0.55
1:A:510:GLU:OE2	1:A:546:ARG:NH1	2.40	0.54
2:C:12:G:H2'	2:C:13:G:O4'	2.08	0.54
1:A:242:LYS:NZ	1:A:243:PRO:O	2.40	0.53
1:A:352:VAL:O	1:A:356:LYS:HD3	2.08	0.53
1:A:471:PHE:HB2	1:A:560:MET:HE3	1.91	0.53
1:A:811:ARG:HG3	1:A:893:VAL:HG23	1.92	0.52
1:A:893:VAL:HG22	1:A:904:LEU:HD21	1.92	0.51
1:A:598:GLU:HG3	1:A:602:ARG:NH2	2.25	0.51
1:A:352:VAL:HG13	1:A:356:LYS:NZ	2.26	0.51
1:A:773:GLU:HG2	1:A:777:ARG:NH1	2.26	0.50
1:A:258:LYS:O	1:A:260:THR:HG22	2.13	0.49
1:A:264:ALA:HB3	1:A:270:LYS:HD2	1.93	0.49
1:A:571:PHE:O	1:A:575:ARG:HG2	2.12	0.48
1:A:579:PHE:CE2	1:A:584:GLN:HG2	2.49	0.48
1:A:913:PHE:O	1:A:915:LYS:N	2.48	0.47
1:A:560:MET:SD	1:A:606:ASN:HB3	2.55	0.47
1:A:560:MET:HE3	1:A:600:VAL:HG12	1.96	0.47
1:A:869:CYS:O	1:A:870:SER:HB2	2.15	0.47
1:A:560:MET:CE	1:A:600:VAL:HG12	2.45	0.46
1:A:657:LYS:N	1:A:694:LEU:HD11	2.29	0.46
1:A:811:ARG:HA	1:A:893:VAL:HG23	1.98	0.46
1:A:447:GLU:N	1:A:447:GLU:OE1	2.49	0.45
1:A:919:ASP:OD1	1:A:920:PRO:HD2	2.16	0.45
1:A:557:HIS:HB2	1:A:639:LEU:HD21	1.98	0.45

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:823:VAL:HG22	1:A:833:VAL:HG22	1.98	0.45
1:A:460:PHE:CE1	1:A:751:GLU:HG2	2.52	0.44
2:C:18:G:H5''	6:C:2003:HOH:O	2.17	0.44
1:A:510:GLU:O	1:A:514:VAL:HG23	2.16	0.44
1:A:899:THR:OG1	1:A:901:VAL:HG23	2.17	0.44
1:A:830:HIS:HE1	2:C:1:GTP:HN22	1.64	0.44
1:A:861:LYS:NZ	1:A:872:ASP:OD1	2.49	0.44
1:A:350:ILE:O	1:A:354:ASN:ND2	2.51	0.44
1:A:766:LEU:HD12	1:A:766:LEU:HA	1.86	0.43
1:A:251:ALA:HB2	1:A:273:VAL:HG13	2.00	0.43
1:A:610:LYS:HE2	1:A:743:SER:HB3	2.01	0.43
1:A:869:CYS:SG	1:A:870:SER:N	2.92	0.42
1:A:578:GLY:HA3	1:A:579:PHE:HA	1.82	0.42
1:A:355:LEU:HD11	1:A:363:LEU:HD21	2.02	0.42
1:A:293:VAL:HG22	1:A:368:LEU:HB3	2.02	0.41
1:A:830:HIS:CE1	2:C:1:GTP:HN22	2.38	0.41
1:A:608:ASN:HA	1:A:609:PRO:HD2	1.87	0.41
1:A:360:ILE:HA	1:A:361:PRO:HD3	1.89	0.41
1:A:807:LYS:HD3	1:A:816:LEU:HD13	2.01	0.41
1:A:454:TYR:HD1	1:A:456:PRO:HD3	1.86	0.41
1:A:560:MET:HB3	1:A:560:MET:HE2	1.73	0.40
1:A:463:VAL:O	1:A:743:SER:HA	2.22	0.40
1:A:694:LEU:H	1:A:694:LEU:HD12	1.87	0.40
1:A:324:ILE:HG22	1:A:350:ILE:HD13	2.04	0.40
2:C:10:U:H5'	2:C:11:C:H5	1.86	0.40
1:A:643:LEU:HD13	1:A:643:LEU:HA	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	618/687 (90%)	573 (93%)	40 (6%)	5 (1%)	24 58

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	880	LYS
1	A	914	GLU
1	A	800	VAL
1	A	867	GLN
1	A	869	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	548/615 (89%)	526 (96%)	22 (4%)	38 73

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	249	GLU
1	A	260	THR
1	A	305	GLN
1	A	328	THR
1	A	359	THR
1	A	393	GLN
1	A	398	SER
1	A	406	ILE
1	A	482	THR
1	A	488	ARG
1	A	500	GLN
1	A	520	CYS
1	A	538	LEU
1	A	559	ARG
1	A	660	ILE
1	A	690	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	694	LEU
1	A	697	THR
1	A	699	VAL
1	A	768	LEU
1	A	881	THR
1	A	911	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	451	GLN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	18/20 (90%)	3 (16%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	7	G
2	C	10	U
2	C	11	C

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GTP	C	1	2	25,34,34	1.04	3 (12%)	34,54,54	1.71	8 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	C	1	2	-	0/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	GTP	O4'-C1'	2.38	1.44	1.41
2	C	1	GTP	C5-C4	2.53	1.46	1.40
2	C	1	GTP	C6-C5	2.90	1.47	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GTP	PA-O3A-PB	-3.67	122.42	132.73
2	C	1	GTP	C2'-C1'-N9	-3.65	108.72	114.29
2	C	1	GTP	C5-C6-N1	-3.16	119.26	123.59
2	C	1	GTP	N3-C2-N1	-3.04	122.81	127.44
2	C	1	GTP	C6-C5-C4	-2.79	117.56	120.90
2	C	1	GTP	C4-C5-N7	-2.38	107.29	109.48
2	C	1	GTP	O3G-PG-O2G	2.37	116.42	107.38
2	C	1	GTP	C6-N1-C2	3.65	121.01	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	GTP	3	0

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ADP	A	1924	5	22,29,29	1.07	1 (4%)	27,45,45	2.14	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	1924	5	-	0/12/32/32	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1924	ADP	C5-C4	3.20	1.47	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1924	ADP	N3-C2-N1	-7.06	123.48	128.89
4	A	1924	ADP	C2'-C1'-N9	-4.26	107.78	114.29
4	A	1924	ADP	C4-C5-N7	-3.68	106.10	109.48
4	A	1924	ADP	PA-O3A-PB	-3.49	120.97	132.67
4	A	1924	ADP	O3A-PA-O5'	-2.79	95.55	102.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1924	ADP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	628/687 (91%)	-0.23	13 (2%) 67 56	21, 51, 91, 117	0
2	C	19/20 (95%)	-0.00	0 100 100	39, 51, 105, 105	0
All	All	647/707 (91%)	-0.22	13 (2%) 68 58	21, 51, 91, 117	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	690	ASP	4.1
1	A	241	PHE	3.6
1	A	691	HIS	3.2
1	A	468	SER	3.1
1	A	868	ASN	2.9
1	A	473	TYR	2.9
1	A	799	PRO	2.7
1	A	454	TYR	2.6
1	A	692	ASN	2.4
1	A	521	MET	2.3
1	A	869	CYS	2.3
1	A	465	SER	2.2
1	A	331	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GTP	C	1	32/32	0.93	0.18	-	50,64,82,84	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ADP	A	1924	27/27	0.93	0.15	-0.32	51,60,68,71	0
3	ZN	A	1923	1/1	0.97	0.03	-2.93	84,84,84,84	0
5	MG	A	1925	1/1	0.97	0.14	-	49,49,49,49	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.