



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 22, 2016 – 08:45 PM EDT

PDB ID : 5B2Q
Title : Crystal structure of Francisella novicida Cas9 RHA in complex with sgRNA and target DNA (TGG PAM)
Authors : Hirano, H.; Nishimasu, H.; Nakane, T.; Ishitani, R.; Nureki, O.
Deposited on : 2016-02-01
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0122
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)
RNA backbone	2183	1045 (2.70-0.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1632	<div> <div>14%</div> <div> <div></div> <div>78%</div> <div>10%</div> <div>11%</div> </div> </div>
2	B	94	<div> <div></div> <div> <div>71%</div> <div>26%</div> <div>3%</div> </div> </div>
3	C	30	<div> <div></div> <div> <div>63%</div> <div>30%</div> <div>7%</div> </div> </div>
4	D	9	<div> <div>22%</div> <div> <div></div> <div>67%</div> <div>33%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	ACT	A	1728	-	-	-	X
9	EDO	A	1722	-	-	-	X
9	EDO	A	1725	-	-	-	X
9	EDO	A	1727	-	-	-	X
9	EDO	B	111	-	-	-	X
9	EDO	B	116	-	-	-	X
9	EDO	B	121	-	-	-	X

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 15694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR-associated endonuclease Cas9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1455	Total	C	N	O	S	0	16	0
			11802	7550	2027	2194	31			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP A0Q5Y3
A	-1	SER	-	expression tag	UNP A0Q5Y3
A	0	HIS	-	expression tag	UNP A0Q5Y3
A	995	ALA	ASN	engineered mutation	UNP A0Q5Y3
A	1369	ARG	GLU	conflict	UNP A0Q5Y3
A	1449	HIS	GLU	conflict	UNP A0Q5Y3
A	1556	ALA	ARG	conflict	UNP A0Q5Y3

- Molecule 2 is a RNA chain called Guide RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	94	Total	C	N	O	P	0	0	0
			1991	886	350	661	94			

- Molecule 3 is a DNA chain called Target DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	30	Total	C	N	O	P	0	0	0
			595	285	105	176	29			

- Molecule 4 is a DNA chain called DNA (5'-D(*TP*GP*GP*TP*AP*TP*CP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	9	Total	C	N	O	P	0	0	0
			185	89	34	54	8			

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	Na	0	0
			2	2		
6	A	2	Total	Na	0	0
			2	2		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Cl	0	0
			2	2		

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	7	Total	Ca	0	0
			7	7		
8	A	10	Total	Ca	0	0
			10	10		

- Molecule 9 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 10 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			4	2	2		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 11 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	588	Total	O	0	0
			588	588		
11	B	334	Total	O	0	0
			334	334		
11	C	64	Total	O	0	0
			64	64		
11	D	7	Total	O	0	0
			7	7		




• Molecule 2: Guide RNA



• Molecule 3: Target DNA



• Molecule 4: DNA (5'-D(*TP*GP*GP*TP*AP*TP*CP*GP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.56Å 158.98Å 96.74Å 90.00° 106.88° 90.00°	Depositor
Resolution (Å)	46.29 – 1.70 46.29 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.29-1.70) 98.9 (46.29-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.70Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.183 , 0.206 0.189 , 0.211	Depositor DCC
R_{free} test set	12762 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.084	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 254951 reflections (0.000%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15694	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL, NA, CA, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/12077	0.55	2/16290 (0.0%)
2	B	0.88	0/2224	1.31	13/3465 (0.4%)
3	C	1.02	0/664	1.23	6/1018 (0.6%)
4	D	0.82	0/207	1.05	1/319 (0.3%)
All	All	0.55	0/15172	0.78	22/21092 (0.1%)

There are no bond length outliers.

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	27	A	O5'-P-OP2	-13.64	93.43	105.70
3	C	11	DC	O5'-P-OP2	-11.75	95.13	105.70
3	C	12	DC	O5'-P-OP2	-9.99	96.71	105.70
2	B	66	U	O5'-P-OP1	-7.66	98.81	105.70
2	B	47	U	C2-N3-C4	-6.86	122.88	127.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11802	0	11583	98	1
2	B	1991	0	997	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	595	0	337	11	0
4	D	185	0	104	1	0
5	A	1	0	0	0	0
6	A	2	0	0	0	0
6	B	2	0	0	0	0
7	A	2	0	0	0	0
8	A	10	0	0	0	0
8	B	7	0	0	0	0
9	A	48	0	72	4	0
9	B	48	0	72	3	0
10	A	8	0	6	0	0
11	A	588	0	0	9	0
11	B	334	0	0	2	0
11	C	64	0	0	1	0
11	D	7	0	0	0	0
All	All	15694	0	13171	116	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 116 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:ASN:O	1:A:291:LEU:N	2.06	0.88
1:A:906:PRO:HA	1:A:916:LEU:HD21	1.59	0.83
1:A:648[A]:ASP:OD2	1:A:1104:TYR:OH	2.03	0.76
1:A:578:LYS:NZ	1:A:582:GLU:OE2	2.20	0.75
1:A:169:LYS:HB3	1:A:187:LEU:HD11	1.68	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:ARG:NH2	1:A:1260:GLU:OE2[2_445]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1444/1632 (88%)	1412 (98%)	30 (2%)	2 (0%)	56 35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	264	PHE
1	A	1128	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1260/1483 (85%)	1218 (97%)	42 (3%)	45 22

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	723	LEU
1	A	856	ARG
1	A	1570	ASP
1	A	724	LEU
1	A	791	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	659	HIS
1	A	720	ASN
1	A	1234	ASN
1	A	642	ASN
1	A	1236	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	92/94 (97%)	5 (5%)	2 (2%)

All (5) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	29	U
2	B	53	C
2	B	67	U
2	B	83	U
2	B	84	U

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	66	U
2	B	67	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 50 ligands modelled in this entry, 24 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	EDO	A	1716	-	3,3,3	0.39	0	2,2,2	0.47	0
9	EDO	A	1717	-	3,3,3	0.43	0	2,2,2	0.51	0
9	EDO	A	1718	-	3,3,3	0.48	0	2,2,2	0.07	0
9	EDO	A	1719	-	3,3,3	0.61	0	2,2,2	0.34	0
9	EDO	A	1720	-	3,3,3	0.55	0	2,2,2	0.25	0
9	EDO	A	1721	-	3,3,3	0.37	0	2,2,2	0.60	0
9	EDO	A	1722	-	3,3,3	0.50	0	2,2,2	0.28	0
9	EDO	A	1723	-	3,3,3	0.49	0	2,2,2	0.12	0
9	EDO	A	1724	-	3,3,3	0.52	0	2,2,2	0.14	0
9	EDO	A	1725	-	3,3,3	0.47	0	2,2,2	0.43	0
9	EDO	A	1726	-	3,3,3	0.48	0	2,2,2	0.21	0
9	EDO	A	1727	-	3,3,3	0.35	0	2,2,2	0.66	0
10	ACT	A	1728	-	0,3,3	0.00	-	0,3,3	0.00	-
10	ACT	A	1729	-	0,3,3	0.00	-	0,3,3	0.00	-
9	EDO	B	110	-	3,3,3	0.43	0	2,2,2	0.44	0
9	EDO	B	111	-	3,3,3	0.52	0	2,2,2	0.46	0
9	EDO	B	112	-	3,3,3	0.62	0	2,2,2	0.04	0
9	EDO	B	113	-	3,3,3	0.64	0	2,2,2	0.11	0
9	EDO	B	114	-	3,3,3	0.48	0	2,2,2	0.39	0
9	EDO	B	115	-	3,3,3	0.51	0	2,2,2	0.16	0
9	EDO	B	116	-	3,3,3	0.51	0	2,2,2	0.15	0
9	EDO	B	117	-	3,3,3	0.43	0	2,2,2	0.60	0
9	EDO	B	118	-	3,3,3	0.43	0	2,2,2	0.62	0
9	EDO	B	119	-	3,3,3	0.48	0	2,2,2	0.04	0
9	EDO	B	120	-	3,3,3	0.37	0	2,2,2	0.37	0
9	EDO	B	121	-	3,3,3	0.48	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	EDO	A	1716	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1717	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1718	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1719	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	EDO	A	1720	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1721	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1722	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1723	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1724	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1725	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1726	-	-	0/1/1/1	0/0/0/0
9	EDO	A	1727	-	-	0/1/1/1	0/0/0/0
10	ACT	A	1728	-	-	0/0/0/0	0/0/0/0
10	ACT	A	1729	-	-	0/0/0/0	0/0/0/0
9	EDO	B	110	-	-	0/1/1/1	0/0/0/0
9	EDO	B	111	-	-	0/1/1/1	0/0/0/0
9	EDO	B	112	-	-	0/1/1/1	0/0/0/0
9	EDO	B	113	-	-	0/1/1/1	0/0/0/0
9	EDO	B	114	-	-	0/1/1/1	0/0/0/0
9	EDO	B	115	-	-	0/1/1/1	0/0/0/0
9	EDO	B	116	-	-	0/1/1/1	0/0/0/0
9	EDO	B	117	-	-	0/1/1/1	0/0/0/0
9	EDO	B	118	-	-	0/1/1/1	0/0/0/0
9	EDO	B	119	-	-	0/1/1/1	0/0/0/0
9	EDO	B	120	-	-	0/1/1/1	0/0/0/0
9	EDO	B	121	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	1721	EDO	1	0
9	A	1725	EDO	2	0
9	A	1727	EDO	1	0
9	B	119	EDO	1	0
9	B	120	EDO	1	0
9	B	121	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1455/1632 (89%)	0.66	232 (15%) 3 3	18, 45, 101, 143	0
2	B	93/94 (98%)	-0.26	0 100 100	19, 29, 58, 79	0
3	C	30/30 (100%)	-0.11	0 100 100	24, 51, 79, 115	0
4	D	9/9 (100%)	0.51	2 (22%) 1 1	36, 52, 110, 111	0
All	All	1587/1765 (89%)	0.59	234 (14%) 3 4	18, 44, 101, 143	0

The worst 5 of 234 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1066	LEU	12.9
1	A	1059	ALA	10.1
1	A	738	CYS	9.5
1	A	1053	THR	9.2
1	A	723	LEU	8.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
9	EDO	A	1727	4/4	0.77	0.36	7.20	81,81,82,83	0
9	EDO	B	116	4/4	0.92	0.11	5.53	39,43,48,50	0
10	ACT	A	1728	4/4	0.95	0.15	4.86	82,84,84,85	0
9	EDO	A	1722	4/4	0.97	0.11	4.77	28,34,39,43	0
9	EDO	B	121	4/4	0.85	0.22	4.53	56,59,61,62	0
9	EDO	B	111	4/4	0.96	0.10	2.46	30,32,32,36	0
9	EDO	A	1725	4/4	0.92	0.12	2.20	42,45,51,57	0
9	EDO	A	1724	4/4	0.87	0.12	1.94	45,47,48,50	0
9	EDO	B	117	4/4	0.95	0.15	1.72	34,46,52,55	0
9	EDO	B	115	4/4	0.89	0.15	1.63	47,51,52,56	0
9	EDO	A	1717	4/4	0.97	0.12	1.10	37,42,46,48	0
9	EDO	B	110	4/4	0.98	0.07	0.90	31,32,35,37	0
9	EDO	A	1720	4/4	0.90	0.13	0.80	33,34,36,38	0
9	EDO	B	113	4/4	0.98	0.11	0.57	29,29,31,33	0
9	EDO	B	112	4/4	0.98	0.14	0.46	23,24,24,26	0
9	EDO	A	1718	4/4	0.98	0.07	0.41	32,37,38,38	0
9	EDO	B	114	4/4	0.94	0.09	0.02	34,40,42,44	0
9	EDO	A	1716	4/4	0.96	0.12	-0.17	39,45,45,47	0
9	EDO	A	1721	4/4	0.92	0.12	-0.25	45,46,47,49	0
5	ZN	A	1701	1/1	1.00	0.11	-0.56	25,25,25,25	0
9	EDO	A	1723	4/4	0.95	0.12	-0.60	45,45,49,52	0
7	CL	A	1705	1/1	0.99	0.07	-0.70	28,28,28,28	0
8	CA	A	1715	1/1	0.97	0.07	-0.75	54,54,54,54	0
7	CL	A	1704	1/1	0.98	0.05	-1.40	43,43,43,43	0
10	ACT	A	1729	4/4	0.95	0.07	-1.54	35,35,36,37	0
9	EDO	A	1719	4/4	0.89	0.12	-1.54	30,36,37,38	0
8	CA	B	109	1/1	0.98	0.07	-1.60	70,70,70,70	0
8	CA	A	1706	1/1	0.98	0.05	-1.61	67,67,67,67	0
6	NA	B	101	1/1	1.00	0.07	-1.90	28,28,28,28	0
8	CA	B	103	1/1	0.99	0.04	-2.22	33,33,33,33	0
6	NA	A	1702	1/1	0.96	0.04	-2.94	47,47,47,47	0
8	CA	A	1709	1/1	0.99	0.05	-3.94	49,49,49,49	0
8	CA	B	106	1/1	0.98	0.06	-5.59	44,44,44,44	0
8	CA	A	1712	1/1	0.94	0.11	-	72,72,72,72	0
8	CA	B	105	1/1	0.99	0.06	-	35,35,35,35	0
9	EDO	B	118	4/4	0.95	0.10	-	45,54,58,58	0
8	CA	A	1711	1/1	0.96	0.04	-	75,75,75,75	0
8	CA	A	1708	1/1	0.99	0.04	-	47,47,47,47	0
8	CA	A	1714	1/1	0.98	0.14	-	69,69,69,69	0
9	EDO	B	119	4/4	0.49	0.28	-	63,65,69,70	0
8	CA	A	1710	1/1	0.97	0.07	-	47,47,47,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	NA	B	102	1/1	0.98	0.15	-	59,59,59,59	0
8	CA	A	1707	1/1	0.99	0.03	-	36,36,36,36	0
9	EDO	B	120	4/4	0.90	0.11	-	50,51,55,57	0
8	CA	A	1713	1/1	0.96	0.11	-	82,82,82,82	0
8	CA	B	104	1/1	0.95	0.04	-	67,67,67,67	0
8	CA	B	107	1/1	0.98	0.08	-	64,64,64,64	0
6	NA	A	1703	1/1	0.78	0.10	-	63,63,63,63	0
8	CA	B	108	1/1	0.95	0.06	-	69,69,69,69	0
9	EDO	A	1726	4/4	0.78	0.17	-	83,85,85,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.