



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:37 AM GMT

PDB ID : 2B65
Title : Crystal structure of the complex of C-lobe of bovine lactoferrin with maltose at 1.5Å resolution
Authors : Singh, N.; Prem kumar, R.; Jabeen, T.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2005-09-30
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	345	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAL	A	3001	X	-	-	-
3	MAN	A	3	X	-	-	-
3	MAN	A	4	X	-	-	-
3	MAN	A	5	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MAN	A	3006	X	-	-	-
5	MAN	A	3008	X	-	-	-
5	MAN	A	3009	X	-	-	-

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 3241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	341	2604	1622	454	507	21	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	565	LYS	ASN	SEE REMARK 999	UNP P24627
A	608	GLU	LYS	SEE REMARK 999	UNP P24627

- Molecule 2 is SUGAR (MALTOSE) (three-letter code: MAL) (formula: C₁₂H₂₂O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	23	12	11	0	0

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 4 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 5 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	6	Total	C	N	O	0	0
			72	40	2	30		

- Molecule 6 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Fe	0	0
			1	1		

- Molecule 7 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	3	Total	Zn	0	0
			3	3		

- Molecule 9 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	O	S	0	0
			5	4	1		
9	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	435	Total	O	0	0
			435	435		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Lactotransferrin



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.48 Å 49.85 Å 65.16 Å 90.00° 107.07° 90.00°	Depositor
Resolution (Å)	20.00 – 1.50	Depositor
% Data completeness (in resolution range)	90.3 (20.00-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.193 , 0.219	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3241	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, BMA, NAG, ZN, FE, MAN, MAL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/2652	0.82	6/3591 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
3	A	3	0
5	A	3	0
All	All	6	2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	684	CYS	C-N-CA	9.54	145.55	121.70
1	A	600	ARG	NE-CZ-NH2	-8.39	116.11	120.30
1	A	685	ALA	N-CA-CB	7.68	120.85	110.10
1	A	600	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	A	682	GLU	O-C-N	-5.24	114.31	122.70
1	A	682	GLU	CA-C-N	5.18	128.59	117.20

All (6) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	3	MAN	C1
3	A	4	MAN	C1
3	A	5	MAN	C1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
5	A	3006	MAN	C1
5	A	3008	MAN	C1
5	A	3009	MAN	C1

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	417	SER	Peptide
1	A	683	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2604	0	2518	48	0
2	A	23	0	22	1	0
3	A	61	0	52	3	0
4	A	28	0	25	0	0
5	A	72	0	61	3	0
6	A	1	0	0	0	0
7	A	4	0	0	0	0
8	A	3	0	0	0	0
9	A	10	0	0	0	0
10	A	435	0	0	14	0
All	All	3241	0	2678	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:658:GLU:HG2	10:A:3410:HOH:O	1.45	1.13
1:A:514:LYS:HD3	10:A:3407:HOH:O	1.60	0.99
5:A:3006:MAN:H3	5:A:3007:BMA:O5	1.75	0.84
1:A:355:GLN:HE21	1:A:355:GLN:HA	1.43	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:635:GLU:O	1:A:636:THR:HG22	1.77	0.82
1:A:638:ASN:HD22	1:A:643:ASP:H	1.32	0.74
5:A:3006:MAN:C3	5:A:3007:BMA:O5	2.36	0.73
1:A:404:LYS:NZ	1:A:682:GLU:HA	2.05	0.71
1:A:671:ASN:HA	1:A:674:LYS:HD2	1.75	0.69
3:A:2:NAG:O3	3:A:3:MAN:H2	1.93	0.68
3:A:2:NAG:H81	10:A:3356:HOH:O	1.95	0.66
1:A:632:PHE:O	1:A:633:LYS:HD2	1.97	0.65
1:A:565:LYS:HD3	1:A:567:GLU:H	1.63	0.64
1:A:514:LYS:HE3	10:A:3370:HOH:O	1.96	0.64
1:A:638:ASN:ND2	1:A:643:ASP:H	1.97	0.62
1:A:416:LYS:HG2	10:A:3194:HOH:O	1.98	0.62
1:A:600:ARG:HD3	10:A:3101:HOH:O	2.00	0.61
1:A:404:LYS:HZ2	1:A:682:GLU:HA	1.67	0.58
1:A:395:ASP:HA	1:A:595:HIS:CD2	2.41	0.55
1:A:635:GLU:O	1:A:636:THR:CG2	2.51	0.55
5:A:3006:MAN:H5	5:A:3007:BMA:O2	2.08	0.54
1:A:566:ARG:HG2	1:A:581:VAL:HG21	1.91	0.53
1:A:626:PRO:HA	1:A:630:CYS:SG	2.49	0.52
1:A:627:ASP:HB3	10:A:3380:HOH:O	2.09	0.52
1:A:423:LEU:HD21	10:A:3424:HOH:O	2.11	0.51
1:A:510:GLN:HG3	10:A:3204:HOH:O	2.11	0.51
1:A:662:GLY:HA2	2:A:3001:MAL:H5'	1.92	0.51
1:A:663:THR:O	1:A:667:THR:HG23	2.11	0.50
1:A:352:PRO:CG	1:A:520:LYS:HD3	2.42	0.50
1:A:553:ASN:HD21	1:A:565:LYS:HE2	1.78	0.49
1:A:355:GLN:NE2	1:A:355:GLN:HA	2.21	0.49
1:A:404:LYS:HZ1	1:A:682:GLU:HA	1.76	0.48
1:A:356:LYS:HG2	10:A:3420:HOH:O	2.12	0.48
1:A:635:GLU:O	1:A:636:THR:CB	2.62	0.47
1:A:420:HIS:CE1	1:A:429:PRO:HG2	2.51	0.46
1:A:632:PHE:C	1:A:633:LYS:HD2	2.36	0.46
1:A:404:LYS:HD2	1:A:683:ALA:H	1.81	0.45
1:A:668:ALA:HB1	3:A:1:NAG:H83	1.99	0.44
1:A:416:LYS:HE3	1:A:620:LYS:HE2	1.99	0.44
1:A:665:TYR:CZ	1:A:669:ILE:HD11	2.52	0.44
1:A:447:THR:HA	1:A:572:LEU:HD22	2.00	0.43
1:A:360:GLN:NE2	10:A:3355:HOH:O	2.46	0.43
1:A:352:PRO:HG2	1:A:520:LYS:HD3	2.01	0.42
1:A:637:LYS:HG2	10:A:3125:HOH:O	2.20	0.42
1:A:455:LYS:HB3	1:A:504:LEU:HD11	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:636:THR:HG21	10:A:3110:HOH:O	2.21	0.41
1:A:414:ASN:HD22	1:A:414:ASN:HA	1.63	0.41
1:A:510:GLN:HB2	1:A:512:LEU:HG	2.02	0.41
1:A:559:ASP:HA	1:A:562:LYS:HD2	2.03	0.41
1:A:363:GLN:HG3	10:A:3415:HOH:O	2.21	0.40
1:A:553:ASN:ND2	1:A:565:LYS:HE2	2.35	0.40
1:A:364:GLN:OE1	1:A:628:LYS:HB3	2.21	0.40
1:A:461:VAL:O	1:A:462:ASP:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	337/345 (98%)	316 (94%)	17 (5%)	4 (1%)	16 2

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	419	LYS
1	A	685	ALA
1	A	682	GLU
1	A	562	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	282/286 (99%)	265 (94%)	17 (6%)	24 2

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	355	GLN
1	A	367	GLN
1	A	414	ASN
1	A	421	SER
1	A	422	SER
1	A	452	LYS
1	A	515	CYS
1	A	520	LYS
1	A	559	ASP
1	A	567	GLU
1	A	602	ASP
1	A	627	ASP
1	A	631	LEU
1	A	633	LYS
1	A	636	THR
1	A	681	LEU
1	A	682	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	355	GLN
1	A	360	GLN
1	A	414	ASN
1	A	621	ASN
1	A	624	ASN
1	A	638	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

13 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1	1,3	14,14,15	0.56	0	15,19,21	0.75	0
3	NAG	A	2	3	14,14,15	0.49	0	15,19,21	0.84	0
3	MAN	A	3	3	11,11,12	0.51	0	14,15,17	1.81	3 (21%)
4	NAG	A	3002	1,4	14,14,15	0.67	1 (7%)	15,19,21	0.79	0
4	NAG	A	3003	4	14,14,15	0.51	0	15,19,21	1.25	1 (6%)
5	NAG	A	3004	1,5	14,14,15	0.56	0	15,19,21	1.19	1 (6%)
5	NAG	A	3005	5	14,14,15	0.58	0	15,19,21	1.02	2 (13%)
5	MAN	A	3006	5	11,11,12	0.66	0	14,15,17	0.96	1 (7%)
5	BMA	A	3007	5	11,11,12	0.67	0	14,15,17	2.06	6 (42%)
5	MAN	A	3008	5	11,11,12	0.97	0	14,15,17	2.08	3 (21%)
5	MAN	A	3009	5	11,11,12	0.44	0	14,15,17	1.13	1 (7%)
3	MAN	A	4	3	11,11,12	0.63	0	14,15,17	2.03	5 (35%)
3	MAN	A	5	3	11,11,12	0.71	0	14,15,17	1.95	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	2	3	-	0/6/23/26	0/1/1/1
3	MAN	A	3	3	1/1/4/5	0/2/19/22	0/1/1/1
4	NAG	A	3002	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	3003	4	-	0/6/23/26	0/1/1/1
5	NAG	A	3004	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	3005	5	-	0/6/23/26	0/1/1/1
5	MAN	A	3006	5	1/1/4/5	0/2/19/22	0/1/1/1
5	BMA	A	3007	5	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MAN	A	3008	5	1/1/4/5	0/2/19/22	0/1/1/1
5	MAN	A	3009	5	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	A	4	3	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	A	5	3	1/1/4/5	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	3002	NAG	O5-C1	-2.03	1.40	1.43

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	3008	MAN	C1-O5-C5	-6.77	103.66	112.25
5	A	3007	BMA	O4-C4-C3	-4.51	100.19	110.34
3	A	5	MAN	O5-C1-C2	-4.08	104.24	110.86
3	A	5	MAN	O4-C4-C3	-3.26	103.00	110.34
5	A	3007	BMA	O5-C5-C6	-2.67	101.57	107.35
5	A	3007	BMA	O5-C1-C2	-2.33	107.07	110.86
3	A	3	MAN	C2-C3-C4	-2.16	107.38	111.04
5	A	3005	NAG	C6-C5-C4	-2.13	107.75	113.02
3	A	4	MAN	C6-C5-C4	-2.01	108.07	113.02
5	A	3005	NAG	C1-O5-C5	2.06	114.86	112.25
5	A	3008	MAN	C2-C3-C4	2.08	114.57	111.04
5	A	3008	MAN	O2-C2-C1	2.14	113.49	109.21
5	A	3006	MAN	O4-C4-C3	2.60	116.19	110.34
5	A	3007	BMA	C2-C3-C4	2.73	115.69	111.04
5	A	3007	BMA	O4-C4-C5	2.75	116.53	109.24
5	A	3007	BMA	C1-O5-C5	2.82	115.83	112.25
3	A	3	MAN	C1-C2-C3	2.84	112.90	109.54
3	A	4	MAN	C2-C3-C4	2.91	115.98	111.04
3	A	5	MAN	C3-C4-C5	2.93	115.30	110.20
5	A	3004	NAG	C1-O5-C5	2.99	116.05	112.25
3	A	4	MAN	C1-O5-C5	3.11	116.20	112.25
3	A	5	MAN	C2-C3-C4	3.25	116.56	111.04
5	A	3009	MAN	C1-C2-C3	3.58	113.78	109.54
3	A	4	MAN	C3-C4-C5	3.62	116.51	110.20
4	A	3003	NAG	C1-O5-C5	3.82	117.10	112.25
3	A	4	MAN	C1-C2-C3	4.15	114.45	109.54
3	A	3	MAN	C1-O5-C5	5.03	118.63	112.25

All (6) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	3	MAN	C1
5	A	3006	MAN	C1
3	A	5	MAN	C1
5	A	3009	MAN	C1
5	A	3008	MAN	C1
3	A	4	MAN	C1

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	NAG	1	0
3	A	2	NAG	2	0
3	A	3	MAN	1	0
5	A	3006	MAN	3	0
5	A	3007	BMA	3	0

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	SO4	A	2001	-	4,4,4	0.35	0	6,6,6	0.33	0
9	SO4	A	2002	8	4,4,4	0.21	0	6,6,6	0.19	0
2	MAL	A	3001	-	24,24,24	0.59	0	35,35,35	1.27	5 (14%)
7	CO3	A	687	6	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	SO4	A	2001	-	-	0/0/0/0	0/0/0/0
9	SO4	A	2002	8	-	0/0/0/0	0/0/0/0
2	MAL	A	3001	-	1/1/10/10	0/8/48/48	0/2/2/2
7	CO3	A	687	6	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3001	MAL	O5'-C1'-C2'	2.08	113.11	109.80
2	A	3001	MAL	C4-C3-C2	2.58	115.61	110.79
2	A	3001	MAL	C2'-C3'-C4'	2.72	115.58	109.60
2	A	3001	MAL	C1'-C2'-C3'	3.36	115.43	110.43
2	A	3001	MAL	C1-C2-C3	3.68	117.23	109.97

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	3001	MAL	C1'

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3001	MAL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.