



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:35 PM GMT

PDB ID : 1BI3
Title : STRUCTURE OF APO-AND HOLO-DIPHThERIA TOXIN REPRESSOR
Authors : Pohl, E.; Hol, W.G.J.
Deposited on : 1998-06-21
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	
1	B	226	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIPHTHERIA TOXIN REPRESSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	0	0
			1586	983	281	315	7			
1	B	137	Total	C	N	O	S	0	0	0
			1069	660	194	208	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	CSS	CYS	MODIFIED	UNP P33120
B	102	CSS	CYS	MODIFIED	UNP P33120

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	98	Total	O	0	0
			98	98		
4	B	71	Total	O	0	0
			71	71		

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	63.40 Å 63.40 Å 213.20 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.40 33.68 – 2.40	Depositor EDS
% Data completeness (in resolution range)	95.8 (8.00-2.40) 96.2 (33.68-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.71 (at 2.39 Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.253 , 0.333 0.270 , 0.337	Depositor DCC
R_{free} test set	919 reflections (4.84%)	DCC
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 129.4	EDS
Estimated twinning fraction	0.048 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 19592 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2836	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.02 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.6180e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, CSS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1594	0.63	1/2166 (0.0%)
1	B	0.40	0/1074	0.65	0/1454
All	All	0.39	0/2668	0.64	1/3620 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	PRO	N-CA-CB	5.52	109.93	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1586	0	1522	59	0
1	B	1069	0	1074	36	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	98	0	0	6	0
4	B	71	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2836	0	2596	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:VAL:HG21	1:B:5:VAL:HG21	1.62	0.81
1:A:89:ILE:HG21	1:B:89:ILE:HG21	1.61	0.80
1:A:4:LEU:HA	4:A:438:HOH:O	1.82	0.79
1:A:187:ILE:HG12	1:A:206:HIS:HB3	1.67	0.77
1:A:195:ILE:HG22	1:A:204:LEU:HD22	1.67	0.75
1:A:196:VAL:HG22	1:A:197:ASP:H	1.53	0.72
1:A:117:VAL:HG21	1:A:140:VAL:HG21	1.71	0.70
1:A:175:GLU:HB3	4:A:423:HOH:O	1.93	0.68
1:B:5:VAL:HA	1:B:106:HIS:HB3	1.81	0.61
1:A:204:LEU:HD11	1:A:213:LEU:HD11	1.81	0.61
1:B:108:MET:CE	1:B:113:GLU:HG2	2.33	0.59
1:A:185:ALA:HB2	4:A:367:HOH:O	2.03	0.59
1:A:73:THR:HG23	1:A:133:PRO:HB2	1.85	0.57
1:B:73:THR:HG23	1:B:133:PRO:HB2	1.87	0.56
1:B:6:ASP:HB2	1:B:9:GLU:HG3	1.88	0.56
1:A:81:LEU:HB3	1:A:108:MET:HE1	1.87	0.56
1:B:103:ARG:NH2	4:B:465:HOH:O	2.37	0.56
1:B:108:MET:HE2	1:B:113:GLU:HG2	1.88	0.55
1:B:10:MET:HG3	4:B:436:HOH:O	2.07	0.55
1:A:134:GLY:HA2	1:A:137:GLU:HG2	1.89	0.54
1:A:67:THR:HG21	4:A:432:HOH:O	2.08	0.54
1:A:55:VAL:HG22	4:A:339:HOH:O	2.07	0.54
1:A:116:LEU:HA	1:A:119:VAL:HG22	1.89	0.54
1:A:84:ARG:HB3	1:A:120:LEU:HD13	1.89	0.53
1:A:38:GLY:O	1:A:41:VAL:HG22	2.09	0.53
1:A:108:MET:CE	1:A:113:GLU:HG2	2.38	0.52
1:A:168:ILE:HD13	1:A:221:ILE:HG23	1.91	0.51
1:B:13:ARG:HD2	4:B:436:HOH:O	2.11	0.51
1:A:5:VAL:HA	1:A:106:HIS:HB3	1.92	0.51
1:B:116:LEU:HA	1:B:119:VAL:HG22	1.92	0.51
1:B:57:ALA:HB2	1:B:63:GLN:HE21	1.75	0.51
1:A:108:MET:HE3	1:A:113:GLU:HG2	1.93	0.50
1:A:87:THR:HG22	1:A:97:VAL:HG21	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:GLN:OE1	1:A:214:LEU:HB2	2.12	0.49
1:A:30:ILE:HD12	1:A:62:LEU:HD12	1.93	0.49
1:B:137:GLU:H	1:B:137:GLU:CD	2.16	0.48
1:A:31:ALA:HB2	1:A:41:VAL:HG11	1.94	0.48
1:B:16:TYR:CD1	1:B:76:MET:HG3	2.50	0.47
1:A:107:VAL:HB	1:B:103:ARG:HB3	1.96	0.47
1:B:37:SER:HB2	1:B:39:PRO:HD2	1.97	0.47
1:B:80:ARG:HB3	1:B:132:ILE:HG12	1.97	0.47
1:A:45:VAL:HG11	1:A:62:LEU:HD21	1.97	0.46
1:A:194:GLU:O	1:A:204:LEU:HA	2.15	0.46
1:A:10:MET:O	1:A:13:ARG:HG3	2.15	0.46
1:A:156:ALA:HB1	1:A:196:VAL:O	2.17	0.45
1:A:187:ILE:HG12	1:A:206:HIS:CB	2.43	0.45
1:B:45:VAL:O	1:B:49:GLU:HG3	2.17	0.45
1:B:134:GLY:O	1:B:137:GLU:HG2	2.17	0.45
1:A:5:VAL:HB	4:A:397:HOH:O	2.17	0.44
1:A:58:SER:C	1:A:60:ARG:H	2.19	0.44
1:A:113:GLU:O	1:A:117:VAL:HG23	2.17	0.44
1:A:56:VAL:HG22	1:A:62:LEU:HD21	1.98	0.44
1:A:166:VAL:HG23	1:A:224:GLU:H	1.82	0.44
1:A:181:GLN:NE2	1:A:211:VAL:HG13	2.33	0.44
1:A:203:THR:HA	1:A:212:GLU:HA	2.00	0.44
1:B:85:LEU:HD22	1:B:116:LEU:HD21	2.00	0.43
1:A:14:THR:OG1	1:A:34:LEU:HD21	2.18	0.43
1:B:123:VAL:HG13	4:B:353:HOH:O	2.19	0.43
1:A:56:VAL:HG22	1:A:62:LEU:CD2	2.49	0.43
1:B:29:ARG:HD2	1:B:29:ARG:HA	1.81	0.43
1:A:107:VAL:HG21	1:B:107:VAL:HG21	2.01	0.42
1:A:27:ARG:HG2	1:A:62:LEU:HD11	2.01	0.42
1:A:59:ASP:N	1:A:59:ASP:OD1	2.52	0.42
1:A:57:ALA:HB2	1:A:63:GLN:NE2	2.34	0.42
1:B:108:MET:HE3	1:B:113:GLU:HG2	2.01	0.42
1:A:57:ALA:HB2	1:A:63:GLN:HE21	1.84	0.41
1:B:139:GLY:HA2	4:B:408:HOH:O	2.20	0.41
1:A:168:ILE:HG22	1:A:168:ILE:O	2.21	0.41
1:A:169:ASN:ND2	1:A:220:THR:O	2.54	0.41
1:A:85:LEU:HD11	1:B:85:LEU:HD11	2.02	0.41
1:A:56:VAL:HG12	1:A:60:ARG:HA	2.02	0.41
1:B:110:ASP:O	1:B:114:ARG:HG3	2.20	0.41
1:B:76:MET:O	1:B:80:ARG:HG3	2.21	0.41
1:B:87:THR:HG23	1:B:88:ASP:OD1	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:55:VAL:HG12	1:B:63:GLN:HB2	2.02	0.41
1:A:187:ILE:CG1	1:A:206:HIS:HB3	2.45	0.41
1:A:31:ALA:HB1	1:A:36:GLN:O	2.21	0.41
1:A:12:LEU:CD1	1:A:71:LEU:HD23	2.51	0.41
1:A:204:LEU:HG	1:A:213:LEU:HG	2.03	0.41
1:A:103:ARG:NH1	1:B:107:VAL:O	2.54	0.41
1:B:87:THR:HG21	1:B:127:PRO:HG3	2.02	0.41
1:B:34:LEU:O	1:B:35:GLU:HB2	2.21	0.41
1:B:29:ARG:HH12	1:B:33:ARG:HH21	1.69	0.41
1:A:86:LEU:HD23	1:A:90:ILE:HD12	2.03	0.40
1:A:153:ILE:HA	1:A:202:ILE:CD1	2.51	0.40
1:A:166:VAL:HG23	1:A:224:GLU:N	2.36	0.40
1:B:81:LEU:HD13	1:B:132:ILE:HD13	2.03	0.40
1:B:92:LEU:HD12	1:B:93:ASP:H	1.85	0.40
1:A:46:ALA:O	1:A:50:ARG:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/226 (91%)	196 (96%)	7 (3%)	2 (1%)	19	28
1	B	134/226 (59%)	129 (96%)	3 (2%)	2 (2%)	13	17
All	All	339/452 (75%)	325 (96%)	10 (3%)	4 (1%)	16	23

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	59	ASP
1	B	59	ASP
1	B	5	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	196	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/197 (83%)	158 (96%)	6 (4%)	41	62
1	B	117/197 (59%)	112 (96%)	5 (4%)	35	55
All	All	281/394 (71%)	270 (96%)	11 (4%)	39	59

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ARG
1	A	55	VAL
1	A	66	PRO
1	A	81	LEU
1	A	84	ARG
1	A	137	GLU
1	B	13	ARG
1	B	29	ARG
1	B	54	VAL
1	B	81	LEU
1	B	84	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	GLN
1	A	169	ASN
1	B	36	GLN
1	B	63	GLN
1	B	130	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSS	A	102	1	4,6,7	0.52	0	3,6,8	1.14	0
1	CSS	B	102	1	4,6,7	0.52	0	3,6,8	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	102	1	-	0/1/5/7	0/0/0/0
1	CSS	B	102	1	-	0/1/5/7	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	292	2	4,4,4	1.08	0	6,6,6	0.08	0
3	SO4	B	1292	2	4,4,4	1.10	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	292	2	-	0/0/0/0	0/0/0/0
3	SO4	B	1292	2	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/226 (93%)	0.94	45 (21%) 1 1	12, 36, 83, 100	0
1	B	136/226 (60%)	0.40	12 (8%) 12 12	13, 27, 66, 82	0
All	All	347/452 (76%)	0.73	57 (16%) 2 2	12, 33, 80, 100	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	LEU	12.1
1	A	211	VAL	8.4
1	A	156	ALA	7.4
1	A	159	MET	7.2
1	B	4	LEU	7.1
1	B	140	VAL	6.4
1	A	140	VAL	5.9
1	A	201	HIS	5.8
1	A	195	ILE	5.5
1	A	158	SER	5.2
1	A	155	ALA	5.0
1	A	167	GLN	5.0
1	A	160	PRO	4.9
1	A	191	SER	4.4
1	A	186	ASP	4.2
1	B	5	VAL	4.1
1	A	192	GLU	4.1
1	A	185	ALA	4.1
1	A	205	SER	4.0
1	A	166	VAL	3.9
1	A	204	LEU	3.9
1	A	202	ILE	3.9
1	A	157	THR	3.7
1	A	193	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	161	ARG	3.7
1	A	214	LEU	3.7
1	A	196	VAL	3.5
1	A	194	GLU	3.5
1	B	39	PRO	3.4
1	A	208	GLY	3.4
1	A	203	THR	3.4
1	B	57	ALA	3.4
1	A	178	GLN	3.3
1	B	43	GLN	3.0
1	A	152	VAL	3.0
1	A	207	ASN	2.9
1	B	35	GLU	2.9
1	A	213	LEU	2.9
1	A	163	VAL	2.8
1	A	190	GLY	2.8
1	A	176	THR	2.8
1	A	210	ASP	2.7
1	B	42	SER	2.6
1	A	206	HIS	2.6
1	A	153	ILE	2.5
1	A	57	ALA	2.5
1	B	59	ASP	2.5
1	A	221	ILE	2.4
1	B	34	LEU	2.3
1	B	136	ASP	2.2
1	A	5	VAL	2.2
1	A	215	ASP	2.2
1	A	182	LEU	2.1
1	B	41	VAL	2.1
1	A	187	ILE	2.1
1	A	212	GLU	2.0
1	A	148	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CSS	A	102	7/8	0.96	0.14	-	14,19,26,29	0
1	CSS	B	102	7/8	0.98	0.12	-	12,17,21,28	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	A	292	5/5	0.98	0.13	-1.69	34,38,43,46	0
3	SO4	B	1292	5/5	0.97	0.13	-1.72	28,35,38,39	0
2	ZN	A	291	1/1	0.87	0.08	-3.34	38,38,38,38	0
2	ZN	B	1291	1/1	0.96	0.08	-	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.