



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:17 PM GMT

PDB ID : 4C0F  
Title : Structure of the NOT-box domain of human CNOT2  
Authors : Boland, A.; Chen, Y.; Raisch, T.; Jonas, S.; Izaurralde, E.; Weichenrieder, O.  
Deposited on : 2013-08-01  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	117	<div> <div>79%</div> <div>18%</div> </div>
1	B	117	<div>4%</div> <div>80%</div> <div>8%</div> <div>12%</div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6868 atoms, of which 3283 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	96	Total	C	H	N	O	S	0	0	0
			1569	529	757	134	145	4			
1	B	103	Total	C	H	N	O	S	0	0	0
			1690	564	823	144	154	5			
1	C	112	Total	C	H	N	O	S	0	0	0
			1845	614	900	160	166	5			
1	D	100	Total	C	H	N	O	S	0	0	0
			1662	559	803	143	153	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	424	GLN	-	EXPRESSION TAG	UNP Q9NZN8
A	425	GLY	-	EXPRESSION TAG	UNP Q9NZN8
A	426	PRO	-	EXPRESSION TAG	UNP Q9NZN8
A	427	HIS	-	EXPRESSION TAG	UNP Q9NZN8
A	428	MET	-	EXPRESSION TAG	UNP Q9NZN8
B	424	GLN	-	EXPRESSION TAG	UNP Q9NZN8
B	425	GLY	-	EXPRESSION TAG	UNP Q9NZN8
B	426	PRO	-	EXPRESSION TAG	UNP Q9NZN8
B	427	HIS	-	EXPRESSION TAG	UNP Q9NZN8
B	428	MET	-	EXPRESSION TAG	UNP Q9NZN8
C	424	GLN	-	EXPRESSION TAG	UNP Q9NZN8
C	425	GLY	-	EXPRESSION TAG	UNP Q9NZN8
C	426	PRO	-	EXPRESSION TAG	UNP Q9NZN8
C	427	HIS	-	EXPRESSION TAG	UNP Q9NZN8
C	428	MET	-	EXPRESSION TAG	UNP Q9NZN8
D	424	GLN	-	EXPRESSION TAG	UNP Q9NZN8
D	425	GLY	-	EXPRESSION TAG	UNP Q9NZN8
D	426	PRO	-	EXPRESSION TAG	UNP Q9NZN8
D	427	HIS	-	EXPRESSION TAG	UNP Q9NZN8
D	428	MET	-	EXPRESSION TAG	UNP Q9NZN8


- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	40	Total 40	O 40	0	0
2	B	18	Total 18	O 18	0	0
2	C	30	Total 30	O 30	0	0
2	D	14	Total 14	O 14	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 2

Chain A: 





- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 2

Chain B: 




- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 2

Chain C: 



- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 2

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.83 Å 64.83 Å 412.29 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.30 – 2.40 49.30 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.3 (49.30-2.40) 99.4 (49.30-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.80 (at 2.39 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.217 , 0.263 0.208 , 0.255	Depositor DCC
$R_{free}$ test set	1066 reflections (5.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	43.8	Xtriage
Anisotropy	0.666	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 41.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 21312 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6868	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/838	0.41	0/1138
1	B	0.23	0/894	0.37	0/1209
1	C	0.24	0/976	0.38	0/1323
1	D	0.24	0/888	0.38	0/1206
All	All	0.24	0/3596	0.39	0/4876

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	812	757	754	2	0
1	B	867	823	819	5	0
1	C	945	900	896	3	1
1	D	859	803	799	3	1
2	A	40	0	0	0	0
2	B	18	0	0	0	0
2	C	30	0	0	0	0
2	D	14	0	0	0	0
All	All	3585	3283	3268	12	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 12 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:500:TYR:CD1	1:B:509:LYS:HG2	2.46	0.50
1:C:439:TYR:CD2	1:C:443:LEU:HD22	2.49	0.47
1:D:491:THR:CG2	1:D:492:ASN:N	2.79	0.46
1:B:494:TYR:CD2	1:B:495:GLU:N	2.85	0.44
1:A:527:LEU:HD21	1:B:428:MET:HE3	2.00	0.44

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:476:ARG:NH1	1:D:507:TRP:O[8_565]	2.08	0.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/117 (80%)	93 (99%)	1 (1%)	0	100	100
1	B	101/117 (86%)	99 (98%)	2 (2%)	0	100	100
1	C	110/117 (94%)	105 (96%)	5 (4%)	0	100	100
1	D	98/117 (84%)	95 (97%)	3 (3%)	0	100	100
All	All	403/468 (86%)	392 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/102 (79%)	81 (100%)	0	100	100
1	B	87/102 (85%)	86 (99%)	1 (1%)	80	92
1	C	96/102 (94%)	96 (100%)	0	100	100
1	D	88/102 (86%)	87 (99%)	1 (1%)	80	92
All	All	352/408 (86%)	350 (99%)	2 (1%)	90	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	526	HIS
1	D	491	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	526	HIS
1	B	515	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	96/117 (82%)	-0.04	0 <span>100</span> <span>100</span>	33, 45, 69, 82	0
1	B	103/117 (88%)	0.20	5 (4%) <span>33</span> <span>34</span>	32, 51, 89, 97	0
1	C	112/117 (95%)	0.15	2 (1%) <span>71</span> <span>71</span>	35, 49, 72, 84	0
1	D	100/117 (85%)	0.24	3 (3%) <span>54</span> <span>53</span>	32, 48, 74, 97	0
All	All	411/468 (87%)	0.14	10 (2%) <span>62</span> <span>61</span>	32, 49, 79, 97	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	484	GLY	9.0
1	D	483	PRO	3.7
1	C	484	GLY	3.6
1	D	485	MET	3.4
1	B	514	PHE	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.