



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2016 – 11:11 AM GMT

PDB ID : 5C1O
Title : Crystal structure of AMP-PNP complexed D-alanine-D-alanine ligase(DDL)
from *Yersinia pestis*
Authors : Tran, H.T.; Kang, L.W.; Hong, M.K.
Deposited on : 2015-06-15
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div> <div>3%</div> <div>84%</div> <div>12%</div> <div>••</div> </div>
1	B	306	<div> <div>3%</div> <div>82%</div> <div>11%</div> <div>• 5%</div> </div>
1	C	306	<div> <div>92%</div> <div>8%</div> <div>•</div> </div>
1	D	306	<div> <div>4%</div> <div>85%</div> <div>11%</div> <div>••</div> </div>

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-alanine–D-alanine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	0	0
			2265	1446	374	434	11			
1	B	290	Total	C	N	O	S	0	0	0
			2203	1403	366	423	11			
1	C	306	Total	C	N	O	S	0	0	0
			2332	1487	384	450	11			
1	D	295	Total	C	N	O	S	0	0	0
			2241	1429	372	429	11			

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Na	0	0
			2	2		
3	A	1	Total	Na	0	0
			1	1		
3	D	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	8	Total	O	0	0
			8	8		
5	B	7	Total	O	0	0
			7	7		
5	C	17	Total	O	0	0
			17	17		
5	D	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: D-alanine–D-alanine ligase




- Molecule 1: D-alanine–D-alanine ligase



- Molecule 1: D-alanine–D-alanine ligase



- Molecule 1: D-alanine–D-alanine ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.03Å 106.80Å 212.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.69 – 2.50 47.69 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.69-2.50) 99.6 (47.69-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	17.07 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.188 , 0.243 0.191 , 0.243	Depositor DCC
R_{free} test set	2560 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	47.7	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 50494 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9210	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/2309	0.84	1/3129 (0.0%)
1	B	0.76	0/2244	0.87	4/3041 (0.1%)
1	C	0.75	0/2379	0.83	1/3226 (0.0%)
1	D	0.67	0/2283	0.79	0/3094
All	All	0.73	0/9215	0.83	6/12490 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	300	ARG	NE-CZ-NH2	-7.55	116.52	120.30
1	C	211	ASP	CB-CG-OD1	5.64	123.37	118.30
1	A	219	ASP	N-CA-C	5.51	125.87	111.00
1	B	306	ASP	CB-CG-OD2	5.16	122.94	118.30
1	B	300	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	B	47	VAL	N-CA-CB	5.08	122.68	111.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2265	0	2260	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2203	0	2206	20	0
1	C	2332	0	2323	10	0
1	D	2241	0	2244	12	0
2	A	31	0	13	0	0
2	B	31	0	13	0	0
2	C	31	0	13	3	0
2	D	31	0	13	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	8	0	0	0	0
5	B	7	0	0	0	0
5	C	17	0	0	0	0
5	D	4	0	0	0	0
All	All	9210	0	9085	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:151:SER:HB3	2:C:401:ANP:O3G	1.87	0.75
1:C:150:SER:O	1:C:152:VAL:N	2.26	0.68
1:A:144:LYS:NZ	1:A:180:GLU:OE1	2.29	0.65
1:A:253:TRP:CH2	1:A:275:PRO:HD2	2.31	0.65
1:B:259:MET:HG3	1:B:269:LEU:HD11	1.77	0.65
1:B:263:ASP:OD1	1:B:265:HIS:CD2	2.52	0.63
1:C:202:ARG:NH1	1:C:204:GLN:OE1	2.33	0.62
1:B:196:GLU:OE2	1:B:300:ARG:NH2	2.33	0.60
1:A:142:ILE:HD11	1:A:182:TRP:HD1	1.69	0.58
1:C:196:GLU:OE1	1:C:290:TYR:OH	2.19	0.57
1:B:255:ARG:NH1	1:B:257:ASP:OD1	2.38	0.55
1:C:253:TRP:CZ3	1:C:275:PRO:HG2	2.42	0.54
1:C:142:ILE:HD11	1:C:182:TRP:HD1	1.75	0.51
1:B:97:LYS:NZ	1:B:148:GLU:OE1	2.43	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:253:TRP:CH2	1:B:275:PRO:HD2	2.46	0.50
1:A:255:ARG:NH1	1:A:257:ASP:OD1	2.45	0.50
1:D:189:THR:OG1	1:D:255:ARG:HD3	2.11	0.50
1:A:223:TYR:CE2	1:A:282:LEU:HD21	2.47	0.49
1:A:58:VAL:O	1:A:83:TYR:HA	2.12	0.49
1:A:118:ASN:OD1	1:A:120:GLN:N	2.45	0.49
1:A:182:TRP:CH2	1:A:184:SER:HA	2.48	0.49
1:D:27:LEU:HD22	1:D:31:LYS:HD3	1.95	0.48
2:C:401:ANP:O5'	2:C:401:ANP:H8	2.13	0.48
1:A:184:SER:OG	1:A:185:GLY:O	2.30	0.47
1:A:218:SER:OG	1:A:219:ASP:N	2.48	0.46
1:D:225:CYS:SG	1:D:289:GLN:HG2	2.55	0.46
1:B:261:ASP:OD1	1:B:263:ASP:OD1	2.32	0.46
1:A:218:SER:O	1:A:220:LYS:N	2.43	0.46
1:D:240:LEU:HG	1:D:268:LEU:HD21	1.98	0.46
1:B:283:VAL:HB	1:B:284:PRO:HD3	1.98	0.45
1:A:117:LEU:HA	1:A:121:GLN:OE1	2.16	0.45
1:C:97:LYS:HE2	2:C:401:ANP:PB	2.57	0.45
1:B:144:LYS:HB3	1:B:154:MET:HG3	1.98	0.45
1:D:132:ALA:HA	1:D:135:ALA:HB2	1.98	0.44
1:B:119:ARG:NH2	1:B:171:PHE:O	2.46	0.44
1:B:46:PRO:O	1:B:47:VAL:HG22	2.17	0.44
1:B:46:PRO:O	1:B:47:VAL:HG13	2.18	0.44
1:A:12:THR:HG22	1:A:43:LYS:HG3	2.00	0.43
1:B:49:GLN:HG2	1:B:52:GLU:OE2	2.17	0.43
1:B:242:LEU:HA	1:B:242:LEU:HD12	1.83	0.43
1:D:143:VAL:O	1:D:154:MET:HA	2.18	0.43
1:D:195:ASP:OD1	1:D:195:ASP:N	2.51	0.42
1:A:105:GLN:HB2	1:A:111:ILE:HD11	2.00	0.42
1:D:242:LEU:HD12	1:D:242:LEU:HA	1.91	0.42
1:D:283:VAL:HB	1:D:284:PRO:HD3	2.01	0.42
1:D:155:SER:HB2	1:D:166:ALA:HB1	2.01	0.42
1:C:272:ASN:C	1:C:272:ASN:OD1	2.58	0.42
1:C:99:ARG:HD2	1:D:89:MET:HE1	2.01	0.42
1:B:53:GLN:O	1:B:53:GLN:HG3	2.20	0.42
1:C:150:SER:C	1:C:152:VAL:H	2.23	0.41
1:D:253:TRP:CH2	1:D:275:PRO:HD2	2.56	0.41
1:B:118:ASN:OD1	1:B:120:GLN:HG3	2.20	0.41
1:B:16:ARG:NH2	1:B:43:LYS:HE3	2.36	0.41
1:B:198:LEU:HB3	1:B:199:PRO:CD	2.50	0.41
1:A:134:VAL:O	1:A:135:ALA:C	2.59	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:255:ARG:HG2	1:B:255:ARG:HH11	1.86	0.41
1:A:253:TRP:CZ3	1:A:275:PRO:HD2	2.56	0.40
1:B:224:PHE:CD1	1:B:224:PHE:N	2.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/306 (95%)	276 (95%)	12 (4%)	3 (1%)	19	34
1	B	284/306 (93%)	277 (98%)	5 (2%)	2 (1%)	26	46
1	C	304/306 (99%)	291 (96%)	11 (4%)	2 (1%)	26	46
1	D	289/306 (94%)	282 (98%)	6 (2%)	1 (0%)	46	68
All	All	1168/1224 (95%)	1126 (96%)	34 (3%)	8 (1%)	26	46

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	GLN
1	A	121	GLN
1	A	219	ASP
1	B	47	VAL
1	C	151	SER
1	C	16	ARG
1	B	46	PRO
1	D	67	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/251 (97%)	229 (94%)	15 (6%)	23	42
1	B	238/251 (95%)	231 (97%)	7 (3%)	50	77
1	C	251/251 (100%)	239 (95%)	12 (5%)	31	55
1	D	242/251 (96%)	227 (94%)	15 (6%)	23	41
All	All	975/1004 (97%)	926 (95%)	49 (5%)	30	53

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	27	LEU
1	A	124	THR
1	A	155	SER
1	A	169	GLU
1	A	204	GLN
1	A	208	VAL
1	A	220	LYS
1	A	225	CYS
1	A	233	SER
1	A	259	MET
1	A	270	GLU
1	A	300	ARG
1	A	303	MET
1	A	306	ASP
1	B	27	LEU
1	B	120	GLN
1	B	155	SER
1	B	169	GLU
1	B	221	THR
1	B	222	GLN
1	B	262	ARG
1	C	105	GLN
1	C	118	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	133	CYS
1	C	150	SER
1	C	199	PRO
1	C	204	GLN
1	C	231	ASP
1	C	235	GLN
1	C	257	ASP
1	C	259	MET
1	C	300	ARG
1	C	306	ASP
1	D	4	LYS
1	D	24	GLN
1	D	27	LEU
1	D	120	GLN
1	D	126	SER
1	D	133	CYS
1	D	151	SER
1	D	152	VAL
1	D	187	GLU
1	D	195	ASP
1	D	200	SER
1	D	219	ASP
1	D	262	ARG
1	D	263	ASP
1	D	306	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	105	GLN
1	B	49	GLN
1	B	243	GLN
1	B	265	HIS
1	C	24	GLN
1	C	105	GLN
1	C	118	ASN
1	C	147	HIS
1	C	173	HIS
1	C	235	GLN
1	D	53	GLN
1	D	105	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 9 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ANP	A	401	4	29,33,33	2.24	7 (24%)	26,52,52	2.34	8 (30%)
2	ANP	B	401	4	29,33,33	2.13	7 (24%)	26,52,52	2.34	7 (26%)
2	ANP	C	401	4	29,33,33	2.20	6 (20%)	26,52,52	2.61	9 (34%)
2	ANP	D	401	4	29,33,33	1.94	7 (24%)	26,52,52	2.07	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	401	4	-	1/13/38/38	0/3/3/3
2	ANP	B	401	4	-	0/13/38/38	0/3/3/3
2	ANP	C	401	4	-	0/13/38/38	0/3/3/3
2	ANP	D	401	4	-	0/13/38/38	0/3/3/3

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	ANP	C2'-C1'	-2.99	1.48	1.53
2	A	401	ANP	PB-O2B	-2.32	1.50	1.56
2	B	401	ANP	O4'-C1'	2.05	1.44	1.41
2	C	401	ANP	PB-N3B	2.52	1.70	1.63
2	A	401	ANP	PB-O3A	2.53	1.62	1.59
2	D	401	ANP	O4'-C1'	2.61	1.45	1.41
2	C	401	ANP	C5-C4	3.09	1.47	1.40
2	C	401	ANP	PG-N3B	3.12	1.71	1.63
2	A	401	ANP	C5-C4	3.52	1.48	1.40
2	D	401	ANP	PG-O1G	3.53	1.49	1.46
2	D	401	ANP	PB-O3A	3.53	1.63	1.59
2	B	401	ANP	C5-C4	3.60	1.48	1.40
2	D	401	ANP	PG-N3B	3.64	1.73	1.63
2	D	401	ANP	C5-C4	3.65	1.48	1.40
2	D	401	ANP	PB-N3B	3.78	1.73	1.63
2	A	401	ANP	PB-N3B	3.82	1.73	1.63
2	B	401	ANP	PG-N3B	3.90	1.74	1.63
2	A	401	ANP	PG-N3B	3.94	1.74	1.63
2	B	401	ANP	PB-N3B	4.02	1.74	1.63
2	B	401	ANP	PB-O3A	4.03	1.64	1.59
2	D	401	ANP	PB-O1B	4.05	1.50	1.46
2	A	401	ANP	PG-O1G	4.96	1.51	1.46
2	C	401	ANP	PB-O1B	4.98	1.51	1.46
2	B	401	ANP	PB-O1B	5.10	1.51	1.46
2	B	401	ANP	PG-O1G	5.22	1.51	1.46
2	A	401	ANP	PB-O1B	6.66	1.53	1.46
2	C	401	ANP	PG-O1G	7.66	1.54	1.46

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	ANP	N3-C2-N1	-7.83	122.72	128.87
2	D	401	ANP	N3-C2-N1	-7.65	122.86	128.87
2	C	401	ANP	N3-C2-N1	-6.55	123.72	128.87
2	A	401	ANP	N3-C2-N1	-6.53	123.74	128.87
2	C	401	ANP	C1'-N9-C4	-4.38	121.92	126.81
2	C	401	ANP	PA-O3A-PB	-4.25	117.30	132.71
2	A	401	ANP	O2G-PG-O1G	-3.38	104.67	113.58
2	B	401	ANP	PA-O3A-PB	-3.07	121.56	132.71
2	D	401	ANP	PA-O3A-PB	-2.91	122.14	132.71
2	A	401	ANP	PA-O3A-PB	-2.89	122.23	132.71
2	B	401	ANP	C2'-C1'-N9	-2.63	106.43	113.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	ANP	C1'-N9-C4	-2.56	123.94	126.81
2	C	401	ANP	O2'-C2'-C1'	-2.05	105.21	111.61
2	B	401	ANP	O3A-PB-N3B	2.06	111.75	106.07
2	C	401	ANP	O2A-PA-O3A	2.09	114.23	105.27
2	D	401	ANP	O2A-PA-O1A	2.10	123.48	112.56
2	B	401	ANP	N6-C6-N1	2.15	122.12	118.52
2	C	401	ANP	O3'-C3'-C4'	2.23	117.67	111.01
2	C	401	ANP	C2-N1-C6	2.31	122.89	118.77
2	A	401	ANP	O3G-PG-O1G	2.35	119.77	113.58
2	A	401	ANP	C2-N1-C6	2.46	123.16	118.77
2	B	401	ANP	C2-N1-C6	3.16	124.40	118.77
2	D	401	ANP	O3G-PG-O2G	3.25	117.10	107.67
2	C	401	ANP	O3G-PG-O1G	3.52	122.86	113.58
2	D	401	ANP	O2B-PB-O1B	3.63	117.16	110.02
2	A	401	ANP	O3G-PG-O2G	4.43	120.54	107.67
2	A	401	ANP	O2B-PB-O1B	4.95	119.77	110.02
2	B	401	ANP	O2B-PB-O1B	5.52	120.89	110.02
2	C	401	ANP	O2B-PB-O1B	6.90	123.61	110.02

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	ANP	O1G-PG-N3B-PB

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	401	ANP	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/306 (97%)	0.03	8 (2%) 58 62	27, 51, 112, 127	0
1	B	290/306 (94%)	-0.09	10 (3%) 49 54	28, 46, 85, 104	0
1	C	306/306 (100%)	-0.20	0 100 100	29, 44, 73, 99	0
1	D	295/306 (96%)	0.10	11 (3%) 45 50	33, 57, 100, 117	0
All	All	1188/1224 (97%)	-0.04	29 (2%) 62 66	27, 50, 95, 127	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	TYR	5.0
1	D	1	MET	4.1
1	B	2	ALA	3.7
1	D	208	VAL	3.6
1	D	122	PHE	3.3
1	B	223	TYR	3.1
1	D	130	LEU	3.0
1	B	279	SER	2.9
1	D	170	ALA	2.9
1	A	170	ALA	2.9
1	B	125	LEU	2.8
1	A	219	ASP	2.7
1	A	172	GLN	2.7
1	A	133	CYS	2.7
1	A	128	GLU	2.7
1	D	209	PHE	2.7
1	D	128	GLU	2.6
1	D	222	GLN	2.5
1	D	166	ALA	2.5
1	B	206	PRO	2.4
1	A	225	CYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	208	VAL	2.4
1	B	224	PHE	2.3
1	B	1	MET	2.3
1	B	280	HIS	2.3
1	D	207	GLY	2.2
1	B	173	HIS	2.1
1	D	226	PRO	2.0
1	B	225	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ANP	C	401	31/31	0.97	0.14	-0.34	39,46,72,78	0
2	ANP	A	401	31/31	0.96	0.11	-1.12	48,58,109,115	0
2	ANP	D	401	31/31	0.95	0.11	-1.33	50,64,80,86	0
2	ANP	B	401	31/31	0.93	0.12	-1.58	43,58,97,100	0
3	NA	A	402	1/1	0.94	0.11	-2.19	35,35,35,35	0
3	NA	C	402	1/1	0.97	0.10	-2.67	38,38,38,38	0
3	NA	D	402	1/1	0.90	0.09	-2.88	47,47,47,47	0
3	NA	B	402	1/1	0.96	0.06	-5.41	36,36,36,36	0
3	NA	B	404	1/1	0.98	0.36	-	39,39,39,39	0
4	MG	A	403	1/1	0.93	0.08	-	77,77,77,77	0
4	MG	B	403	1/1	0.79	0.14	-	88,88,88,88	0
4	MG	C	403	1/1	0.77	0.15	-	63,63,63,63	0
4	MG	D	403	1/1	0.92	0.06	-	76,76,76,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.