



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 8, 2016 – 03:39 PM EDT

PDB ID : 5C3Y
Title : Structure of human ribokinase crystallized with AMPPNP
Authors : Park, J.; Lee, T.-W.; Chakrabarti, J.; Singh, B.; Gupta, R.S.; Junop, M.S.
Deposited on : 2015-06-17
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div><div>88%</div><div><div></div><div></div><div></div><div></div></div><div>7%</div></div>
1	B	330	<div><div>86%</div><div><div></div><div></div><div></div><div></div></div><div>5%</div></div>
1	C	330	<div><div>88%</div><div><div></div><div></div><div></div><div></div></div><div>5%</div></div>
1	D	330	<div><div>88%</div><div><div></div><div></div><div></div><div></div></div><div>6%</div></div>
1	E	330	<div><div>87%</div><div><div></div><div></div><div></div><div></div></div><div>6%</div></div>
1	F	330	<div><div>87%</div><div><div></div><div></div><div></div><div></div></div><div>6%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	330	 86%8%5%
1	H	330	 87%7%6%
1	I	330	 2%89%5%5%
1	J	330	 89%6%5%
1	K	330	 89%5%5%
1	L	330	 89%5%6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NA	B	403	-	-	-	X
3	NA	E	403	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 28644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	0	0
			2242	1414	371	443	14			
1	B	314	Total	C	N	O	S	0	1	0
			2313	1459	385	454	15			
1	C	313	Total	C	N	O	S	0	1	0
			2307	1454	383	455	15			
1	D	311	Total	C	N	O	S	0	1	0
			2281	1443	374	449	15			
1	E	311	Total	C	N	O	S	0	1	0
			2287	1447	372	453	15			
1	F	312	Total	C	N	O	S	0	0	0
			2292	1446	381	451	14			
1	G	314	Total	C	N	O	S	0	0	0
			2312	1460	383	455	14			
1	H	311	Total	C	N	O	S	0	0	0
			2275	1439	374	448	14			
1	I	313	Total	C	N	O	S	0	0	0
			2303	1453	384	452	14			
1	J	314	Total	C	N	O	S	0	0	0
			2289	1446	376	453	14			
1	K	312	Total	C	N	O	S	0	4	0
			2316	1463	380	457	16			
1	L	311	Total	C	N	O	S	0	1	0
			2269	1434	372	449	14			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	323	LEU	-	expression tag	UNP Q9H477
A	324	GLU	-	expression tag	UNP Q9H477
A	325	HIS	-	expression tag	UNP Q9H477
A	326	HIS	-	expression tag	UNP Q9H477
A	327	HIS	-	expression tag	UNP Q9H477

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	328	HIS	-	expression tag	UNP Q9H477
A	329	HIS	-	expression tag	UNP Q9H477
A	330	HIS	-	expression tag	UNP Q9H477
B	323	LEU	-	expression tag	UNP Q9H477
B	324	GLU	-	expression tag	UNP Q9H477
B	325	HIS	-	expression tag	UNP Q9H477
B	326	HIS	-	expression tag	UNP Q9H477
B	327	HIS	-	expression tag	UNP Q9H477
B	328	HIS	-	expression tag	UNP Q9H477
B	329	HIS	-	expression tag	UNP Q9H477
B	330	HIS	-	expression tag	UNP Q9H477
C	323	LEU	-	expression tag	UNP Q9H477
C	324	GLU	-	expression tag	UNP Q9H477
C	325	HIS	-	expression tag	UNP Q9H477
C	326	HIS	-	expression tag	UNP Q9H477
C	327	HIS	-	expression tag	UNP Q9H477
C	328	HIS	-	expression tag	UNP Q9H477
C	329	HIS	-	expression tag	UNP Q9H477
C	330	HIS	-	expression tag	UNP Q9H477
D	323	LEU	-	expression tag	UNP Q9H477
D	324	GLU	-	expression tag	UNP Q9H477
D	325	HIS	-	expression tag	UNP Q9H477
D	326	HIS	-	expression tag	UNP Q9H477
D	327	HIS	-	expression tag	UNP Q9H477
D	328	HIS	-	expression tag	UNP Q9H477
D	329	HIS	-	expression tag	UNP Q9H477
D	330	HIS	-	expression tag	UNP Q9H477
E	323	LEU	-	expression tag	UNP Q9H477
E	324	GLU	-	expression tag	UNP Q9H477
E	325	HIS	-	expression tag	UNP Q9H477
E	326	HIS	-	expression tag	UNP Q9H477
E	327	HIS	-	expression tag	UNP Q9H477
E	328	HIS	-	expression tag	UNP Q9H477
E	329	HIS	-	expression tag	UNP Q9H477
E	330	HIS	-	expression tag	UNP Q9H477
F	323	LEU	-	expression tag	UNP Q9H477
F	324	GLU	-	expression tag	UNP Q9H477
F	325	HIS	-	expression tag	UNP Q9H477
F	326	HIS	-	expression tag	UNP Q9H477
F	327	HIS	-	expression tag	UNP Q9H477
F	328	HIS	-	expression tag	UNP Q9H477
F	329	HIS	-	expression tag	UNP Q9H477

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	330	HIS	-	expression tag	UNP Q9H477
G	323	LEU	-	expression tag	UNP Q9H477
G	324	GLU	-	expression tag	UNP Q9H477
G	325	HIS	-	expression tag	UNP Q9H477
G	326	HIS	-	expression tag	UNP Q9H477
G	327	HIS	-	expression tag	UNP Q9H477
G	328	HIS	-	expression tag	UNP Q9H477
G	329	HIS	-	expression tag	UNP Q9H477
G	330	HIS	-	expression tag	UNP Q9H477
H	323	LEU	-	expression tag	UNP Q9H477
H	324	GLU	-	expression tag	UNP Q9H477
H	325	HIS	-	expression tag	UNP Q9H477
H	326	HIS	-	expression tag	UNP Q9H477
H	327	HIS	-	expression tag	UNP Q9H477
H	328	HIS	-	expression tag	UNP Q9H477
H	329	HIS	-	expression tag	UNP Q9H477
H	330	HIS	-	expression tag	UNP Q9H477
I	323	LEU	-	expression tag	UNP Q9H477
I	324	GLU	-	expression tag	UNP Q9H477
I	325	HIS	-	expression tag	UNP Q9H477
I	326	HIS	-	expression tag	UNP Q9H477
I	327	HIS	-	expression tag	UNP Q9H477
I	328	HIS	-	expression tag	UNP Q9H477
I	329	HIS	-	expression tag	UNP Q9H477
I	330	HIS	-	expression tag	UNP Q9H477
J	323	LEU	-	expression tag	UNP Q9H477
J	324	GLU	-	expression tag	UNP Q9H477
J	325	HIS	-	expression tag	UNP Q9H477
J	326	HIS	-	expression tag	UNP Q9H477
J	327	HIS	-	expression tag	UNP Q9H477
J	328	HIS	-	expression tag	UNP Q9H477
J	329	HIS	-	expression tag	UNP Q9H477
J	330	HIS	-	expression tag	UNP Q9H477
K	323	LEU	-	expression tag	UNP Q9H477
K	324	GLU	-	expression tag	UNP Q9H477
K	325	HIS	-	expression tag	UNP Q9H477
K	326	HIS	-	expression tag	UNP Q9H477
K	327	HIS	-	expression tag	UNP Q9H477
K	328	HIS	-	expression tag	UNP Q9H477
K	329	HIS	-	expression tag	UNP Q9H477
K	330	HIS	-	expression tag	UNP Q9H477
L	323	LEU	-	expression tag	UNP Q9H477

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	324	GLU	-	expression tag	UNP Q9H477
L	325	HIS	-	expression tag	UNP Q9H477
L	326	HIS	-	expression tag	UNP Q9H477
L	327	HIS	-	expression tag	UNP Q9H477
L	328	HIS	-	expression tag	UNP Q9H477
L	329	HIS	-	expression tag	UNP Q9H477
L	330	HIS	-	expression tag	UNP Q9H477

- # AN2

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 27	C 10	N 6	O 9	P 2	0	0
2	B	1	Total 27	C 10	N 6	O 9	P 2	0	0
2	C	1	Total 27	C 10	N 6	O 9	P 2	0	0
2	D	1	Total 27	C 10	N 6	O 9	P 2	0	0
2	E	1	Total 27	C 10	N 6	O 9	P 2	0	0
2	F	1	Total 27	C 10	N 6	O 9	P 2	0	0
2	G	1	Total 27	C 10	N 6	O 9	P 2	0	0
2	H	1	Total 27	C 10	N 6	O 9	P 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	I	1	Total	C	N	O	P	0	0
			27	10	6	9	2		
2	J	1	Total	C	N	O	P	0	0
			27	10	6	9	2		
2	K	1	Total	C	N	O	P	0	0
			27	10	6	9	2		
2	L	1	Total	C	N	O	P	0	0
			27	10	6	9	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	2	Total	Na	0	0
			2	2		
3	J	2	Total	Na	0	0
			2	2		
3	D	2	Total	Na	0	0
			2	2		
3	K	2	Total	Na	0	0
			2	2		
3	E	2	Total	Na	0	0
			2	2		
3	H	2	Total	Na	0	0
			2	2		
3	B	2	Total	Na	0	0
			2	2		
3	I	2	Total	Na	0	0
			2	2		
3	C	2	Total	Na	0	0
			2	2		
3	A	2	Total	Na	0	0
			2	2		
3	L	2	Total	Na	0	0
			2	2		
3	F	2	Total	Na	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	56	Total	O	0	0
			56	56		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	88	Total 88	O 88	0	0
4	C	96	Total 96	O 96	0	0
4	D	71	Total 71	O 71	0	0
4	E	90	Total 90	O 90	0	0
4	F	46	Total 46	O 46	0	0
4	G	83	Total 83	O 83	0	0
4	H	79	Total 79	O 79	0	0
4	I	54	Total 54	O 54	0	0
4	J	40	Total 40	O 40	0	0
4	K	58	Total 58	O 58	0	0
4	L	49	Total 49	O 49	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribokinase

Chain A:  88% 7%





- Molecule 1: Ribokinase

Chain B:  86% 8% 5%





- Molecule 1: Ribokinase

Chain C:  88% 6% 5%





- Molecule 1: Ribokinase

Chain D:  88% 5% 6%





- Molecule 1: Ribokinase



Chain E:  87% 6% 6%




- Molecule 1: Ribokinase





- Molecule 1: Ribokinase




- Molecule 1: Ribokinase



- Molecule 1: Ribokinase




- Molecule 1: Ribokinase






● Molecule 1: Ribokinase



● Molecule 1: Ribokinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.31Å 168.14Å 161.50Å 90.00° 90.82° 90.00°	Depositor
Resolution (Å)	51.26 – 2.60 51.26 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (51.26-2.60) 99.7 (51.26-2.60)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.205 , 0.232 0.209 , 0.234	Depositor DCC
R_{free} test set	6493 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	32.6	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 14.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for -h,-l,-k 0.020 for -h,l,k 0.150 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	28644	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, AN2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/2274	1.00	11/3095 (0.4%)
1	B	0.94	3/2348 (0.1%)	1.00	14/3197 (0.4%)
1	C	0.99	5/2341 (0.2%)	0.97	10/3187 (0.3%)
1	D	0.97	5/2314 (0.2%)	1.09	10/3150 (0.3%)
1	E	0.99	4/2320 (0.2%)	0.97	6/3158 (0.2%)
1	F	0.93	2/2326 (0.1%)	0.95	9/3167 (0.3%)
1	G	1.02	4/2347 (0.2%)	1.00	10/3194 (0.3%)
1	H	0.96	6/2308 (0.3%)	0.96	6/3143 (0.2%)
1	I	0.86	3/2339 (0.1%)	0.97	7/3185 (0.2%)
1	J	0.89	2/2323 (0.1%)	0.99	9/3166 (0.3%)
1	K	0.86	1/2351 (0.0%)	0.98	5/3202 (0.2%)
1	L	0.89	1/2302 (0.0%)	0.90	5/3138 (0.2%)
All	All	0.93	36/27893 (0.1%)	0.98	102/37982 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	86	GLU	CD-OE2	11.10	1.37	1.25
1	G	200	GLU	CD-OE2	7.93	1.34	1.25
1	F	200	GLU	CG-CD	7.89	1.63	1.51
1	I	134	GLU	CD-OE2	7.58	1.33	1.25
1	D	134	GLU	CD-OE1	7.44	1.33	1.25

The worst 5 of 102 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	169	ARG	NE-CZ-NH2	20.71	130.66	120.30
1	D	169	ARG	NE-CZ-NH1	-20.22	110.19	120.30
1	K	145	ARG	NE-CZ-NH1	-17.53	111.54	120.30
1	A	289	ASP	CB-CG-OD1	15.00	131.80	118.30
1	D	168	ARG	NE-CZ-NH1	12.49	126.55	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	325	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2242	0	2245	6	0
1	B	2313	0	2295	12	0
1	C	2307	0	2295	6	0
1	D	2281	0	2274	4	0
1	E	2287	0	2275	8	0
1	F	2292	0	2285	8	0
1	G	2312	0	2304	9	0
1	H	2275	0	2266	9	0
1	I	2303	0	2293	10	0
1	J	2289	0	2262	4	0
1	K	2316	0	2292	6	0
1	L	2269	0	2246	4	0
2	A	27	0	14	0	0
2	B	27	0	14	0	0
2	C	27	0	14	0	0
2	D	27	0	14	1	0
2	E	27	0	14	0	0
2	F	27	0	14	0	0
2	G	27	0	14	0	0
2	H	27	0	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	I	27	0	14	0	0
2	J	27	0	14	0	0
2	K	27	0	14	0	0
2	L	27	0	14	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
3	G	2	0	0	0	0
3	H	2	0	0	0	0
3	I	2	0	0	0	0
3	J	2	0	0	0	0
3	K	2	0	0	0	0
3	L	2	0	0	0	0
4	A	56	0	0	2	0
4	B	88	0	0	1	0
4	C	96	0	0	1	0
4	D	71	0	0	2	0
4	E	90	0	0	0	0
4	F	46	0	0	0	0
4	G	83	0	0	1	0
4	H	79	0	0	1	0
4	I	54	0	0	0	0
4	J	40	0	0	0	0
4	K	58	0	0	0	0
4	L	49	0	0	2	0
All	All	28644	0	27500	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:145:ARG:NH1	1:E:129:LEU:HD13	1.96	0.80
2:L:401:AN2:O1B	4:L:501:HOH:O	2.07	0.71
1:C:151[B]:CYS:SG	1:C:176:PHE:HD1	2.13	0.71
1:H:188:GLN:NE2	1:I:284:ASN:HD21	1.88	0.69
1:B:115:ASN:OD1	1:B:115:ASN:N	2.22	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/330 (92%)	296 (97%)	8 (3%)	0	100	100
1	B	313/330 (95%)	305 (97%)	8 (3%)	0	100	100
1	C	312/330 (94%)	306 (98%)	6 (2%)	0	100	100
1	D	310/330 (94%)	303 (98%)	7 (2%)	0	100	100
1	E	310/330 (94%)	304 (98%)	6 (2%)	0	100	100
1	F	310/330 (94%)	302 (97%)	8 (3%)	0	100	100
1	G	312/330 (94%)	304 (97%)	8 (3%)	0	100	100
1	H	309/330 (94%)	301 (97%)	8 (3%)	0	100	100
1	I	311/330 (94%)	303 (97%)	8 (3%)	0	100	100
1	J	312/330 (94%)	305 (98%)	7 (2%)	0	100	100
1	K	314/330 (95%)	307 (98%)	7 (2%)	0	100	100
1	L	310/330 (94%)	302 (97%)	8 (3%)	0	100	100
All	All	3727/3960 (94%)	3638 (98%)	89 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/266 (90%)	235 (98%)	5 (2%)	61	85
1	B	246/266 (92%)	238 (97%)	8 (3%)	45	73
1	C	247/266 (93%)	240 (97%)	7 (3%)	51	78
1	D	242/266 (91%)	234 (97%)	8 (3%)	45	73
1	E	243/266 (91%)	235 (97%)	8 (3%)	45	73
1	F	245/266 (92%)	237 (97%)	8 (3%)	45	73
1	G	247/266 (93%)	237 (96%)	10 (4%)	38	67
1	H	241/266 (91%)	233 (97%)	8 (3%)	45	73
1	I	247/266 (93%)	239 (97%)	8 (3%)	46	74
1	J	242/266 (91%)	236 (98%)	6 (2%)	55	81
1	K	247/266 (93%)	239 (97%)	8 (3%)	46	74
1	L	240/266 (90%)	233 (97%)	7 (3%)	50	77
All	All	2927/3192 (92%)	2836 (97%)	91 (3%)	47	76

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	148	VAL
1	G	168	ARG
1	L	15	VAL
1	F	168	ARG
1	G	68	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	325	HIS
1	H	188	GLN
1	H	230	GLN
1	L	45	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 36 ligands modelled in this entry, 24 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	AN2	A	401	-	24,29,29	3.19	6 (25%)	22,45,45	2.26	5 (22%)
2	AN2	B	401	-	24,29,29	2.06	5 (20%)	22,45,45	1.88	3 (13%)
2	AN2	C	401	-	24,29,29	3.27	3 (12%)	22,45,45	2.49	6 (27%)
2	AN2	D	401	-	24,29,29	2.94	5 (20%)	22,45,45	1.82	3 (13%)
2	AN2	E	401	-	24,29,29	2.88	5 (20%)	22,45,45	1.98	5 (22%)
2	AN2	F	401	-	24,29,29	1.77	5 (20%)	22,45,45	2.02	4 (18%)
2	AN2	G	401	-	24,29,29	1.60	4 (16%)	22,45,45	2.10	2 (9%)
2	AN2	H	401	-	24,29,29	3.54	6 (25%)	22,45,45	2.26	4 (18%)
2	AN2	I	401	-	24,29,29	1.59	3 (12%)	22,45,45	2.30	6 (27%)
2	AN2	J	401	-	24,29,29	1.97	4 (16%)	22,45,45	2.51	5 (22%)
2	AN2	K	401	-	24,29,29	1.52	3 (12%)	22,45,45	2.25	4 (18%)
2	AN2	L	401	-	24,29,29	1.56	4 (16%)	22,45,45	2.56	10 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AN2	A	401	-	-	0/9/32/32	0/3/3/3
2	AN2	B	401	-	-	0/9/32/32	0/3/3/3
2	AN2	C	401	-	-	0/9/32/32	0/3/3/3
2	AN2	D	401	-	-	0/9/32/32	0/3/3/3
2	AN2	E	401	-	-	0/9/32/32	0/3/3/3
2	AN2	F	401	-	-	0/9/32/32	0/3/3/3
2	AN2	G	401	-	-	0/9/32/32	0/3/3/3
2	AN2	H	401	-	-	0/9/32/32	0/3/3/3
2	AN2	I	401	-	-	0/9/32/32	0/3/3/3
2	AN2	J	401	-	-	0/9/32/32	0/3/3/3
2	AN2	K	401	-	-	0/9/32/32	0/3/3/3
2	AN2	L	401	-	-	0/9/32/32	0/3/3/3

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	AN2	C2'-C1'	-2.84	1.49	1.53
2	H	401	AN2	PB-O2B	-2.76	1.49	1.56
2	E	401	AN2	PB-O2B	-2.58	1.49	1.56
2	A	401	AN2	PB-O2B	-2.35	1.50	1.56
2	H	401	AN2	C5-N7	-2.31	1.31	1.39

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	AN2	N3-C2-N1	-9.36	121.52	128.87
2	J	401	AN2	N3-C2-N1	-9.07	121.74	128.87
2	H	401	AN2	N3-C2-N1	-8.76	121.99	128.87
2	K	401	AN2	N3-C2-N1	-8.46	122.23	128.87
2	L	401	AN2	N3-C2-N1	-8.42	122.25	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	AN2	1	0
2	L	401	AN2	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/330 (92%)	-0.42	1 (0%) 94 93	18, 42, 95, 110	0
1	B	314/330 (95%)	-0.68	0 100 100	17, 33, 62, 84	0
1	C	313/330 (94%)	-0.66	0 100 100	18, 33, 64, 80	0
1	D	311/330 (94%)	-0.65	0 100 100	20, 34, 59, 77	0
1	E	311/330 (94%)	-0.68	0 100 100	16, 29, 54, 75	0
1	F	312/330 (94%)	-0.28	3 (0%) 84 81	22, 44, 81, 93	0
1	G	314/330 (95%)	-0.67	0 100 100	16, 31, 51, 67	0
1	H	311/330 (94%)	-0.65	1 (0%) 94 93	17, 30, 52, 90	0
1	I	313/330 (94%)	-0.41	6 (1%) 70 64	24, 43, 88, 121	0
1	J	314/330 (95%)	-0.49	0 100 100	27, 45, 71, 99	0
1	K	312/330 (94%)	-0.55	0 100 100	23, 39, 74, 90	0
1	L	311/330 (94%)	-0.57	0 100 100	24, 41, 70, 90	0
All	All	3742/3960 (94%)	-0.56	11 (0%) 94 93	16, 37, 72, 121	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	118	GLN	2.9
1	I	113	VAL	2.5
1	I	212	GLY	2.3
1	I	112	ILE	2.3
1	F	226	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NA	E	403	1/1	0.98	0.14	3.37	26,26,26,26	0
3	NA	B	403	1/1	0.98	0.15	2.73	31,31,31,31	0
3	NA	I	403	1/1	0.95	0.13	1.49	38,38,38,38	0
3	NA	H	403	1/1	0.99	0.12	0.63	21,21,21,21	0
2	AN2	B	401	27/27	0.97	0.12	0.18	26,31,38,40	0
3	NA	F	402	1/1	0.85	0.15	-0.03	50,50,50,50	0
2	AN2	C	401	27/27	0.98	0.12	-0.07	29,36,43,43	0
2	AN2	G	401	27/27	0.98	0.12	-0.13	24,26,31,34	0
2	AN2	E	401	27/27	0.98	0.12	-0.26	24,29,37,39	0
2	AN2	I	401	27/27	0.95	0.12	-0.26	35,39,48,50	0
2	AN2	A	401	27/27	0.97	0.14	-0.27	35,45,58,59	0
2	AN2	F	401	27/27	0.95	0.14	-0.31	32,48,56,63	0
2	AN2	H	401	27/27	0.98	0.10	-0.37	18,20,29,30	0
2	AN2	K	401	27/27	0.96	0.12	-0.40	42,46,60,60	0
2	AN2	D	401	27/27	0.97	0.12	-0.45	29,34,38,38	0
2	AN2	L	401	27/27	0.98	0.12	-0.53	33,37,51,54	0
2	AN2	J	401	27/27	0.98	0.11	-0.69	26,36,41,47	0
3	NA	C	402	1/1	0.95	0.09	-1.05	29,29,29,29	0
3	NA	K	402	1/1	0.95	0.09	-1.26	40,40,40,40	0
3	NA	C	403	1/1	0.97	0.09	-1.53	28,28,28,28	0
3	NA	F	403	1/1	0.94	0.08	-1.64	35,35,35,35	0
3	NA	J	402	1/1	0.98	0.11	-1.66	38,38,38,38	0
3	NA	L	403	1/1	0.91	0.08	-1.76	47,47,47,47	0
3	NA	G	402	1/1	0.98	0.08	-1.86	28,28,28,28	0
3	NA	H	402	1/1	0.94	0.08	-2.13	33,33,33,33	0
3	NA	D	403	1/1	0.99	0.07	-2.15	34,34,34,34	0
3	NA	L	402	1/1	0.97	0.04	-3.18	26,26,26,26	0
3	NA	D	402	1/1	0.98	0.06	-3.60	26,26,26,26	0
3	NA	G	403	1/1	0.96	0.08	-3.85	17,17,17,17	0
3	NA	A	402	1/1	0.98	0.05	-3.89	26,26,26,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NA	A	403	1/1	0.96	0.09	-3.92	31,31,31,31	0
3	NA	I	402	1/1	0.95	0.06	-4.02	33,33,33,33	0
3	NA	K	403	1/1	0.99	0.05	-4.61	37,37,37,37	0
3	NA	B	402	1/1	0.96	0.05	-5.55	22,22,22,22	0
3	NA	J	403	1/1	0.99	0.09	-5.62	27,27,27,27	0
3	NA	E	402	1/1	0.95	0.05	-7.36	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.