



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:14 AM GMT

PDB ID : 2C63
Title : 14-3-3 Protein Eta (Human) Complexed to Peptide
Authors : Elkins, J.M.; Yang, X.; Smee, C.E.A.; Johansson, C.; Sundstrom, M.; Edwards, A.; Weigelt, J.; Arrowsmith, C.; Doyle, D.A.
Deposited on : 2005-11-07
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	<div> <div>2%</div> <div>85% 9% 6%</div> </div>
1	B	247	<div> <div>5%</div> <div>86% 8% 6%</div> </div>
1	C	247	<div> <div>3%</div> <div>85% 9% 6%</div> </div>
1	D	247	<div> <div>2%</div> <div>88% 6% 6%</div> </div>
2	P	6	<div> <div>67% 33%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	Q	6	 67%33%
2	R	6	 50%17%33%
2	S	6	 17%67%33%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 PROTEIN ETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	0	0
			1838	1160	310	360	8			
1	B	233	Total	C	N	O	S	0	0	0
			1833	1157	308	360	8			
1	C	233	Total	C	N	O	S	0	0	0
			1850	1164	310	367	9			
1	D	233	Total	C	N	O	S	0	0	0
			1832	1155	307	362	8			

- Molecule 2 is a protein called CONSENSUS PEPTIDE FOR 14-3-3 PROTEINS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	4	Total	C	N	O	P	0	0	0
			34	20	4	9	1			
2	Q	4	Total	C	N	O	P	0	0	0
			34	20	4	9	1			
2	R	4	Total	C	N	O	P	0	0	0
			34	20	4	9	1			
2	S	4	Total	C	N	O	P	0	0	0
			34	20	4	9	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	92	Total	O	0	0
			92	92		
3	B	67	Total	O	0	0
			67	67		
3	C	113	Total	O	0	0
			113	113		
3	D	107	Total	O	0	0
			107	107		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	P	3	Total 3	O 3	0	0
3	Q	5	Total 5	O 5	0	0
3	R	4	Total 4	O 4	0	0
3	S	6	Total 6	O 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 14-3-3 PROTEIN ETA




• Molecule 1: 14-3-3 PROTEIN ETA



• Molecule 1: 14-3-3 PROTEIN ETA



• Molecule 1: 14-3-3 PROTEIN ETA



• Molecule 2: CONSENSUS PEPTIDE FOR 14-3-3 PROTEINS

Chain P:  67% 33%




- Molecule 2: CONSENSUS PEPTIDE FOR 14-3-3 PROTEINS

Chain Q:  67% 33%




- Molecule 2: CONSENSUS PEPTIDE FOR 14-3-3 PROTEINS

Chain R:  50% 17% 33%



- Molecule 2: CONSENSUS PEPTIDE FOR 14-3-3 PROTEINS

Chain S:  17% 67% 33%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.25Å 79.40Å 125.11Å 90.00° 95.15° 90.00°	Depositor
Resolution (Å)	125.00 – 2.15 58.02 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (125.00-2.15) 99.9 (58.02-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.184 , 0.248 0.193 , 0.250	Depositor DCC
R_{free} test set	1883 reflections (3.14%)	DCC
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 61897 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7886	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.79 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.0911e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/1865	0.76	2/2519 (0.1%)
1	B	0.73	0/1860	0.74	2/2514 (0.1%)
1	C	0.88	2/1877 (0.1%)	0.82	3/2535 (0.1%)
1	D	0.81	0/1859	0.79	1/2514 (0.0%)
2	P	0.77	0/23	0.57	0/28
2	Q	0.81	0/23	0.55	0/28
2	R	0.83	0/23	0.59	0/28
2	S	0.80	0/23	0.80	0/28
All	All	0.80	2/7553 (0.0%)	0.78	8/10194 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	75	ASN	CG-OD1	5.67	1.36	1.24
1	C	194	CYS	CB-SG	-5.12	1.73	1.81

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	LEU	CA-CB-CG	7.39	132.30	115.30
1	B	42	ARG	NE-CZ-NH2	-7.30	116.65	120.30
1	D	56	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	C	12	ARG	NE-CZ-NH2	-5.79	117.41	120.30
1	C	56	ARG	NE-CZ-NH1	5.31	122.95	120.30
1	C	19	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	A	196	LEU	CB-CG-CD1	5.15	119.76	111.00
1	B	42	ARG	NE-CZ-NH1	5.12	122.86	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1838	0	1788	9	0
1	B	1833	0	1772	10	0
1	C	1850	0	1796	6	0
1	D	1832	0	1763	7	0
2	P	34	0	31	0	0
2	Q	34	0	31	0	0
2	R	34	0	31	1	0
2	S	34	0	31	0	0
3	A	92	0	0	2	0
3	B	67	0	0	2	0
3	C	113	0	0	1	0
3	D	107	0	0	4	0
3	P	3	0	0	0	0
3	Q	5	0	0	0	0
3	R	4	0	0	0	0
3	S	6	0	0	0	0
All	All	7886	0	7243	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:227:ARG:O	1:C:231:THR:HG23	1.83	0.78
1:A:43:ASN:ND2	3:A:2017:HOH:O	2.26	0.68
1:C:50:LYS:NZ	2:R:4:SEP:O1P	2.30	0.64
1:C:187:GLN:NE2	3:C:2094:HOH:O	2.30	0.63
1:A:42:ARG:HG2	1:A:122:PHE:CE1	2.36	0.61
1:B:205:ILE:HD11	3:B:2063:HOH:O	2.05	0.56
1:D:74:GLY:O	3:D:2040:HOH:O	2.19	0.51
1:A:205:ILE:HD13	1:A:208:LEU:HD11	1.94	0.50
1:D:37:LEU:HB3	1:D:41:ASP:HB2	1.94	0.50
1:A:118:GLU:HA	1:A:165:MET:CE	2.42	0.49
1:C:117:TYR:CD1	1:C:164:GLN:HG3	2.48	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:50:LYS:HE2	3:D:2026:HOH:O	2.14	0.47
1:B:227:ARG:O	1:B:231:THR:HG23	2.14	0.46
1:B:221:LEU:HD12	3:B:2066:HOH:O	2.16	0.46
1:B:114:ASP:O	1:B:117:TYR:CE1	2.69	0.46
1:B:132:ARG:O	1:B:136:GLU:HG3	2.17	0.44
1:C:118:GLU:HA	1:C:165:MET:CE	2.47	0.44
1:B:188:ASN:O	1:B:190:PRO:HD3	2.19	0.43
1:D:144:ASN:HB3	3:D:2084:HOH:O	2.19	0.42
1:D:231:THR:O	1:D:235:SER:OG	2.36	0.42
1:B:33:LEU:HD23	1:D:106:LYS:HD2	2.02	0.42
1:A:113:ASN:C	1:A:115:PHE:H	2.23	0.42
1:A:199:GLN:HB3	3:A:2072:HOH:O	2.19	0.42
1:A:108:LEU:HD22	1:A:123:TYR:CZ	2.55	0.41
1:B:78:LYS:O	1:B:82:VAL:HG23	2.21	0.41
1:B:158:PHE:CE2	1:B:162:LYS:HE3	2.56	0.41
1:C:212:ASN:HB2	1:C:215:SER:OG	2.21	0.41
1:B:196:LEU:HA	1:B:199:GLN:HG2	2.03	0.41
1:A:117:TYR:CD1	1:A:164:GLN:HG3	2.56	0.40
1:D:3:ASP:HB2	3:D:2002:HOH:O	2.21	0.40
1:A:90:GLU:O	1:A:94:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/247 (94%)	229 (99%)	2 (1%)	0	100	100
1	B	231/247 (94%)	228 (99%)	3 (1%)	0	100	100
1	C	231/247 (94%)	229 (99%)	2 (1%)	0	100	100
1	D	231/247 (94%)	226 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	P	1/6 (17%)	1 (100%)	0	0	100	100
2	Q	1/6 (17%)	1 (100%)	0	0	100	100
2	R	1/6 (17%)	1 (100%)	0	0	100	100
2	S	1/6 (17%)	1 (100%)	0	0	100	100
All	All	928/1012 (92%)	916 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/215 (89%)	182 (95%)	9 (5%)	32	28
1	B	189/215 (88%)	185 (98%)	4 (2%)	61	65
1	C	195/215 (91%)	186 (95%)	9 (5%)	33	29
1	D	189/215 (88%)	184 (97%)	5 (3%)	54	55
2	P	3/4 (75%)	3 (100%)	0	100	100
2	Q	3/4 (75%)	3 (100%)	0	100	100
2	R	3/4 (75%)	3 (100%)	0	100	100
2	S	3/4 (75%)	3 (100%)	0	100	100
All	All	776/876 (89%)	749 (96%)	27 (4%)	43	41

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ARG
1	A	25	SER
1	A	50	LYS
1	A	56	ARG
1	A	139	SER
1	A	179	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	196	LEU
1	A	205	ILE
1	A	227	ARG
1	B	112	CYS
1	B	115	PHE
1	B	218	ASP
1	B	231	THR
1	C	3	ASP
1	C	10	ARG
1	C	25	SER
1	C	179	PHE
1	C	191	GLU
1	C	195	LEU
1	C	196	LEU
1	C	231	THR
1	C	235	SER
1	D	10	ARG
1	D	105	ASP
1	D	112	CYS
1	D	196	LEU
1	D	221	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	9	GLN
1	B	144	ASN
1	B	199	GLN
1	C	9	GLN
1	C	75	ASN
1	C	144	ASN
1	D	9	GLN
1	D	43	ASN
1	D	113	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SEP	P	4	2	8,9,10	1.31	2 (25%)	8,12,14	1.63	3 (37%)
2	SEP	Q	4	2	8,9,10	1.41	1 (12%)	8,12,14	1.71	3 (37%)
2	SEP	R	4	2	8,9,10	1.28	1 (12%)	8,12,14	1.32	1 (12%)
2	SEP	S	4	2	8,9,10	1.27	0	8,12,14	1.46	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	P	4	2	-	0/6/8/10	0/0/0/0
2	SEP	Q	4	2	-	0/6/8/10	0/0/0/0
2	SEP	R	4	2	-	0/6/8/10	0/0/0/0
2	SEP	S	4	2	-	0/6/8/10	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	4	SEP	P-O3P	2.01	1.61	1.54
2	R	4	SEP	P-O1P	2.40	1.59	1.51
2	P	4	SEP	P-O1P	2.57	1.59	1.51
2	Q	4	SEP	P-O1P	3.36	1.62	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	4	SEP	O3P-P-O1P	-2.60	102.21	110.58
2	P	4	SEP	O-C-CA	-2.05	120.16	125.49
2	S	4	SEP	O3P-P-O2P	2.13	115.48	107.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	4	SEP	OG-CB-CA	2.21	110.16	108.27
2	P	4	SEP	O3P-P-O2P	2.26	115.98	107.38
2	Q	4	SEP	OG-CB-CA	2.33	110.26	108.27
2	P	4	SEP	O2P-P-O1P	2.39	118.26	110.58
2	Q	4	SEP	O3P-P-O2P	2.79	118.01	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	R	4	SEP	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	233/247 (94%)	0.44	5 (2%) 67 74	34, 41, 55, 72	0
1	B	233/247 (94%)	0.52	12 (5%) 31 41	35, 41, 56, 74	0
1	C	233/247 (94%)	0.46	8 (3%) 49 59	34, 41, 56, 77	0
1	D	233/247 (94%)	0.34	5 (2%) 67 74	34, 41, 55, 75	0
2	P	3/6 (50%)	0.84	0 100 100	34, 34, 45, 49	0
2	Q	3/6 (50%)	1.10	0 100 100	34, 34, 46, 49	0
2	R	3/6 (50%)	0.21	0 100 100	33, 33, 45, 48	0
2	S	3/6 (50%)	0.44	1 (33%) 0 1	33, 33, 45, 49	0
All	All	944/1012 (93%)	0.45	31 (3%) 50 60	33, 41, 56, 77	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	74	GLY	4.7
1	B	73	ASP	4.4
1	B	235	SER	4.3
1	D	232	LEU	4.0
1	C	235	SER	3.9
1	C	115	PHE	3.7
1	C	71	MET	3.4
1	A	235	SER	3.3
1	B	117	TYR	3.3
1	D	235	SER	3.3
1	B	72	ALA	3.3
1	B	114	ASP	3.2
1	B	70	THR	3.1
1	C	117	TYR	2.9
1	D	233	TRP	2.9
1	D	193	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	112	CYS	2.5
1	B	234	THR	2.4
1	B	64	SER	2.4
1	C	112	CYS	2.4
1	B	79	LEU	2.3
1	A	72	ALA	2.3
1	A	75	ASN	2.2
1	B	113	ASN	2.2
1	B	63	ILE	2.2
1	A	3	ASP	2.1
1	C	114	ASP	2.1
1	C	233	TRP	2.1
1	B	74	GLY	2.1
2	S	3	ILE	2.1
1	A	232	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SEP	R	4	10/11	0.98	0.07	-	24,31,34,34	0
2	SEP	S	4	10/11	0.98	0.07	-	27,31,34,34	0
2	SEP	P	4	10/11	0.98	0.09	-	24,31,34,34	0
2	SEP	Q	4	10/11	0.98	0.08	-	26,34,35,35	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.