



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:45 PM GMT

PDB ID : 1CBO  
Title : CHOLESTEROL OXIDASE FROM STREPTOMYCES HIS447ASN MUTANT  
Authors : Vrielink, A.; Yue, Q.K.  
Deposited on : 1999-02-26  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	504	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4185 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (CHOLESTEROL OXIDASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	498	3837	2423	666	732	16	0	8	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	ASN	HIS	SEE REMARK 999	UNP P12676

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	295	Total 295	O 295	0	0



Note EDS was not executed.

- Chain A:  87% 10%
- | ASP | ASN  | GLY  | ... |
|-----|------|------|-----|
| G9  | M46  | Q47  |     |
|     | G57  | M58  |     |
|     | L59  | M60  |     |
|     | M86  | Q103 |     |
|     | V108 | M119 |     |
|     | G120 | G121 |     |
|     | M122 | A123 |     |
|     | V124 | K127 |     |
|     | R156 | V157 |     |
|     | M158 | H159 |     |
|     | I160 | K172 |     |
|     | V176 | Q180 |     |
|     | K183 | V191 |     |
|     | T215 | E216 |     |
|     | V217 | I218 |     |
|     | Q226 | D229 |     |
|     | L233 |      |     |
- |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |     |     |     |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|-----|-----|
| T295 | V299 | W351 | N353 | S357 | E361 | Y390 | R403 | M406 | N424 | T435 | Q436 | D474 | G475 | S476 | G480 | S481 | V482 | M485 | P486 | F487 | R486 | N497 | V498 | E499 | R500 | I501 | I502 | R503 | D505 | V506 | THR | ALA | SER |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|-----|-----|

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.58 Å   74.15 Å   63.74 Å 90.00°   105.08°   90.00°	Depositor
Resolution (Å)	20.00 – 1.80	Depositor
% Data completeness (in resolution range)	90.1 (20.00-1.80)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.843	Depositor
R, $R_{free}$	0.159 , 0.194	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4185	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/3966	0.63	1/5392 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	486	PRO	N-CA-C	6.36	128.63	112.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3837	0	3743	36	0
2	A	53	0	31	3	0
3	A	295	0	0	6	0
All	All	4185	0	3774	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:VAL:HB	3:A:783:HOH:O	1.82	0.78
1:A:299[B]:VAL:HG12	1:A:390:TYR:HB2	1.65	0.78
1:A:103:GLN:NE2	1:A:403:ARG:H	1.86	0.71
1:A:361:GLU:HB3	3:A:676:HOH:O	1.96	0.65
1:A:498:VAL:O	1:A:502:ILE:HG12	2.01	0.59
1:A:435:THR:HG22	1:A:436:GLN:H	1.67	0.58
1:A:160:ILE:HG21	3:A:804:HOH:O	2.03	0.58
1:A:119:ASN:C	1:A:119:ASN:HD22	2.08	0.56
1:A:183:LYS:HD2	1:A:424:ASN:OD1	2.09	0.53
1:A:60:ASN:N	1:A:60:ASN:HD22	2.08	0.51
1:A:217:VAL:HG13	1:A:218:ILE:N	2.26	0.50
1:A:60:ASN:H	1:A:60:ASN:HD22	1.59	0.48
1:A:59:LEU:HD13	1:A:215[B]:THR:HG21	1.95	0.48
1:A:127:LYS:HE3	1:A:351:TRP:O	2.14	0.48
1:A:57:GLY:H	1:A:60:ASN:HD21	1.62	0.47
1:A:500:ARG:HH21	1:A:504:GLN:HE21	1.63	0.47
1:A:119:ASN:HB2	2:A:510:FAD:C4X	2.45	0.47
1:A:122:MET:SD	1:A:361:GLU:OE2	2.73	0.46
2:A:510:FAD:H2B	3:A:599:HOH:O	2.16	0.45
1:A:253:ILE:HG12	1:A:263:LEU:HD22	1.99	0.45
1:A:406:ASN:HD22	1:A:406:ASN:HA	1.56	0.44
1:A:124:VAL:HG22	1:A:191:VAL:HG13	2.00	0.44
1:A:58:MET:HB2	1:A:215[B]:THR:HG22	1.99	0.44
1:A:496:ARG:NH1	1:A:497:ASN:HD21	2.16	0.44
1:A:295:THR:O	1:A:299[A]:VAL:HG13	2.17	0.43
1:A:86:ASN:ND2	3:A:657:HOH:O	2.51	0.43
1:A:485:ASN:HB3	2:A:510:FAD:C2	2.49	0.43
1:A:158:ASN:N	1:A:158:ASN:HD22	2.18	0.42
1:A:351:TRP:CD2	1:A:357:SER:HB3	2.54	0.42
1:A:180:GLN:HG3	3:A:661:HOH:O	2.20	0.42
1:A:229:ASP:HA	1:A:233:LEU:HB2	2.01	0.42
1:A:352:ASP:O	1:A:353:ASN:HB2	2.20	0.41
1:A:119:ASN:ND2	1:A:121:GLY:H	2.19	0.41
1:A:474:ASP:HB2	1:A:475:GLY:H	1.73	0.41
1:A:172:LYS:O	1:A:176:VAL:HG23	2.21	0.41
1:A:291:SER:HA	1:A:476:SER:HB3	2.03	0.41
1:A:156:ARG:HH11	1:A:226:GLN:HE21	1.69	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	504/504 (100%)	484 (96%)	18 (4%)	2 (0%)	39	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	217	VAL
1	A	480	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	403/401 (100%)	393 (98%)	10 (2%)	55	39

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	47	GLN
1	A	60	ASN
1	A	119	ASN
1	A	158	ASN
1	A	361	GLU
1	A	406	ASN
1	A	435	THR
1	A	482	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	487	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	60	ASN
1	A	86	ASN
1	A	103	GLN
1	A	119	ASN
1	A	226	GLN
1	A	249	GLN
1	A	406	ASN
1	A	411	ASN
1	A	469	ASN
1	A	497	ASN
1	A	504	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	510	-	48,58,58	1.67	8 (16%)	54,89,89	1.80	11 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	510	-	-	0/30/50/50	0/6/6/6

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	510	FAD	PA-O2A	-4.61	1.35	1.54
2	A	510	FAD	C10-N10	-4.11	1.34	1.39
2	A	510	FAD	P-O2P	-3.97	1.38	1.54
2	A	510	FAD	C9A-N10	2.00	1.41	1.38
2	A	510	FAD	C4-N3	2.53	1.37	1.33
2	A	510	FAD	C4X-C10	3.05	1.46	1.41
2	A	510	FAD	O5'-C5'	3.58	1.59	1.44
2	A	510	FAD	O4B-C1B	3.78	1.46	1.41

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	510	FAD	C4X-C4-N3	-5.29	116.36	123.59
2	A	510	FAD	O4B-C1B-N9A	-3.23	101.34	108.10
2	A	510	FAD	C4B-O4B-C1B	-3.22	106.18	109.72
2	A	510	FAD	O5B-PA-O1A	-3.20	97.18	109.62
2	A	510	FAD	N3A-C2A-N1A	-2.39	127.06	128.89
2	A	510	FAD	O5'-P-O1P	-2.25	100.87	109.62
2	A	510	FAD	C6-C5X-N5	-2.13	116.22	118.96
2	A	510	FAD	O3B-C3B-C4B	2.00	117.06	111.05
2	A	510	FAD	C2A-N1A-C6A	2.10	122.53	118.77
2	A	510	FAD	O3P-PA-O5B	2.53	109.66	102.94
2	A	510	FAD	C4-N3-C2	7.07	121.36	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	510	FAD	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.