



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:45 PM GMT

PDB ID : 1CCB
Title : THE ASP-HIS-FE TRIAD OF CYTOCHROME C PEROXIDASE CONTROLS THE REDUCTION POTENTIAL, ELECTRONIC STRUCTURE, AND COUPLING OF THE TRYPTOPHAN FREE-RADICAL TO THE HEME
Authors : Goodin, D.B.; Mcree, D.E.
Deposited on : 1993-01-04
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	297	 79% 15% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2947 atoms, of which 505 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	291	2854	1502	505	390	451	6	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ILE	THR	CONFLICT	UNP P00431
A	152	GLY	ASP	CONFLICT	UNP P00431
A	235	GLU	ASP	CONFLICT	UNP P00431

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	50	Total	O	0	0
			50	50		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CYTOCHROME C PEROXIDASE

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.20 Å 74.30 Å 45.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.190 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2947	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/2415	1.62	42/3269 (1.3%)

There are no bond length outliers.

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	ARG	NE-CZ-NH2	-9.57	115.52	120.30
1	A	127	ARG	NE-CZ-NH1	9.40	125.00	120.30
1	A	51	TRP	CD1-CG-CD2	9.31	113.75	106.30
1	A	223	TRP	CD1-CG-CD2	9.18	113.65	106.30
1	A	191	TRP	CD1-CG-CD2	8.88	113.41	106.30
1	A	72	ARG	NE-CZ-NH1	8.77	124.68	120.30
1	A	31	ARG	NE-CZ-NH1	8.59	124.60	120.30
1	A	251	TYR	CB-CG-CD2	-8.58	115.85	121.00
1	A	51	TRP	CE2-CD2-CG	-8.52	100.48	107.30
1	A	101	TRP	CD1-CG-CD2	8.43	113.05	106.30
1	A	51	TRP	CB-CG-CD1	-8.21	116.33	127.00
1	A	191	TRP	CE2-CD2-CG	-8.18	100.75	107.30
1	A	101	TRP	CE2-CD2-CG	-7.87	101.00	107.30
1	A	223	TRP	CE2-CD2-CG	-7.85	101.02	107.30
1	A	166	ARG	NE-CZ-NH2	-7.58	116.51	120.30
1	A	126	TRP	CD1-CG-CD2	7.56	112.35	106.30
1	A	48	ARG	NE-CZ-NH1	7.46	124.03	120.30
1	A	163	MET	CG-SD-CE	-7.26	88.58	100.20
1	A	31	ARG	NE-CZ-NH2	-7.11	116.75	120.30
1	A	143	ARG	NE-CZ-NH1	7.07	123.83	120.30
1	A	211	TRP	CD1-CG-CD2	6.98	111.88	106.30
1	A	51	TRP	CG-CD2-CE3	6.97	140.18	133.90
1	A	166	ARG	NE-CZ-NH1	6.93	123.77	120.30
1	A	126	TRP	CE2-CD2-CG	-6.65	101.98	107.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	130	ARG	CG-CD-NE	-6.58	97.99	111.80
1	A	211	TRP	CE2-CD2-CG	-6.44	102.15	107.30
1	A	223	TRP	CG-CD1-NE1	-6.25	103.85	110.10
1	A	51	TRP	CG-CD1-NE1	-6.23	103.87	110.10
1	A	162	ASN	CA-C-N	-6.23	103.50	117.20
1	A	229	TYR	CB-CG-CD2	-6.12	117.33	121.00
1	A	57	TRP	CD1-CG-CD2	6.06	111.15	106.30
1	A	57	TRP	CE2-CD2-CG	-6.05	102.46	107.30
1	A	72	ARG	NE-CZ-NH2	-5.73	117.43	120.30
1	A	48	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	A	101	TRP	CB-CG-CD1	-5.63	119.69	127.00
1	A	153	TYR	CB-CG-CD2	-5.55	117.67	121.00
1	A	120	GLN	CA-CB-CG	5.48	125.45	113.40
1	A	101	TRP	CG-CD2-CE3	5.42	138.78	133.90
1	A	101	TRP	CG-CD1-NE1	-5.30	104.80	110.10
1	A	191	TRP	CG-CD1-NE1	-5.21	104.89	110.10
1	A	14	ARG	NE-CZ-NH2	-5.13	117.73	120.30
1	A	127	ARG	NE-CZ-NH2	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2349	505	2220	29	0
2	A	43	0	30	1	0
3	A	50	0	0	1	0
All	All	2442	505	2250	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:ASN:H	1:A:195:ASN:HD22	1.34	0.75
1:A:172:MET:HE1	2:A:295:HEM:HHC	1.82	0.60
1:A:195:ASN:H	1:A:195:ASN:ND2	2.01	0.59
1:A:74:LYS:HE2	1:A:74:LYS:H	1.68	0.57
1:A:20:GLN:HE22	1:A:287:LYS:N	2.04	0.56
1:A:195:ASN:HD22	1:A:195:ASN:N	2.04	0.54
1:A:20:GLN:HE22	1:A:287:LYS:H	1.55	0.53
1:A:212:LYS:HE3	1:A:224:ASP:O	2.07	0.53
1:A:271:GLU:HA	1:A:274:ILE:HD12	1.91	0.51
1:A:239:ILE:HG22	1:A:245:LEU:HD13	1.93	0.51
1:A:20:GLN:NE2	1:A:287:LYS:H	2.13	0.47
1:A:212:LYS:HE3	1:A:225:SER:HA	1.97	0.47
1:A:212:LYS:HE2	1:A:226:LYS:HD2	1.96	0.47
1:A:168:VAL:O	1:A:172:MET:HG2	2.16	0.46
1:A:206:LEU:HD13	1:A:231:MET:SD	2.55	0.46
1:A:130:ARG:NH1	1:A:270:LEU:O	2.48	0.46
1:A:146:ASP:O	1:A:234:THR:HG21	2.15	0.46
1:A:130:ARG:NE	3:A:326:HOH:O	2.44	0.46
1:A:213:LEU:HD11	1:A:221:GLU:HB3	1.96	0.46
1:A:125:PRO:HG3	1:A:285:ILE:HD11	2.00	0.44
1:A:51:TRP:CD1	1:A:52:HIS:HD2	2.35	0.44
1:A:125:PRO:HG3	1:A:285:ILE:CD1	2.48	0.43
1:A:264:LYS:HB2	1:A:264:LYS:HE3	1.89	0.43
1:A:149:LYS:O	1:A:234:THR:HA	2.19	0.42
1:A:163:MET:HA	1:A:167:GLU:OE1	2.20	0.42
1:A:167:GLU:O	1:A:171:LEU:HB2	2.20	0.42
1:A:14:ARG:HG3	1:A:101:TRP:CE3	2.56	0.41
1:A:149:LYS:H	1:A:234:THR:HG23	1.86	0.40
1:A:236:TYR:CE2	1:A:239:ILE:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/297 (97%)	278 (96%)	10 (4%)	1 (0%)	46 45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	247/255 (97%)	233 (94%)	14 (6%)	25 22

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	28	LEU
1	A	31	ARG
1	A	35	GLU
1	A	47	VAL
1	A	51	TRP
1	A	97	LYS
1	A	156	THR
1	A	159	GLN
1	A	166	ARG
1	A	171	LEU
1	A	195	ASN
1	A	212	LYS
1	A	287	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	24	ASN
1	A	87	ASN
1	A	195	ASN
1	A	208	ASN
1	A	255	GLN
1	A	292	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	295	1,3	30,50,50	3.99	13 (43%)	24,82,82	2.58	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	295	1,3	-	0/10/54/54	0/0/8/8

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	295	HEM	C3B-C4B	-13.12	1.40	1.51
2	A	295	HEM	C3D-C4D	-9.30	1.39	1.51
2	A	295	HEM	C2C-C1C	-6.91	1.39	1.52
2	A	295	HEM	C3C-CAC	-6.03	1.40	1.51
2	A	295	HEM	C3B-CAB	-5.94	1.40	1.51
2	A	295	HEM	C2D-C3D	-5.14	1.39	1.54
2	A	295	HEM	C2D-C1D	-3.97	1.38	1.51
2	A	295	HEM	C2B-C1B	-3.59	1.40	1.51
2	A	295	HEM	FE-ND	2.03	2.08	1.97
2	A	295	HEM	CHD-C4C	2.22	1.41	1.36
2	A	295	HEM	FE-NC	2.54	2.05	1.95
2	A	295	HEM	CBB-CAB	2.57	1.44	1.29
2	A	295	HEM	CBC-CAC	3.03	1.46	1.29

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	295	HEM	CAA-C2A-C1A	-2.10	124.72	127.01
2	A	295	HEM	CMD-C2D-C3D	2.36	124.77	114.35
2	A	295	HEM	C3B-CAB-CBB	2.59	128.43	124.46
2	A	295	HEM	C2D-C3D-C4D	2.89	106.40	101.50
2	A	295	HEM	C4B-CHC-C1C	3.15	131.09	125.82
2	A	295	HEM	CAD-C3D-C2D	4.35	125.72	113.22
2	A	295	HEM	CAD-C3D-C4D	4.37	127.88	112.47
2	A	295	HEM	CMB-C2B-C3B	5.53	130.35	116.53
2	A	295	HEM	CMC-C2C-C3C	5.88	131.20	116.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	295	HEM	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.