



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:46 PM GMT

PDB ID : 1CDL  
Title : TARGET ENZYME RECOGNITION BY CALMODULIN: 2.4  
ANGSTROMS STRUCTURE OF A CALMODULIN-PEPTIDE COM-  
PLEX  
Authors : Meador, W.E.; Quiococho, F.A.  
Deposited on : 1993-10-08  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	147	
1	B	147	
1	C	147	
1	D	147	
2	E	20	
2	F	20	
2	G	20	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	20	 A horizontal bar chart showing the quality of chain H. The bar is divided into four segments: green (50%), yellow (35%), orange (5%), and grey (10%). The percentages are labeled below the bar.

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CALMODULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	142	Total	C	N	O	S	0	0	0
			1097	672	177	239	9			
1	B	142	Total	C	N	O	S	0	0	0
			1112	680	178	245	9			
1	C	142	Total	C	N	O	S	0	0	0
			1106	676	178	243	9			
1	D	142	Total	C	N	O	S	0	0	0
			1087	664	177	237	9			

- Molecule 2 is a protein called CALCIUM/CALMODULIN-DEPENDENT PROTEIN KINASE TYPE II ALPHA CHAIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	19	Total	C	N	O	0	0	0
			156	95	37	24			
2	F	19	Total	C	N	O	0	0	0
			156	95	37	24			
2	G	20	Total	C	N	O	0	0	0
			154	94	35	25			
2	H	18	Total	C	N	O	0	0	0
			133	81	29	23			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	4	Total	Ca	0	0
			4	4		
3	A	4	Total	Ca	0	0
			4	4		
3	D	4	Total	Ca	0	0
			4	4		
3	C	4	Total	Ca	0	0
			4	4		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	72	Total 72	O 72	0	0
4	B	83	Total 83	O 83	0	0
4	C	50	Total 50	O 50	0	0
4	D	46	Total 46	O 46	0	0
4	E	13	Total 13	O 13	0	0
4	F	4	Total 4	O 4	0	0
4	G	1	Total 1	O 1	0	0
4	H	1	Total 1	O 1	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: CALMODULIN




#### • Molecule 1: CALMODULIN




#### • Molecule 1: CALMODULIN



#### • Molecule 1: CALMODULIN






- Molecule 2: CALCIUM/CALMODULIN-DEPENDENT PROTEIN KINASE TYPE II ALPHA CHAIN



- Molecule 2: CALCIUM/CALMODULIN-DEPENDENT PROTEIN KINASE TYPE II ALPHA CHAIN



- Molecule 2: CALCIUM/CALMODULIN-DEPENDENT PROTEIN KINASE TYPE II ALPHA CHAIN



- Molecule 2: CALCIUM/CALMODULIN-DEPENDENT PROTEIN KINASE TYPE II ALPHA CHAIN



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	166.10 Å   104.90 Å   43.32 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	10.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.00)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.208 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5287	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1109	0.67	0/1491
1	B	0.47	0/1124	0.70	1/1511 (0.1%)
1	C	0.45	0/1118	0.63	0/1502
1	D	0.42	0/1099	0.66	0/1476
2	E	0.45	0/158	0.65	0/207
2	F	0.45	0/158	0.50	0/207
2	G	0.44	0/156	0.58	0/205
2	H	0.37	0/135	0.53	0/178
All	All	0.45	0/5057	0.65	1/6777 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	95	ASP	CB-CG-OD2	5.04	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1097	0	1004	40	0
1	B	1112	0	1024	33	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1106	0	1018	36	0
1	D	1087	0	992	44	0
2	E	156	0	166	10	0
2	F	156	0	166	9	0
2	G	154	0	159	8	0
2	H	133	0	129	6	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0
3	C	4	0	0	0	0
3	D	4	0	0	0	0
4	A	72	0	0	6	0
4	B	83	0	0	3	0
4	C	50	0	0	2	0
4	D	46	0	0	2	0
4	E	13	0	0	3	0
4	F	4	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
All	All	5287	0	4658	164	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 164 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:6:GLU:HA	4:D:184:HOH:O	1.68	0.94
1:A:144:MET:SD	2:E:800:TRP:HB3	2.11	0.90
1:C:106:ARG:HG3	1:C:121:VAL:HG11	1.63	0.80
1:C:100:ILE:HB	1:C:136:VAL:HG13	1.66	0.77
1:D:136:VAL:HG13	1:D:140:GLU:HB3	1.66	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/147 (95%)	131 (94%)	7 (5%)	2 (1%)	14	6
1	B	140/147 (95%)	137 (98%)	3 (2%)	0	100	100
1	C	140/147 (95%)	129 (92%)	7 (5%)	4 (3%)	6	2
1	D	140/147 (95%)	124 (89%)	13 (9%)	3 (2%)	9	3
2	E	17/20 (85%)	15 (88%)	2 (12%)	0	100	100
2	F	17/20 (85%)	17 (100%)	0	0	100	100
2	G	18/20 (90%)	16 (89%)	2 (11%)	0	100	100
2	H	16/20 (80%)	15 (94%)	1 (6%)	0	100	100
All	All	628/668 (94%)	584 (93%)	35 (6%)	9 (1%)	14	6

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	56	ASP
1	D	129	ASP
1	C	129	ASP
1	D	145	MET
1	A	93	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/125 (93%)	109 (94%)	7 (6%)	24	17
1	B	120/125 (96%)	105 (88%)	15 (12%)	6	3
1	C	119/125 (95%)	107 (90%)	12 (10%)	9	5
1	D	115/125 (92%)	102 (89%)	13 (11%)	7	4
2	E	15/15 (100%)	15 (100%)	0	100	100
2	F	15/15 (100%)	13 (87%)	2 (13%)	5	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	G	14/15 (93%)	13 (93%)	1 (7%)	18	12
2	H	12/15 (80%)	10 (83%)	2 (17%)	3	1
All	All	526/560 (94%)	474 (90%)	52 (10%)	10	5

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	801	GLN
1	C	64	ASP
1	D	140	GLU
1	C	5	THR
1	C	24	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	111	ASN
1	C	137	ASN
1	D	143	GLN
1	B	49	GLN
1	D	49	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.