



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:54 AM GMT

PDB ID : 3CM7
Title : Crystal Structure of XIAP-BIR3 domain in complex with Smac-mimetic compound, Smac005
Authors : Mastrangelo, E.; Cossu, F.; Milani, M.
Deposited on : 2008-03-21
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	130	<div> <div>2%</div> <div>55% 20% • 22%</div> </div>
1	B	130	<div> <div>2%</div> <div>52% 25% • 22%</div> </div>
1	C	130	<div> <div>2%</div> <div>55% 20% • • 22%</div> </div>
1	D	130	<div> <div>%</div> <div>61% 15% • 23%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	A	6	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3532 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Baculoviral IAP repeat-containing protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	102	Total	C	N	O	S	0	0	0
			831	531	142	153	5			
1	A	101	Total	C	N	O	S	0	0	0
			825	528	141	151	5			
1	B	101	Total	C	N	O	S	0	0	0
			825	528	141	151	5			
1	D	100	Total	C	N	O	S	0	0	0
			814	522	137	150	5			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	227	MET	-	EXPRESSION TAG	UNP P98170
C	228	ALA	-	EXPRESSION TAG	UNP P98170
C	229	SER	-	EXPRESSION TAG	UNP P98170
C	230	MET	-	EXPRESSION TAG	UNP P98170
C	231	THR	-	EXPRESSION TAG	UNP P98170
C	232	GLY	-	EXPRESSION TAG	UNP P98170
C	233	GLY	-	EXPRESSION TAG	UNP P98170
C	234	GLN	-	EXPRESSION TAG	UNP P98170
C	235	GLN	-	EXPRESSION TAG	UNP P98170
C	236	MET	-	EXPRESSION TAG	UNP P98170
C	237	GLY	-	EXPRESSION TAG	UNP P98170
C	238	ARG	-	EXPRESSION TAG	UNP P98170
C	239	GLY	-	EXPRESSION TAG	UNP P98170
C	240	SER	-	EXPRESSION TAG	UNP P98170
A	227	MET	-	EXPRESSION TAG	UNP P98170
A	228	ALA	-	EXPRESSION TAG	UNP P98170
A	229	SER	-	EXPRESSION TAG	UNP P98170
A	230	MET	-	EXPRESSION TAG	UNP P98170
A	231	THR	-	EXPRESSION TAG	UNP P98170
A	232	GLY	-	EXPRESSION TAG	UNP P98170
A	233	GLY	-	EXPRESSION TAG	UNP P98170

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	234	GLN	-	EXPRESSION TAG	UNP P98170
A	235	GLN	-	EXPRESSION TAG	UNP P98170
A	236	MET	-	EXPRESSION TAG	UNP P98170
A	237	GLY	-	EXPRESSION TAG	UNP P98170
A	238	ARG	-	EXPRESSION TAG	UNP P98170
A	239	GLY	-	EXPRESSION TAG	UNP P98170
A	240	SER	-	EXPRESSION TAG	UNP P98170
B	227	MET	-	EXPRESSION TAG	UNP P98170
B	228	ALA	-	EXPRESSION TAG	UNP P98170
B	229	SER	-	EXPRESSION TAG	UNP P98170
B	230	MET	-	EXPRESSION TAG	UNP P98170
B	231	THR	-	EXPRESSION TAG	UNP P98170
B	232	GLY	-	EXPRESSION TAG	UNP P98170
B	233	GLY	-	EXPRESSION TAG	UNP P98170
B	234	GLN	-	EXPRESSION TAG	UNP P98170
B	235	GLN	-	EXPRESSION TAG	UNP P98170
B	236	MET	-	EXPRESSION TAG	UNP P98170
B	237	GLY	-	EXPRESSION TAG	UNP P98170
B	238	ARG	-	EXPRESSION TAG	UNP P98170
B	239	GLY	-	EXPRESSION TAG	UNP P98170
B	240	SER	-	EXPRESSION TAG	UNP P98170
D	227	MET	-	EXPRESSION TAG	UNP P98170
D	228	ALA	-	EXPRESSION TAG	UNP P98170
D	229	SER	-	EXPRESSION TAG	UNP P98170
D	230	MET	-	EXPRESSION TAG	UNP P98170
D	231	THR	-	EXPRESSION TAG	UNP P98170
D	232	GLY	-	EXPRESSION TAG	UNP P98170
D	233	GLY	-	EXPRESSION TAG	UNP P98170
D	234	GLN	-	EXPRESSION TAG	UNP P98170
D	235	GLN	-	EXPRESSION TAG	UNP P98170
D	236	MET	-	EXPRESSION TAG	UNP P98170
D	237	GLY	-	EXPRESSION TAG	UNP P98170
D	238	ARG	-	EXPRESSION TAG	UNP P98170
D	239	GLY	-	EXPRESSION TAG	UNP P98170
D	240	SER	-	EXPRESSION TAG	UNP P98170

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0

Continued on next page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0

- [illegible]

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total 36	C 28	N 4	O 4	0	0
3	A	1	Total 36	C 28	N 4	O 4	0	0
3	B	1	Total 36	C 28	N 4	O 4	0	0
3	D	1	Total 36	C 28	N 4	O 4	0	0
3	D	1	Total 36	C 28	N 4	O 4	0	0
3	D	1	Total 36	C 28	N 4	O 4	0	0

- 

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total 4	O 4	0	0
4	B	6	Total 6	O 6	0	0
4	C	3	Total 3	O 3	0	0
4	D	4	Total 4	O 4	0	0

L348	
L352	
V353	
ARG	
THR	
THR	

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.39Å 92.39Å 175.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.10 39.57 – 3.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (40.00-3.10) 100.0 (39.57-3.10)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.233 , 0.300 0.232 , 0.294	Depositor DCC
R_{free} test set	725 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	51.3	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 14399 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3532	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, X22

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/851	0.56	0/1153
1	B	0.45	0/851	0.56	0/1153
1	C	0.54	1/857 (0.1%)	0.53	0/1161
1	D	0.42	0/840	0.55	0/1139
All	All	0.47	1/3399 (0.0%)	0.55	0/4606

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	253	SER	CB-OG	9.67	1.54	1.42

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	256	LEU	Peptide
1	C	343	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	825	0	775	16	1
1	B	825	0	774	20	1
1	C	831	0	780	24	0
1	D	814	0	761	11	0
2	A	1	0	0	2	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	36	0	36	2	0
3	B	36	0	36	2	0
3	C	36	0	36	2	0
3	D	108	0	108	5	0
4	A	4	0	0	0	0
4	B	6	0	0	0	0
4	C	3	0	0	0	0
4	D	4	0	0	0	0
All	All	3532	0	3306	71	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:VAL:HG21	1:A:310:TRP:HB3	1.61	0.81
1:A:308:THR:O	3:A:600:X22:HB	1.80	0.80
1:C:297:LYS:HG2	1:C:308:THR:HG23	1.62	0.80
1:C:279:VAL:HG21	1:C:310:TRP:HB3	1.62	0.79
1:A:327:CYS:SG	2:A:6:ZN:ZN	1.73	0.77

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:337:GLU:OE2	1:B:258:ARG:NH1[6_544]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/130 (76%)	88 (89%)	10 (10%)	1 (1%)	19	58
1	B	99/130 (76%)	82 (83%)	15 (15%)	2 (2%)	9	38
1	C	100/130 (77%)	91 (91%)	8 (8%)	1 (1%)	19	58
1	D	98/130 (75%)	90 (92%)	8 (8%)	0	100	100
All	All	396/520 (76%)	351 (89%)	41 (10%)	4 (1%)	19	58

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	312	PRO
1	B	273	GLY
1	B	257	PRO
1	A	273	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/109 (79%)	74 (86%)	12 (14%)	4	18
1	B	86/109 (79%)	80 (93%)	6 (7%)	19	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	87/109 (80%)	78 (90%)	9 (10%)	9	32
1	D	85/109 (78%)	78 (92%)	7 (8%)	14	47
All	All	344/436 (79%)	310 (90%)	34 (10%)	10	34

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	308	THR
1	A	337	GLU
1	D	345	THR
1	A	327	CYS
1	C	344	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	259	ASN
1	B	343	HIS
1	D	340	ASN
1	A	343	HIS
1	D	336	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	X22	A	600	-	37,39,39	0.59	0	44,54,54	1.36	6 (13%)
3	X22	B	600	-	37,39,39	0.70	0	44,54,54	1.33	4 (9%)
3	X22	C	600	-	37,39,39	0.71	0	44,54,54	1.50	7 (15%)
3	X22	D	600	-	37,39,39	0.86	1 (2%)	44,54,54	1.59	8 (18%)
3	X22	D	700	-	37,39,39	1.37	4 (10%)	44,54,54	2.66	12 (27%)
3	X22	D	701	-	37,39,39	0.82	1 (2%)	44,54,54	1.91	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	X22	A	600	-	-	0/28/58/58	0/4/4/4
3	X22	B	600	-	-	0/28/58/58	0/4/4/4
3	X22	C	600	-	-	0/28/58/58	0/4/4/4
3	X22	D	600	-	-	0/28/58/58	0/4/4/4
3	X22	D	700	-	-	0/28/58/58	0/4/4/4
3	X22	D	701	-	-	0/28/58/58	0/4/4/4

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	600	X22	CBA-NBJ	-2.87	1.32	1.35
3	D	701	X22	CBA-NBJ	-2.87	1.32	1.35
3	D	700	X22	CBB-CBH	2.73	1.56	1.52
3	D	700	X22	CBC-CBH	2.86	1.56	1.52
3	D	700	X22	CBG-NBJ	3.09	1.50	1.46

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	701	X22	CBB-CBH-NAW	-5.52	100.87	111.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	600	X22	CBC-CBH-NAW	-4.04	103.65	111.18
3	D	700	X22	CBE-CBI-NAX	-3.66	106.06	110.70
3	D	700	X22	OAD-CAZ-CBG	-3.56	112.33	120.51
3	C	600	X22	CBC-CBH-NAW	-3.34	104.95	111.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	X22	2	0
3	B	600	X22	2	0
3	C	600	X22	2	0
3	D	600	X22	2	0
3	D	700	X22	2	0
3	D	701	X22	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	101/130 (77%)	0.01	3 (2%) 54 29	23, 41, 70, 72	0
1	B	101/130 (77%)	0.25	3 (2%) 54 29	28, 48, 76, 78	0
1	C	102/130 (78%)	-0.21	3 (2%) 55 31	18, 35, 54, 66	0
1	D	100/130 (76%)	-0.26	1 (1%) 84 69	23, 38, 53, 60	0
All	All	404/520 (77%)	-0.05	10 (2%) 61 37	18, 40, 71, 78	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	355	THR	4.9
1	A	274	THR	3.7
1	C	254	THR	3.6
1	A	270	PHE	3.2
1	C	253	SER	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	X22	D	700	36/36	0.90	0.26	1.95	24,35,38,40	0
3	X22	B	600	36/36	0.91	0.30	1.62	52,54,58,58	0
3	X22	D	600	36/36	0.94	0.21	1.27	19,22,27,29	0
3	X22	A	600	36/36	0.94	0.25	0.95	27,28,30,31	0
3	X22	D	701	36/36	0.94	0.21	0.77	19,23,25,26	0
3	X22	C	600	36/36	0.95	0.19	-0.01	23,25,30,31	0
2	ZN	B	5	1/1	0.99	0.08	-1.77	35,35,35,35	0
2	ZN	C	4	1/1	0.99	0.10	-1.78	28,28,28,28	0
2	ZN	A	6	1/1	0.98	0.06	-1.87	51,51,51,51	0
2	ZN	D	3	1/1	0.99	0.07	-	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.