



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:36 PM GMT

PDB ID : 4CQN
Title : Crystal structure of the E.coli LeuRS-tRNA complex with the non- cognate
isoleucyl adenylate analogue
Authors : Palencia, A.; Cusack, S.; Cvetesic, N.; Haslaz, I.; Gruic-Sovulj, I.
Deposited on : 2014-02-20
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)
RNA backbone	2183	1172 (3.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	880	<div> <div></div> <div> <div></div> <div>58%</div> <div>36%</div> <div>..</div> </div> </div>
1	D	880	<div> <div>16%</div> <div>59%</div> <div>34%</div> <div>..</div> </div>
2	B	82	<div> <div></div> <div> <div></div> <div>45%</div> <div>38%</div> <div>17%</div> </div> </div>
2	E	82	<div> <div>11%</div> <div>32%</div> <div>51%</div> <div>13%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ILA	A	1863	X	-	X	-
4	ILA	D	1863	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 17558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LEUCINE-TRNA LIGASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	860	Total	C	N	O	S	0	2	0
			6844	4345	1161	1293	45			
1	D	860	Total	C	N	O	S	0	0	0
			6834	4339	1158	1292	45			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP P07813
A	-18	GLY	-	EXPRESSION TAG	UNP P07813
A	-17	SER	-	EXPRESSION TAG	UNP P07813
A	-16	SER	-	EXPRESSION TAG	UNP P07813
A	-15	HIS	-	EXPRESSION TAG	UNP P07813
A	-14	HIS	-	EXPRESSION TAG	UNP P07813
A	-13	HIS	-	EXPRESSION TAG	UNP P07813
A	-12	HIS	-	EXPRESSION TAG	UNP P07813
A	-11	HIS	-	EXPRESSION TAG	UNP P07813
A	-10	HIS	-	EXPRESSION TAG	UNP P07813
A	-9	SER	-	EXPRESSION TAG	UNP P07813
A	-8	SER	-	EXPRESSION TAG	UNP P07813
A	-7	GLY	-	EXPRESSION TAG	UNP P07813
A	-6	LEU	-	EXPRESSION TAG	UNP P07813
A	-5	VAL	-	EXPRESSION TAG	UNP P07813
A	-4	PRO	-	EXPRESSION TAG	UNP P07813
A	-3	ARG	-	EXPRESSION TAG	UNP P07813
A	-2	GLY	-	EXPRESSION TAG	UNP P07813
A	-1	SER	-	EXPRESSION TAG	UNP P07813
A	0	HIS	-	EXPRESSION TAG	UNP P07813
D	-19	MET	-	EXPRESSION TAG	UNP P07813
D	-18	GLY	-	EXPRESSION TAG	UNP P07813
D	-17	SER	-	EXPRESSION TAG	UNP P07813
D	-16	SER	-	EXPRESSION TAG	UNP P07813
D	-15	HIS	-	EXPRESSION TAG	UNP P07813

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-14	HIS	-	EXPRESSION TAG	UNP P07813
D	-13	HIS	-	EXPRESSION TAG	UNP P07813
D	-12	HIS	-	EXPRESSION TAG	UNP P07813
D	-11	HIS	-	EXPRESSION TAG	UNP P07813
D	-10	HIS	-	EXPRESSION TAG	UNP P07813
D	-9	SER	-	EXPRESSION TAG	UNP P07813
D	-8	SER	-	EXPRESSION TAG	UNP P07813
D	-7	GLY	-	EXPRESSION TAG	UNP P07813
D	-6	LEU	-	EXPRESSION TAG	UNP P07813
D	-5	VAL	-	EXPRESSION TAG	UNP P07813
D	-4	PRO	-	EXPRESSION TAG	UNP P07813
D	-3	ARG	-	EXPRESSION TAG	UNP P07813
D	-2	GLY	-	EXPRESSION TAG	UNP P07813
D	-1	SER	-	EXPRESSION TAG	UNP P07813
D	0	HIS	-	EXPRESSION TAG	UNP P07813


- Molecule 2 is a RNA chain called ESCHERICHIA COLI TRNA-LEU UAA ISOACCEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	82	Total	C	N	O	P	0	0	0
			1755	781	317	575	82			
2	E	79	Total	C	N	O	P	0	0	0
			1691	752	305	555	79			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is N-[ISOLEUCINYL]-N'-[ADENOSYL]-DIAMINOSUFONE (three-letter code: ILA) (formula: C₁₆H₂₆N₈O₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 16	N 8	O 6	S 1	0	0
4	A	1	Total 31	C 16	N 8	O 6	S 1	0	0
4	D	1	Total 31	C 16	N 8	O 6	S 1	0	0
4	D	1	Total 31	C 16	N 8	O 6	S 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	E	1	Total Mg 1 1	0	0


- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	233	Total O 233 233	0	0
6	B	57	Total O 57 57	0	0
6	D	16	Total O 16 16	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LEUCINE-TRNA LIGASE





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	155.62Å 67.58Å 224.59Å 90.00° 105.00° 90.00°	Depositor
Resolution (Å)	46.49 – 2.50 46.45 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.1 (46.49-2.50) 97.1 (46.45-2.50)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.214 , 0.290 0.214 , 0.290	Depositor DCC
R_{free} test set	3828 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.592	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 45.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 76243 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	17558	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, ILA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.71	0/7014	0.88	7/9520 (0.1%)
1	D	0.38	0/6995	0.60	1/9495 (0.0%)
2	B	0.61	2/1959 (0.1%)	0.90	4/3047 (0.1%)
2	E	0.37	1/1887 (0.1%)	0.64	0/2935
All	All	0.56	3/17855 (0.0%)	0.76	12/24997 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	G	OP3-P	-9.82	1.49	1.61
2	B	1	G	OP3-P	-9.46	1.49	1.61
2	B	12	G	O3'-P	-6.47	1.53	1.61

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	58	A	O5'-P-OP1	-7.04	99.36	105.70
1	A	345	ASP	CB-CG-OD1	6.49	124.14	118.30
2	B	72	U	O5'-P-OP1	6.16	118.09	110.70
1	A	345	ASP	CB-CG-OD2	-5.78	113.10	118.30
1	A	791	ASP	CB-CG-OD1	5.62	123.36	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	592	ILE	Peptide
1	A	80	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6844	0	6696	258	0
1	D	6834	0	6683	216	0
2	B	1755	0	890	37	0
2	E	1691	0	858	38	0
3	A	1	0	0	0	0
3	D	1	0	0	0	0
4	A	62	0	52	12	0
4	D	62	0	52	7	0
5	B	1	0	0	0	0
5	E	1	0	0	0	0
6	A	233	0	0	15	0
6	B	57	0	0	1	0
6	D	16	0	0	2	0
All	All	17558	0	15231	537	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 537 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:422:VAL:HG11	1:A:494:MET:HE1	1.35	1.07
1:A:267:LEU:HD21	1:A:305:THR:HG21	1.40	1.00
1:A:553:MET:CE	6:A:2052:HOH:O	2.10	0.99
1:A:342:ASP:HB2	4:A:1863:ILA:CG2	1.96	0.95
2:B:41:C:H5''	2:B:41:C:H6	1.32	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	860/880 (98%)	797 (93%)	54 (6%)	9 (1%)	19	34
1	D	858/880 (98%)	716 (83%)	119 (14%)	23 (3%)	6	9
All	All	1718/1760 (98%)	1513 (88%)	173 (10%)	32 (2%)	10	16

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	439	ASP
1	A	693	ALA
1	D	237	ASN
1	D	463	ILE
1	A	161	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	726/741 (98%)	661 (91%)	65 (9%)	12	22
1	D	723/741 (98%)	655 (91%)	68 (9%)	11	20
All	All	1449/1482 (98%)	1316 (91%)	133 (9%)	11	21

5 of 133 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	798	ASP
1	D	125	LEU
1	D	702	ARG
1	A	816	VAL
1	D	21	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	190	GLN
1	D	222	ASN
1	D	533	HIS
1	D	157	ASN
1	D	507	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	81/82 (98%)	15 (18%)	6 (7%)
2	E	76/82 (92%)	21 (27%)	3 (3%)
All	All	157/164 (95%)	36 (22%)	9 (5%)

5 of 36 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	9	G
2	B	11	U
2	B	15	A
2	B	16	U
2	B	17	C

5 of 9 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	47(D)	C
2	E	48	U
2	E	19	G
2	B	19	G
2	B	47(E)	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ILA	A	1862	-	27,33,33	2.84	5 (18%)	31,49,49	3.36	10 (32%)
4	ILA	A	1863	-	27,33,33	2.57	5 (18%)	31,49,49	3.43	14 (45%)
4	ILA	D	1862	-	27,33,33	2.40	5 (18%)	31,49,49	2.36	5 (16%)
4	ILA	D	1863	-	27,33,33	2.41	6 (22%)	31,49,49	2.43	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ILA	A	1862	-	-	0/19/41/41	0/3/3/3
4	ILA	A	1863	-	1/1/8/10	1/19/41/41	0/3/3/3
4	ILA	D	1862	-	-	0/19/41/41	0/3/3/3
4	ILA	D	1863	-	1/1/8/10	1/19/41/41	0/3/3/3

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1862	ILA	SA-N5'	-10.39	1.51	1.61
4	D	1862	ILA	SA-N5'	-8.69	1.52	1.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1863	ILA	SA-N5'	-8.25	1.53	1.61
4	A	1863	ILA	SA-N5'	-8.21	1.53	1.61
4	A	1862	ILA	SA-N3A	-5.57	1.52	1.63

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1863	ILA	N3-C2-N1	-14.16	118.05	128.89
4	A	1862	ILA	N3-C2-N1	-10.55	120.82	128.89
4	D	1863	ILA	N3-C2-N1	-10.22	121.07	128.89
4	A	1862	ILA	O2A-SA-O1A	-9.90	103.88	120.04
4	D	1862	ILA	N3-C2-N1	-8.86	122.11	128.89

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	D	1863	ILA	CB
4	A	1863	ILA	CB

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	1863	ILA	N3A-SA-N5'-C5'
4	A	1863	ILA	N3A-SA-N5'-C5'

There are no ring outliers.

4 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1862	ILA	1	0
4	A	1863	ILA	11	0
4	D	1862	ILA	3	0
4	D	1863	ILA	4	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	860/880 (97%)	-0.24	6 (0%) 89 90	6, 19, 48, 99	0
1	D	860/880 (97%)	1.04	141 (16%) 2 2	36, 71, 111, 142	0
2	B	82/82 (100%)	-0.43	1 (1%) 81 83	9, 21, 72, 116	0
2	E	79/82 (96%)	0.79	9 (11%) 7 6	56, 79, 125, 142	0
All	All	1881/1924 (97%)	0.38	157 (8%) 14 15	6, 47, 105, 142	0

The worst 5 of 157 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	860	GLY	13.2
1	D	809	LYS	6.1
1	D	330	TYR	5.9
1	D	322	ALA	5.9
1	D	321	TRP	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ILA	A	1863	31/31	0.95	0.14	1.00	21,28,32,36	0
4	ILA	D	1863	31/31	0.65	0.31	0.64	104,151,160,160	0
4	ILA	A	1862	31/31	0.97	0.15	0.38	7,10,11,11	0
4	ILA	D	1862	31/31	0.95	0.16	-0.68	38,48,56,61	0
3	ZN	D	1861	1/1	0.86	0.06	-1.45	100,100,100,100	0
3	ZN	A	1861	1/1	0.96	0.05	-3.07	65,65,65,65	0
5	MG	B	1077	1/1	0.93	0.09	-	25,25,25,25	0
5	MG	E	1077	1/1	0.93	0.22	-	52,52,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.