



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:30 PM GMT

PDB ID : 5DBG
Title : Crystal Structure of Iridoid Synthase from Cantharanthus roseus in complex with NAD+
Authors : Liu, W.D.; Hu, Y.M.; Zheng, Y.Y.; Xu, Z.X.; Ko, T.P.; Chen, C.C.; Guo, R.T.
Deposited on : 2015-08-21
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	 3% 85% 10% ..
1	B	374	 3% 84% 10% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iridoid synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	0	0
			2858	1841	471	527	19			
1	B	361	Total	C	N	O	S	0	0	0
			2865	1846	472	528	19			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	ALA	-	expression tag	UNP K7WDL7
A	390	SER	-	expression tag	UNP K7WDL7
A	391	ALA	-	expression tag	UNP K7WDL7
A	392	TRP	-	expression tag	UNP K7WDL7
A	393	SER	-	expression tag	UNP K7WDL7
A	394	HIS	-	expression tag	UNP K7WDL7
A	395	PRO	-	expression tag	UNP K7WDL7
A	396	GLN	-	expression tag	UNP K7WDL7
A	397	PHE	-	expression tag	UNP K7WDL7
A	398	GLU	-	expression tag	UNP K7WDL7
A	399	LYS	-	expression tag	UNP K7WDL7
B	389	ALA	-	expression tag	UNP K7WDL7
B	390	SER	-	expression tag	UNP K7WDL7
B	391	ALA	-	expression tag	UNP K7WDL7
B	392	TRP	-	expression tag	UNP K7WDL7
B	393	SER	-	expression tag	UNP K7WDL7
B	394	HIS	-	expression tag	UNP K7WDL7
B	395	PRO	-	expression tag	UNP K7WDL7
B	396	GLN	-	expression tag	UNP K7WDL7
B	397	PHE	-	expression tag	UNP K7WDL7
B	398	GLU	-	expression tag	UNP K7WDL7
B	399	LYS	-	expression tag	UNP K7WDL7

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	262	Total	O	0	0
			262	262		
3	B	299	Total	O	0	0
			299	299		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Iridoid synthase



• Molecule 1: Iridoid synthase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	92.12Å 95.58Å 173.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.95 24.71 – 1.94	Depositor EDS
% Data completeness (in resolution range)	95.6 (25.00-1.95) 95.6 (24.71-1.94)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.177 , 0.230 0.186 , 0.238	Depositor DCC
R_{free} test set	2694 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.429	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.5	EDS
Estimated twinning fraction	0.058 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 54097 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6372	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	1/2932 (0.0%)	0.91	5/3980 (0.1%)
1	B	0.88	0/2939	0.91	7/3990 (0.2%)
All	All	0.91	1/5871 (0.0%)	0.91	12/7970 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	336	GLU	CD-OE1	7.99	1.34	1.25

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	249	VAL	CB-CA-C	-7.23	97.66	111.40
1	A	249	VAL	CB-CA-C	-6.67	98.73	111.40
1	A	249	VAL	CG1-CB-CG2	6.61	121.48	110.90
1	B	200	ARG	NE-CZ-NH2	-5.99	117.31	120.30
1	B	249	VAL	CG1-CB-CG2	5.61	119.88	110.90
1	A	168	ASP	CB-CG-OD1	5.52	123.27	118.30
1	B	87	ASP	CB-CG-OD1	5.47	123.22	118.30
1	B	192	ASN	C-N-CA	-5.32	108.40	121.70
1	B	172	LEU	CA-CB-CG	5.31	127.52	115.30
1	B	289	LEU	CB-CG-CD2	5.17	119.79	111.00
1	A	325	LYS	CD-CE-NZ	5.10	123.43	111.70
1	A	151	ILE	CG1-CB-CG2	5.01	122.43	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2858	0	2825	21	0
1	B	2865	0	2834	28	0
2	A	44	0	26	0	0
2	B	44	0	26	0	0
3	A	262	0	0	6	2
3	B	299	0	0	6	1
All	All	6372	0	5711	49	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:315:ASP:HB3	3:B:504:HOH:O	1.37	1.23
1:B:140:CYS:HB3	3:B:750:HOH:O	0.81	0.98
1:B:151:ILE:HD11	1:B:176:ASN:HA	1.57	0.85
1:B:328:ASN:O	1:B:385:ARG:NH1	2.09	0.83
1:A:151:ILE:HD11	3:A:529:HOH:O	1.80	0.79
1:A:256:GLU:OE2	3:A:501:HOH:O	2.00	0.78
1:A:26:LYS:HD3	3:A:694:HOH:O	1.85	0.77
1:B:222:TYR:HA	1:B:313:MET:HE1	1.70	0.73
1:A:110:ILE:HD12	1:A:112:SER:H	1.55	0.70
1:B:249:VAL:HG13	1:B:275:ASN:HA	1.75	0.66
1:B:72:TRP:HZ2	1:B:210:CYS:HG	1.47	0.62
1:B:214:ASN:C	1:B:214:ASN:HD22	2.03	0.61
1:A:214:ASN:C	1:A:214:ASN:HD22	2.06	0.59
1:A:249:VAL:HG13	1:A:275:ASN:HA	1.84	0.59
1:B:222:TYR:HA	1:B:313:MET:CE	2.35	0.57
1:B:222:TYR:HD1	1:B:313:MET:CE	2.18	0.57
1:B:151:ILE:HD11	1:B:176:ASN:CA	2.33	0.55
1:B:95:LEU:HA	1:B:98:LEU:HD22	1.89	0.55
1:B:88:ASN:HB2	3:B:672:HOH:O	2.06	0.55
1:A:151:ILE:CD1	3:A:529:HOH:O	2.46	0.54
1:B:265:PRO:O	1:B:268:LYS:HD2	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:140:CYS:CB	3:B:750:HOH:O	1.70	0.52
1:A:382:ARG:HG3	1:A:387:ILE:HD11	1.93	0.51
1:B:132:ASN:ND2	3:B:507:HOH:O	2.43	0.51
1:A:72:TRP:HZ2	1:A:210:CYS:HG	1.59	0.51
1:B:173:ASN:HD22	1:B:173:ASN:C	2.16	0.49
1:A:81:ILE:HD12	3:A:660:HOH:O	2.13	0.48
1:B:382:ARG:HG3	1:B:387:ILE:HD11	1.95	0.47
1:B:110:ILE:HD12	1:B:121:MET:SD	2.54	0.47
1:A:163:SER:HB2	1:A:164:PRO:HA	1.96	0.47
1:A:80:TYR:C	1:A:81:ILE:HD13	2.35	0.46
1:B:249:VAL:HG13	1:B:275:ASN:CA	2.43	0.45
1:B:214:ASN:ND2	1:B:217:SER:H	2.14	0.44
1:B:222:TYR:HD1	1:B:313:MET:HE1	1.83	0.44
1:A:222:TYR:CD1	1:A:309:LEU:HD11	2.53	0.44
1:B:72:TRP:HZ2	1:B:210:CYS:SG	2.40	0.44
1:B:385:ARG:NH1	3:B:502:HOH:O	2.31	0.43
1:A:151:ILE:HD13	1:A:151:ILE:HA	1.85	0.43
1:A:237:GLY:HA3	1:A:347:PHE:HB3	1.99	0.43
1:B:306:GLN:OE1	1:B:348:CYS:HB3	2.19	0.43
1:B:237:GLY:HA3	1:B:347:PHE:HB3	1.99	0.43
1:A:306:GLN:OE1	1:A:348:CYS:HB3	2.19	0.42
1:A:137:GLN:HA	3:A:693:HOH:O	2.18	0.42
1:B:380:LYS:HE3	1:B:384:TYR:CZ	2.55	0.42
1:A:214:ASN:ND2	1:A:217:SER:H	2.19	0.41
1:A:216:VAL:HG12	1:A:381:MET:HE1	2.02	0.41
1:A:220:CYS:SG	1:A:378:ILE:HG12	2.60	0.41
1:A:110:ILE:HG23	1:A:121:MET:SD	2.61	0.40
1:B:220:CYS:SG	1:B:378:ILE:HG12	2.62	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:535:HOH:O	3:A:698:HOH:O[3_455]	2.03	0.17
3:A:722:HOH:O	3:B:509:HOH:O[5_455]	2.08	0.12

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	356/374 (95%)	350 (98%)	6 (2%)	0	100	100
1	B	357/374 (96%)	350 (98%)	7 (2%)	0	100	100
All	All	713/748 (95%)	700 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/331 (96%)	305 (96%)	14 (4%)	35	19
1	B	320/331 (97%)	302 (94%)	18 (6%)	26	11
All	All	639/662 (96%)	607 (95%)	32 (5%)	30	14

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	SER
1	A	44	SER
1	A	81	ILE
1	A	88	ASN
1	A	110	ILE
1	A	135	ASN
1	A	151	ILE
1	A	214	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	234	VAL
1	A	249	VAL
1	A	268	LYS
1	A	289	LEU
1	A	329	LEU
1	A	334	LEU
1	B	26	LYS
1	B	27	SER
1	B	93	SER
1	B	98	LEU
1	B	110	ILE
1	B	151	ILE
1	B	172	LEU
1	B	173	ASN
1	B	185	LEU
1	B	214	ASN
1	B	249	VAL
1	B	268	LYS
1	B	289	LEU
1	B	297	MET
1	B	315	ASP
1	B	329	LEU
1	B	334	LEU
1	B	362	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	88	ASN
1	A	142	GLN
1	A	192	ASN
1	A	214	ASN
1	A	284	HIS
1	B	82	GLN
1	B	132	ASN
1	B	137	GLN
1	B	173	ASN
1	B	192	ASN
1	B	214	ASN
1	B	284	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	A	400	-	42,48,48	1.17	4 (9%)	46,73,73	1.61	8 (17%)
2	NAD	B	401	-	42,48,48	1.16	6 (14%)	46,73,73	1.74	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	400	-	-	0/22/62/62	0/5/5/5
2	NAD	B	401	-	-	0/22/62/62	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAD	C2N-N1N	-2.94	1.30	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	NAD	O2D-C2D	-2.28	1.37	1.43
2	B	401	NAD	C2B-C1B	-2.28	1.50	1.53
2	A	400	NAD	C4A-N3A	-2.21	1.32	1.35
2	B	401	NAD	C5A-N7A	-2.19	1.31	1.39
2	B	401	NAD	C5N-C4N	-2.02	1.35	1.38
2	B	401	NAD	C5A-C4A	2.16	1.45	1.40
2	B	401	NAD	C2A-N3A	2.58	1.36	1.32
2	A	400	NAD	C5A-C4A	2.81	1.46	1.40
2	A	400	NAD	C4N-C3N	2.87	1.44	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	NAD	N3A-C2A-N1A	-6.04	124.13	128.87
2	B	401	NAD	N3A-C2A-N1A	-5.59	124.48	128.87
2	A	400	NAD	C4D-O4D-C1D	-4.29	105.10	109.64
2	B	401	NAD	C1B-N9A-C4A	-3.84	122.52	126.81
2	A	400	NAD	C1B-N9A-C4A	-3.33	123.09	126.81
2	B	401	NAD	C4B-O4B-C1B	-3.21	106.24	109.64
2	B	401	NAD	C3N-C7N-N7N	-2.66	114.82	117.82
2	A	400	NAD	C2B-C1B-N9A	-2.42	106.98	113.47
2	A	400	NAD	O5B-C5B-C4B	-2.26	100.94	109.09
2	A	400	NAD	O2N-PN-O1N	2.18	123.90	112.56
2	A	400	NAD	O2A-PA-O1A	2.28	124.43	112.56
2	A	400	NAD	O4B-C1B-N9A	2.31	112.46	108.11
2	B	401	NAD	O7N-C7N-C3N	3.04	122.97	119.60
2	B	401	NAD	N6A-C6A-N1A	3.48	124.35	118.52
2	B	401	NAD	O4D-C1D-N1N	4.55	113.02	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/374 (96%)	0.17	12 (3%) 50 61	15, 25, 46, 83	0
1	B	361/374 (96%)	0.09	11 (3%) 54 64	15, 24, 41, 71	0
All	All	721/748 (96%)	0.13	23 (3%) 51 61	15, 25, 44, 83	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	26	LYS	4.4
1	B	158	VAL	3.7
1	A	159	VAL	3.5
1	A	140	CYS	3.4
1	A	305	GLU	3.2
1	B	26	LYS	3.1
1	A	151	ILE	2.8
1	B	140	CYS	2.8
1	B	173	ASN	2.7
1	A	318	GLN	2.6
1	B	107	VAL	2.5
1	B	27	SER	2.5
1	A	304	LYS	2.4
1	A	27	SER	2.3
1	B	221	VAL	2.3
1	A	35	VAL	2.3
1	B	35	VAL	2.3
1	A	139	VAL	2.2
1	B	104	ILE	2.2
1	A	221	VAL	2.2
1	B	141	LEU	2.1
1	A	141	LEU	2.1
1	B	219	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAD	B	401	44/44	0.97	0.09	-0.79	16,21,23,25	0
2	NAD	A	400	44/44	0.97	0.08	-0.94	14,20,22,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.