



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:41 AM GMT

PDB ID : 2DWN
Title : Crystal structure of the PriA protein complexed with oligonucleotides
Authors : Sasaki, K.; Ose, T.; Tanaka, T.; Masai, H.; Maenaka, K.; Kohda, D.
Deposited on : 2006-08-15
Resolution : 3.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1005 (3.42-3.30)
Clashscore	102246	1076 (3.42-3.30)
Ramachandran outliers	100387	1059 (3.42-3.30)
Sidechain outliers	100360	1058 (3.42-3.30)
RSRZ outliers	91569	1010 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	105	<div> <div>8%</div> <div>41%</div> <div>53%</div> <div>6%</div> </div>
1	B	105	<div> <div>26%</div> <div>42%</div> <div>50%</div> <div>8%</div> </div>
1	C	105	<div> <div>8%</div> <div>48%</div> <div>43%</div> <div>7%</div> </div>
1	D	105	<div> <div>10%</div> <div>45%</div> <div>44%</div> <div>8%</div> </div>
2	E	2	<div> <div>50%</div> <div>50%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3334 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Primosomal protein N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	105	Total	C	N	O	S	0	0	0
			832	546	141	142	3			
1	B	104	Total	C	N	O	S	0	0	0
			820	540	137	140	3			
1	C	105	Total	C	N	O	S	0	0	0
			832	546	141	142	3			
1	D	103	Total	C	N	O	S	0	0	0
			812	535	136	139	2			


- Molecule 2 is a DNA chain called 5'-D(*AP*(DGP))-3'.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	1	Total	C	N	O	0	0	0
			19	10	5	4			
2	F	1	Total	C	N	O	0	0	0
			19	10	5	4			


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Primosomal protein N




• Molecule 1: Primosomal protein N



• Molecule 1: Primosomal protein N



• Molecule 1: Primosomal protein N





- Molecule 2: 5'-D(*AP*(DGP))-3'



- Molecule 2: 5'-D(*AP*(DGP))-3'



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	112.25Å 112.25Å 260.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 3.35 47.78 – 3.35	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-3.35) 99.9 (47.78-3.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.63 (at 3.33Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.281 , 0.333 0.270 , 0.323	Depositor DCC
R_{free} test set	646 reflections (7.34%)	DCC
Wilson B-factor (Å ²)	116.0	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 114.8	EDS
Estimated twinning fraction	0.024 for -h,1/3*h-1/3*k-1/3*l,-4/3*h-8/3*k+1/3*l 0.016 for -1/3*h+1/3*k+1/3*l,-k,8/3*h+4/3*k+1/3*l 0.029 for -2/3*h-1/3*k-1/3*l,-1/3*h-2/3*k+1/3*l,-4/3*h+4/3*k+1/3*l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 9392 reflections	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3334	wwPDB-VP
Average B, all atoms (Å ²)	108.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DGP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/857	0.82	0/1173
1	B	0.44	0/845	0.71	0/1159
1	C	0.49	0/857	0.77	0/1173
1	D	0.56	0/837	0.86	0/1148
All	All	0.52	0/3396	0.79	0/4653

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	832	0	853	85	0
1	B	820	0	840	98	0
1	C	832	0	853	77	0
1	D	812	0	829	66	0
2	E	19	0	12	0	0
2	F	19	0	12	0	0
All	All	3334	0	3399	275	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 41.

The worst 5 of 275 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:LEU:HB3	1:B:13:PRO:HD3	1.27	1.11
1:B:1:MET:HB3	1:B:2:PRO:HD2	1.32	1.08
1:A:31:ARG:HD3	1:A:68:ASP:OD2	1.56	1.05
1:C:12:LEU:HB3	1:C:13:PRO:HD3	1.43	0.98
1:C:12:LEU:HD21	1:C:86:ASP:HB2	1.47	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/105 (98%)	92 (89%)	7 (7%)	4 (4%)	4	29
1	B	102/105 (97%)	80 (78%)	17 (17%)	5 (5%)	3	22
1	C	103/105 (98%)	82 (80%)	15 (15%)	6 (6%)	2	17
1	D	101/105 (96%)	81 (80%)	12 (12%)	8 (8%)	1	9
All	All	409/420 (97%)	335 (82%)	51 (12%)	23 (6%)	2	18

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	PRO
1	A	12	LEU
1	B	11	PRO
1	B	39	GLN
1	C	2	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/92 (100%)	89 (97%)	3 (3%)	45	79
1	B	91/92 (99%)	87 (96%)	4 (4%)	35	72
1	C	92/92 (100%)	83 (90%)	9 (10%)	10	37
1	D	90/92 (98%)	83 (92%)	7 (8%)	16	51
All	All	365/368 (99%)	342 (94%)	23 (6%)	22	61

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	22	GLU
1	C	60	LEU
1	D	58	ASN
1	C	54	GLU
1	C	66	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	58	ASN
1	C	89	HIS
1	D	58	ASN
1	B	58	ASN
1	C	98	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	DGP	E	2	-	17,21,25	1.20	1 (5%)	20,31,38	2.68	4 (20%)
2	DGP	F	2	-	17,21,25	1.10	2 (11%)	20,31,38	2.67	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DGP	E	2	-	-	0/2/18/22	0/3/3/3
2	DGP	F	2	-	-	0/2/18/22	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2	DGP	C8-N7	-2.15	1.30	1.34
2	F	2	DGP	C6-N1	3.52	1.39	1.33
2	E	2	DGP	C6-N1	3.75	1.40	1.33

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	DGP	C5-C6-N1	-8.88	111.44	123.59
2	F	2	DGP	C5-C6-N1	-8.88	111.45	123.59
2	E	2	DGP	N3-C2-N1	-2.42	123.76	127.44
2	F	2	DGP	N3-C2-N1	-2.35	123.87	127.44
2	E	2	DGP	O5'-C5'-C4'	2.67	120.16	111.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	105/105 (100%)	0.68	8 (7%) 17 17	40, 89, 176, 200	0
1	B	104/105 (99%)	1.38	27 (25%) 1 1	47, 141, 200, 200	0
1	C	105/105 (100%)	0.48	8 (7%) 17 17	31, 77, 177, 200	0
1	D	103/105 (98%)	0.86	10 (9%) 10 11	29, 87, 187, 200	0
2	E	0/2	-	-	-	-
2	F	0/2	-	-	-	-
All	All	417/424 (98%)	0.85	53 (12%) 5 4	29, 95, 199, 200	0

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	53	SER	12.6
1	D	12	LEU	11.4
1	A	12	LEU	11.4
1	A	13	PRO	11.0
1	D	13	PRO	8.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	DGP	F	2	19/23	0.72	0.36	-	85,147,147,147	0
2	DGP	E	2	19/23	0.85	0.41	-	100,173,173,173	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.