



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:58 PM GMT

PDB ID : 4DYN  
Title : Crystal Structure of WSN/A Influenza Nucleoprotein with BMS-885838 Lig-  
and Bound  
Authors : Lewis, H.A.; Baldwin, E.T.; Steinbacher, S.; Maskos, K.; Mortl, M.; Kiefer-  
sauer, R.; Edavettal, S.; McDonnell, P.A.; Pearce, B.C.; Langley, D.R.  
Deposited on : 2012-02-29  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	499	 2% 80% 7% 12%
1	B	499	 5% 81% 7% 12%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nucleocapsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	1	0
			3284	2060	576	623	25			
1	B	437	Total	C	N	O	S	0	1	0
			3162	1990	555	592	25			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	499	LEU	-	EXPRESSION TAG	UNP B4URF1
A	500	GLU	-	EXPRESSION TAG	UNP B4URF1
A	501	HIS	-	EXPRESSION TAG	UNP B4URF1
A	502	HIS	-	EXPRESSION TAG	UNP B4URF1
A	503	HIS	-	EXPRESSION TAG	UNP B4URF1
A	504	HIS	-	EXPRESSION TAG	UNP B4URF1
A	505	HIS	-	EXPRESSION TAG	UNP B4URF1
A	506	HIS	-	EXPRESSION TAG	UNP B4URF1
B	499	LEU	-	EXPRESSION TAG	UNP B4URF1
B	500	GLU	-	EXPRESSION TAG	UNP B4URF1
B	501	HIS	-	EXPRESSION TAG	UNP B4URF1
B	502	HIS	-	EXPRESSION TAG	UNP B4URF1
B	503	HIS	-	EXPRESSION TAG	UNP B4URF1
B	504	HIS	-	EXPRESSION TAG	UNP B4URF1
B	505	HIS	-	EXPRESSION TAG	UNP B4URF1
B	506	HIS	-	EXPRESSION TAG	UNP B4URF1

- Molecule 2 is N-[4-CHLORANYL-5-[4-[[3-(2-METHOXYPHENYL)-5-METHYL-1,2-OXAZOL-4-YL]CARBONYL]PIPERAZIN-1-YL]-2-NITRO-PHENYL]PYRIDINE-2-CARBOXYMIDE (three-letter code: OMR) (formula: C<sub>28</sub>H<sub>25</sub>ClN<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 41	C 28	Cl 1	N 6	O 6	0	0
2	B	1	Total 41	C 28	Cl 1	N 6	O 6	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	64	Total O 64 64	0	0
3	B	31	Total O 31 31	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Nucleocapsid protein



#### • Molecule 1: Nucleocapsid protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.61Å 145.61Å 145.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.90 – 2.40 48.54 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.90-2.40) 99.9 (48.54-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.49 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.242 , 0.291 0.239 , 0.286	Depositor DCC
$R_{free}$ test set	981 reflections (2.49%)	DCC
Wilson B-factor (Å <sup>2</sup> )	43.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.3	EDS
Estimated twinning fraction	0.029 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 40358 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6623	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 0MR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/3340	0.46	0/4513
1	B	0.31	0/3217	0.45	0/4359
All	All	0.31	0/6557	0.45	0/8872

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3284	0	3117	19	0
1	B	3162	0	2892	18	0
2	A	41	0	25	5	0
2	B	41	0	25	5	0
3	A	64	0	0	0	0
3	B	31	0	0	1	0
All	All	6623	0	6059	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:311:GLN:HE22	1:B:309:ASN:HA	1.29	0.96
1:A:311:GLN:NE2	1:B:309:ASN:HA	2.01	0.75
1:A:184:LYS:HD3	1:A:189:MET:HE1	1.69	0.74
1:A:189:MET:HE2	1:A:189:MET:HA	1.83	0.60
1:A:184:LYS:HD3	1:A:189:MET:CE	2.34	0.57
1:B:309:ASN:OD1	2:B:601:0MR:H16	2.06	0.54
2:B:601:0MR:O37	2:B:601:0MR:H8	2.09	0.52
2:B:601:0MR:C18	2:B:601:0MR:H18	2.39	0.52
2:A:601:0MR:C18	2:A:601:0MR:H18	2.41	0.50
1:B:422:ARG:N	1:B:423:PRO:HD2	2.27	0.50
1:A:154:LEU:HD13	1:A:161:PRO:HA	1.92	0.49
1:A:408:ILE:HG12	1:A:419:PRO:HD2	1.95	0.48
1:A:309:ASN:ND2	2:A:601:0MR:H16	2.28	0.48
1:A:273:LYS:HD3	1:A:298:LEU:HD23	1.96	0.47
1:B:324:HIS:HB3	1:B:358:LEU:HG	1.96	0.47
1:B:39:PHE:CZ	1:B:67:VAL:HB	2.49	0.46
1:A:302:ASP:HB2	1:A:303:PRO:HD3	1.95	0.46
1:A:221:ARG:O	1:A:225:ILE:HG12	2.16	0.46
1:A:141:SER:OG	1:A:169:GLY:HA2	2.15	0.46
1:B:36:ILE:HD13	1:B:120:TRP:HH2	1.81	0.45
2:A:601:0MR:C11	2:A:601:0MR:H18	2.47	0.44
2:B:601:0MR:H18	2:B:601:0MR:C11	2.48	0.44
2:A:601:0MR:H21	1:B:313:TYR:CG	2.53	0.44
1:B:334:HIS:HE1	3:B:708:HOH:O	2.01	0.44
1:A:51:ASP:HB3	1:B:287:SER:HA	1.99	0.44
1:A:313:TYR:CG	2:B:601:0MR:H21	2.53	0.43
1:A:424:THR:OG1	1:A:452:ARG:HD2	2.18	0.43
1:B:133:LEU:O	1:B:137:MET:HG3	2.19	0.43
1:A:287:SER:O	1:B:50[A]:SER:OG	2.37	0.42
2:A:601:0MR:O37	2:A:601:0MR:H8	2.19	0.42
1:B:160:ASP:HB3	1:B:163:MET:HG3	2.01	0.42
1:B:302:ASP:HB2	1:B:303:PRO:HD3	2.01	0.42
1:A:164:CYS:HA	1:A:167:MET:HG2	2.02	0.42
1:B:321:ASN:HD22	1:B:324:HIS:CE1	2.39	0.41
1:A:308:GLN:HA	1:A:382:ARG:HG2	2.02	0.41
1:B:321:ASN:HA	1:B:322:PRO:HD2	1.96	0.41
1:B:190:VAL:HG13	1:B:256:LEU:HB3	2.03	0.41
1:A:89:PRO:HB2	1:A:90:LYS:H	1.62	0.41
1:B:470:LYS:O	1:B:472:THR:N	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	428/499 (86%)	419 (98%)	9 (2%)	0	100	100
1	B	426/499 (85%)	417 (98%)	7 (2%)	2 (0%)	34	48
All	All	854/998 (86%)	836 (98%)	16 (2%)	2 (0%)	52	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	471	ALA
1	B	17	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/425 (77%)	316 (97%)	10 (3%)	47	69
1	B	292/425 (69%)	286 (98%)	6 (2%)	61	80
All	All	618/850 (73%)	602 (97%)	16 (3%)	54	74

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ASP
1	A	151	THR
1	A	157	THR
1	A	176	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	302	ASP
1	A	339	GLU
1	A	389	ARG
1	A	395	ASN
1	A	416	ARG
1	A	472	THR
1	B	105	ARG
1	B	111	TYR
1	B	348	ARG
1	B	389	ARG
1	B	416	ARG
1	B	454	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	ASN
1	A	250	ASN
1	A	309	ASN
1	A	311	GLN
1	A	399	GLN
1	A	409	GLN
1	B	168	GLN
1	B	272	HIS
1	B	334	HIS
1	B	459	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	0MR	A	601	-	40,45,45	1.12	4 (10%)	48,64,64	1.40	9 (18%)
2	0MR	B	601	-	40,45,45	1.07	4 (10%)	48,64,64	1.31	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0MR	A	601	-	-	0/23/40/40	0/4/5/5
2	0MR	B	601	-	-	0/23/40/40	0/4/5/5

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	0MR	C11-C18	-4.84	1.44	1.49
2	B	601	0MR	C11-C18	-4.44	1.44	1.49
2	A	601	0MR	C12-C21	-2.31	1.47	1.51
2	B	601	0MR	C12-C21	-2.21	1.48	1.51
2	A	601	0MR	C22-N33	2.01	1.40	1.35
2	B	601	0MR	C22-N33	2.15	1.41	1.35
2	B	601	0MR	C21-N32	2.40	1.40	1.34
2	A	601	0MR	C21-N32	2.43	1.40	1.34

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	0MR	O40-C16-C6	-3.35	118.72	124.35
2	A	601	0MR	C8-C13-N31	-3.28	118.28	122.72
2	A	601	0MR	O40-C16-C6	-3.19	118.98	124.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	OMR	C8-C13-N31	-2.86	118.84	122.72
2	A	601	OMR	C5-C11-C16	2.01	120.02	117.36
2	A	601	OMR	C18-C12-C21	2.10	131.14	126.03
2	B	601	OMR	C12-C21-N32	2.16	120.01	117.76
2	B	601	OMR	C8-C14-C15	2.33	119.84	116.57
2	A	601	OMR	C17-C13-N31	2.36	121.99	119.97
2	A	601	OMR	C12-C21-N32	2.39	120.25	117.76
2	A	601	OMR	C8-C14-C15	2.49	120.07	116.57
2	B	601	OMR	C13-C17-CL4	2.86	122.58	120.18
2	A	601	OMR	O40-C16-C11	3.00	121.09	116.30
2	B	601	OMR	O40-C16-C11	3.19	121.39	116.30
2	A	601	OMR	C13-C17-CL4	3.69	123.27	120.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	OMR	5	0
2	B	601	OMR	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	437/499 (87%)	0.13	8 (1%) 71 71	26, 36, 50, 59	0
1	B	437/499 (87%)	0.38	27 (6%) 24 25	29, 47, 71, 76	0
All	All	874/998 (87%)	0.26	35 (4%) 42 43	26, 41, 68, 76	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	368	ASN	3.6
1	B	239	VAL	3.5
1	B	261	ARG	3.2
1	B	83	PRO	3.2
1	B	356	GLY	3.2
1	B	15	THR	3.1
1	B	222	MET	3.0
1	B	223	CYS	3.0
1	A	126	GLY	3.0
1	B	248	PRO	2.9
1	B	214	ARG	2.9
1	B	403	SER	2.8
1	B	406	ILE	2.7
1	B	262	SER	2.6
1	A	72	ASP	2.6
1	B	216	ARG	2.5
1	B	10	TYR	2.5
1	B	247	ASN	2.5
1	A	429	PHE	2.5
1	B	242	VAL	2.5
1	B	249	GLY	2.4
1	B	238	MET	2.4
1	B	424	THR	2.4
1	A	127	ASP	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	353	VAL	2.3
1	B	487	TYR	2.3
1	B	453	PRO	2.3
1	A	365	ILE	2.2
1	B	244	GLU	2.2
1	B	258	PHE	2.2
1	A	318	PRO	2.2
1	A	428	ALA	2.1
1	B	265	ILE	2.1
1	B	255	ASP	2.0
1	B	241	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	0MR	A	601	41/41	0.93	0.17	1.10	28,30,31,32	0
2	0MR	B	601	41/41	0.94	0.14	-0.45	31,34,35,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.