



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:42 AM GMT

PDB ID : 2E1W
Title : Crystal structure of adenosine deaminase complexed with potent inhibitors
Authors : Kinoshita, T.
Deposited on : 2006-10-30
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2935 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosine deaminase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	0	0	0
			2789	1772	471	534	12			


There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	ASP	ASN	SEE REMARK 999	UNP P56658
A	32	LYS	ARG	SEE REMARK 999	UNP P56658
A	33	ARG	LYS	SEE REMARK 999	UNP P56658
A	57	THR	SER	SEE REMARK 999	UNP P56658
A	60	ASP	GLU	SEE REMARK 999	UNP P56658
A	77	ASP	GLU	SEE REMARK 999	UNP P56658
A	79	ILE	VAL	SEE REMARK 999	UNP P56658
A	199	GLN	LYS	SEE REMARK 999	UNP P56658
A	246	THR	ALA	SEE REMARK 999	UNP P56658
A	261	ILE	VAL	SEE REMARK 999	UNP P56658
A	279	ALA	PRO	SEE REMARK 999	UNP P56658
A	281	ILE	VAL	SEE REMARK 999	UNP P56658
A	313	LYS	ASN	SEE REMARK 999	UNP P56658
A	314	ASP	GLU	SEE REMARK 999	UNP P56658
A	352	ARG	GLY	SEE REMARK 999	UNP P56658

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 1-[(1R,2S)-2-HYDROXY-1-[2-(1-NAPHTHYL)ETHYL]PROPYL]-1H-IMIDAZOLE-4-CARBOXAMIDE (three-letter code: FR6) (formula: C₁₉H₂₁N₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			24	19	3	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	121	Total	O	0	0
			121	121		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenosine deaminase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.41 Å 78.41 Å 137.46 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.79 – 2.50 51.68 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.79-2.50) 75.4 (51.68-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.11 (at 2.51 Å)	Xtriage
Refinement program	CNX	Depositor
R, R_{free}	0.218 , 0.246 0.199 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.209	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 64.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 11689 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2935	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FR6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/2853	1.78	51/3867 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	12

There are no bond length outliers.

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	CYS	CA-CB-SG	-11.74	92.86	114.00
1	A	33	ARG	NE-CZ-NH2	11.02	125.81	120.30
1	A	185	ASP	N-CA-CB	-10.79	91.17	110.60
1	A	184	GLY	N-CA-C	-9.91	88.33	113.10
1	A	81	ARG	NE-CZ-NH2	9.18	124.89	120.30
1	A	267	TYR	CB-CG-CD1	-8.82	115.71	121.00
1	A	283	PHE	CB-CG-CD2	-8.06	115.16	120.80
1	A	317	PHE	CB-CG-CD2	-7.81	115.33	120.80
1	A	235	ARG	NE-CZ-NH1	-7.74	116.43	120.30
1	A	283	PHE	CB-CG-CD1	7.51	126.06	120.80
1	A	317	PHE	N-CA-CB	6.85	122.92	110.60
1	A	253	ARG	NE-CZ-NH2	6.83	123.72	120.30
1	A	325	LEU	CA-CB-CG	6.77	130.86	115.30
1	A	253	ARG	NE-CZ-NH1	-6.57	117.01	120.30
1	A	194	PHE	CB-CG-CD2	-6.50	116.25	120.80
1	A	173	ARG	NE-CZ-NH2	-6.42	117.09	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	290	TYR	CB-CG-CD2	-6.28	117.23	121.00
1	A	300	PHE	CB-CG-CD2	-6.28	116.40	120.80
1	A	38	LEU	CB-CA-C	6.28	122.12	110.20
1	A	351	TYR	CB-CG-CD1	-6.18	117.29	121.00
1	A	310	MET	CG-SD-CE	-6.15	90.36	100.20
1	A	324	ARG	NE-CZ-NH2	6.11	123.36	120.30
1	A	68	TYR	CB-CG-CD2	-5.99	117.41	121.00
1	A	67	TYR	CB-CG-CD1	-5.95	117.43	121.00
1	A	352	ARG	NE-CZ-NH2	5.91	123.26	120.30
1	A	40	ALA	CB-CA-C	-5.91	101.24	110.10
1	A	235	ARG	NE-CZ-NH2	5.89	123.25	120.30
1	A	69	MET	CG-SD-CE	-5.88	90.78	100.20
1	A	30	TYR	CB-CG-CD2	-5.84	117.49	121.00
1	A	288	VAL	CA-CB-CG2	5.77	119.55	110.90
1	A	33	ARG	CD-NE-CZ	5.70	131.59	123.60
1	A	278	HIS	CA-CB-CG	-5.65	103.99	113.60
1	A	313	LYS	C-N-CA	-5.61	107.68	121.70
1	A	283	PHE	C-N-CA	-5.60	107.69	121.70
1	A	175	GLN	C-N-CA	5.54	135.56	121.70
1	A	73	ALA	C-N-CA	-5.46	110.84	122.30
1	A	175	GLN	CA-CB-CG	5.44	125.37	113.40
1	A	156	ARG	NE-CZ-NH2	-5.44	117.58	120.30
1	A	325	LEU	CB-CG-CD2	5.37	120.12	111.00
1	A	131	SER	N-CA-CB	5.35	118.52	110.50
1	A	96	VAL	CG1-CB-CG2	5.33	119.43	110.90
1	A	33	ARG	NE-CZ-NH1	-5.28	117.66	120.30
1	A	277	GLU	C-N-CA	-5.22	108.65	121.70
1	A	325	LEU	CB-CA-C	5.20	120.08	110.20
1	A	132	LEU	CA-CB-CG	-5.19	103.37	115.30
1	A	316	GLY	C-N-CA	-5.18	108.76	121.70
1	A	81	ARG	NE-CZ-NH1	-5.15	117.72	120.30
1	A	172	TYR	CD1-CG-CD2	5.12	123.53	117.90
1	A	290	TYR	CD1-CG-CD2	5.11	123.53	117.90
1	A	259	PHE	CB-CG-CD2	-5.11	117.22	120.80
1	A	32	LYS	C-N-CA	-5.04	109.11	121.70

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	173	ARG	Sidechain
1	A	201	TYR	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	267	TYR	Sidechain
1	A	282	ARG	Sidechain
1	A	324	ARG	Sidechain
1	A	33	ARG	Sidechain
1	A	348	TYR	Sidechain
1	A	36	ILE	Peptide
1	A	37	ALA	Peptide
1	A	67	TYR	Sidechain
1	A	68	TYR	Sidechain
1	A	84	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2789	0	2743	101	0
2	A	1	0	0	0	0
3	A	24	0	21	0	0
4	A	121	0	0	22	0
All	All	2935	0	2764	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:CYS:SG	1:A:180:ILE:HD11	2.23	0.78
1:A:152:LEU:HD23	1:A:169:CYS:SG	2.25	0.77
1:A:83:ALA:HB1	1:A:137:LEU:HG	1.68	0.74
1:A:261:ILE:O	1:A:293:ASN:HB2	1.88	0.74
1:A:180:ILE:HG12	1:A:181:ASP:N	2.09	0.68
1:A:131:SER:HA	1:A:134:ASN:OD1	1.94	0.67
1:A:80:LYS:HA	4:A:441:HOH:O	1.93	0.67
1:A:249:TYR:CE2	1:A:283:PHE:HE1	2.14	0.65
1:A:166:VAL:HG11	1:A:203:GLU:HB3	1.78	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:GLU:O	1:A:176:THR:HA	1.99	0.62
1:A:180:ILE:HD13	1:A:201:TYR:CE1	2.34	0.62
1:A:180:ILE:HD13	1:A:201:TYR:CZ	2.34	0.62
1:A:38:LEU:HB2	1:A:39:PRO:HD3	1.82	0.60
1:A:278:HIS:O	1:A:281:ILE:HG13	2.01	0.60
1:A:302:SER:HB2	1:A:306:THR:OG1	2.01	0.60
1:A:133:VAL:HA	4:A:441:HOH:O	2.00	0.60
1:A:312:LYS:HE2	1:A:313:LYS:HZ1	1.67	0.60
1:A:352:ARG:HG2	1:A:352:ARG:HH21	1.66	0.60
1:A:158:GLN:HG2	1:A:161:TRP:CE2	2.37	0.59
1:A:15:HIS:O	1:A:15:HIS:CD2	2.56	0.59
1:A:101:ARG:HB3	1:A:151:ILE:HB	1.84	0.59
1:A:292:LEU:HD13	1:A:325:LEU:HD11	1.83	0.59
1:A:281:ILE:HG22	1:A:284:LYS:HE3	1.84	0.59
1:A:237:GLY:HA2	1:A:260:GLU:HB2	1.85	0.59
1:A:56:LEU:HD13	1:A:60:ASP:HB3	1.85	0.59
1:A:241:HIS:O	1:A:244:GLU:HG2	2.03	0.58
1:A:310:MET:HG3	4:A:500:HOH:O	2.03	0.58
1:A:312:LYS:HE2	1:A:313:LYS:NZ	2.19	0.58
1:A:260:GLU:HA	4:A:516:HOH:O	2.02	0.58
1:A:50:ILE:HG22	4:A:413:HOH:O	2.03	0.58
1:A:15:HIS:HD2	1:A:15:HIS:O	1.86	0.57
1:A:185:ASP:OD1	1:A:187:THR:HB	2.05	0.56
1:A:47:GLN:HE21	1:A:47:GLN:HA	1.70	0.56
1:A:166:VAL:HB	4:A:448:HOH:O	2.07	0.54
1:A:235:ARG:HD2	1:A:260:GLU:OE1	2.08	0.53
1:A:15:HIS:HA	1:A:101:ARG:NH2	2.23	0.52
1:A:46:LEU:HD13	4:A:509:HOH:O	2.07	0.52
1:A:63:ALA:HB1	4:A:453:HOH:O	2.09	0.51
1:A:313:LYS:HB3	1:A:313:LYS:NZ	2.26	0.51
1:A:83:ALA:HB2	1:A:133:VAL:HG13	1.93	0.51
1:A:152:LEU:CD2	1:A:169:CYS:SG	2.99	0.50
1:A:249:TYR:CE2	1:A:283:PHE:CE1	2.98	0.50
1:A:101:ARG:HB2	1:A:153:CYS:SG	2.52	0.50
1:A:31:GLY:HA3	1:A:38:LEU:HD22	1.94	0.50
1:A:178:VAL:O	1:A:210:HIS:HB2	2.12	0.49
1:A:308:TYR:O	1:A:312:LYS:HB2	2.12	0.49
1:A:259:PHE:HB2	4:A:427:HOH:O	2.12	0.48
1:A:117:TRP:HA	4:A:481:HOH:O	2.13	0.48
1:A:181:ASP:HA	1:A:212:THR:O	2.14	0.48
1:A:260:GLU:HG3	4:A:516:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:VAL:HG21	1:A:200:ALA:HA	1.95	0.48
1:A:39:PRO:HG2	1:A:49:ILE:HD11	1.96	0.47
1:A:46:LEU:HD12	4:A:440:HOH:O	2.13	0.47
1:A:293:ASN:HB3	4:A:420:HOH:O	2.13	0.47
1:A:235:ARG:HD3	1:A:258:HIS:CD2	2.50	0.47
1:A:173:ARG:O	1:A:175:GLN:N	2.48	0.47
1:A:55:PRO:HA	1:A:268:LEU:O	2.14	0.46
1:A:239:GLY:O	1:A:242:THR:HG23	2.16	0.46
1:A:249:TYR:HE2	1:A:283:PHE:CE1	2.32	0.46
1:A:263:PRO:HB2	1:A:310:MET:CE	2.46	0.46
1:A:224:VAL:HG21	1:A:242:THR:HG22	1.98	0.45
1:A:63:ALA:O	1:A:66:ASP:HB2	2.16	0.45
1:A:39:PRO:HB3	4:A:464:HOH:O	2.16	0.45
1:A:278:HIS:CD2	1:A:280:VAL:HG12	2.52	0.45
1:A:9:LYS:HE3	1:A:9:LYS:HB3	1.85	0.45
1:A:188:ILE:HD13	1:A:188:ILE:HG21	1.75	0.44
1:A:61:PHE:CE2	1:A:65:PHE:HE1	2.36	0.44
1:A:180:ILE:HD12	4:A:472:HOH:O	2.17	0.44
1:A:213:VAL:N	1:A:233:THR:HG21	2.32	0.44
1:A:250:ASN:O	1:A:253:ARG:HG3	2.17	0.44
1:A:258:HIS:HA	1:A:289:ASN:O	2.18	0.43
1:A:113:GLU:HA	1:A:114:PRO:HA	1.76	0.43
1:A:317:PHE:HB2	4:A:468:HOH:O	2.18	0.43
1:A:22:ILE:HD13	1:A:72:ILE:HD13	2.00	0.43
1:A:313:LYS:HB3	1:A:313:LYS:HZ2	1.82	0.43
1:A:117:TRP:HZ3	4:A:482:HOH:O	2.02	0.42
1:A:330:ALA:O	1:A:340:LYS:HD2	2.19	0.42
1:A:33:ARG:HB3	1:A:33:ARG:HH21	1.83	0.42
1:A:238:HIS:HB3	1:A:240:TYR:CZ	2.55	0.42
1:A:158:GLN:HG2	1:A:161:TRP:CD2	2.54	0.42
1:A:27:ILE:HG21	1:A:68:TYR:HB2	2.02	0.42
1:A:33:ARG:CB	1:A:33:ARG:HH21	2.33	0.41
1:A:128:GLU:O	1:A:132:LEU:HD12	2.20	0.41
1:A:312:LYS:HB3	1:A:313:LYS:NZ	2.36	0.41
1:A:288:VAL:HG22	4:A:478:HOH:O	2.20	0.41
1:A:201:TYR:HD2	1:A:211:ARG:HD2	1.85	0.41
1:A:263:PRO:HB2	1:A:310:MET:HE3	2.02	0.41
1:A:97:TYR:OH	1:A:335:LEU:HD13	2.20	0.41
1:A:21:ALA:HB2	1:A:297:PRO:HD2	2.02	0.41
1:A:236:LEU:O	1:A:260:GLU:HB2	2.20	0.41
1:A:120:ALA:HB1	4:A:423:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:VAL:HG22	4:A:495:HOH:O	2.21	0.41
1:A:21:ALA:HB3	4:A:497:HOH:O	2.20	0.41
1:A:33:ARG:HB3	1:A:33:ARG:NH2	2.36	0.41
1:A:57:THR:OG1	1:A:59:PRO:HD2	2.21	0.41
1:A:228:VAL:HG22	4:A:475:HOH:O	2.20	0.41
1:A:278:HIS:HD2	1:A:280:VAL:HG12	1.86	0.40
1:A:262:CYS:SG	1:A:295:ASP:HB2	2.62	0.40
1:A:28:LEU:O	1:A:32:LYS:HG2	2.22	0.40
1:A:238:HIS:HB3	1:A:240:TYR:CE2	2.56	0.40
1:A:27:ILE:HD13	1:A:68:TYR:CB	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	347/356 (98%)	305 (88%)	35 (10%)	7 (2%)	9 15

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	ALA
1	A	174	GLU
1	A	112	VAL
1	A	186	GLU
1	A	238	HIS
1	A	38	LEU
1	A	173	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	304/309 (98%)	244 (80%)	60 (20%)	1 3

All (60) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ASP
1	A	11	LYS
1	A	25	GLU
1	A	28	LEU
1	A	32	LYS
1	A	33	ARG
1	A	41	ASP
1	A	45	GLU
1	A	46	LEU
1	A	47	GLN
1	A	49	ILE
1	A	50	ILE
1	A	53	ASP
1	A	57	THR
1	A	62	LEU
1	A	75	CYS
1	A	80	LYS
1	A	93	ASP
1	A	101	ARG
1	A	103	SER
1	A	107	LEU
1	A	124	LEU
1	A	148	VAL
1	A	152	LEU
1	A	160	SER
1	A	168	LEU
1	A	174	GLU
1	A	180	ILE
1	A	187	THR
1	A	188	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	192	SER
1	A	193	LEU
1	A	199	GLN
1	A	205	VAL
1	A	218	VAL
1	A	222	ASN
1	A	225	LYS
1	A	233	THR
1	A	242	THR
1	A	253	ARG
1	A	266	SER
1	A	267	TYR
1	A	277	GLU
1	A	280	VAL
1	A	281	ILE
1	A	284	LYS
1	A	288	VAL
1	A	294	THR
1	A	298	LEU
1	A	305	ASP
1	A	312	LYS
1	A	313	LYS
1	A	317	PHE
1	A	319	GLU
1	A	325	LEU
1	A	335	LEU
1	A	344	LEU
1	A	347	LEU
1	A	349	LYS
1	A	352	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	GLN
1	A	175	GLN
1	A	197	HIS
1	A	210	HIS
1	A	309	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	FR6	A	1	-	22,26,26	1.89	2 (9%)	25,36,36	1.56	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FR6	A	1	-	-	0/9/17/17	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1	FR6	C5-N4	3.45	1.48	1.37
3	A	1	FR6	C1-N2	6.53	1.47	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	1	FR6	O9-C8-N10	-3.02	118.34	122.59
3	A	1	FR6	C2-C28-C27	-2.37	117.63	123.22
3	A	1	FR6	O9-C8-C5	5.47	124.72	119.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	349/356 (98%)	-0.28	1 (0%) 94 95	7, 20, 35, 44	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	GLU	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	FR6	A	1	24/24	0.95	0.13	0.32	10,16,20,21	0
2	ZN	A	400	1/1	0.96	0.11	-	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.