



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:19 AM GMT

PDB ID : 3E6N  
Title : Structure of murine INOS oxygenase domain with inhibitor AR-C125813  
Authors : Garcin, E.D.; Arvai, A.S.; Rosenfeld, R.J.; Kroeger, M.D.; Crane, B.R.; Andersson, G.; Andrews, G.; Hamley, P.J.; Mallinder, P.R.; Nicholls, D.J.; St-Gallay, S.A.; Tinker, A.C.; Gensmantel, N.P.; Mete, A.; Cheshire, D.R.; Connolly, S.; Stueh, D.J.; Aberg, A.; Wallace, A.V.; Tainer, J.A.; Getzoff, E.D.  
Deposited on : 2008-08-15  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	
1	B	433	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	H4B	A	902	X	-	-	-
3	H4B	B	902	X	-	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, inducible.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	413	Total	C	N	O	S	0	0	0
			3363	2157	580	606	20			
1	B	413	Total	C	N	O	S	0	0	0
			3363	2157	580	606	20			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 4-METHYL-6-PROPYLPYRIDIN-2-AMINE (three-letter code: AS3) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			11	9	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	0	0
			11	9	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	253	Total	O	0	0
			253	253		
5	B	240	Total	O	0	0
			240	240		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, inducible



- Molecule 1: Nitric oxide synthase, inducible



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	213.96 Å   213.96 Å   112.08 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	19.99 – 2.40 19.99 – 2.40	Depositor EDS
% Data completeness (in resolution range)	87.0 (19.99-2.40) 87.0 (19.99-2.40)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.33 (at 2.41 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.233   ,   0.270 0.224   ,   0.261	Depositor DCC
$R_{free}$ test set	2923 reflections (5.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.0	Xtriage
Anisotropy	0.651	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 50.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 57844 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7361	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AS3, HEM, H4B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/3461	0.64	0/4706
1	B	0.38	0/3461	0.63	0/4706
All	All	0.39	0/6922	0.63	0/9412

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3363	0	3256	49	0
1	B	3363	0	3256	50	0
2	A	43	0	30	3	0
2	B	43	0	30	1	0
3	A	17	0	14	0	0
3	B	17	0	14	0	0
4	A	11	0	14	2	0
4	B	11	0	14	1	0
5	A	253	0	0	8	0
5	B	240	0	0	4	0
All	All	7361	0	6628	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:488:GLU:HG3	5:A:952:HOH:O	1.78	0.82
1:B:83:ASN:HD22	1:B:85:GLY:H	1.33	0.76
1:B:285:ILE:HG12	1:B:291:PRO:HG3	1.65	0.76
1:B:83:ASN:ND2	1:B:85:GLY:H	1.85	0.75
1:A:410:THR:O	1:A:414:VAL:HG23	1.85	0.75
1:B:215:GLN:HE21	1:B:219:GLN:HE21	1.36	0.71
1:B:308:GLN:NE2	1:B:496:GLN:HE22	1.89	0.70
1:B:330:TYR:HD2	1:B:332:TRP:HE1	1.41	0.69
1:B:188:TRP:CE3	1:B:200:TRP:HA	2.29	0.68
1:B:342:ALA:HB1	1:B:425:VAL:HG11	1.77	0.67
1:B:260:ARG:HG3	5:B:925:HOH:O	1.96	0.66
1:A:165:LYS:HG2	5:A:1060:HOH:O	1.96	0.65
1:A:465:SER:O	1:A:471:HIS:HE1	1.81	0.63
1:A:326:GLU:HG2	1:A:334:GLN:HG2	1.80	0.63
1:A:215:GLN:HE21	1:A:219:GLN:HE21	1.47	0.61
1:A:188:TRP:CE3	1:A:200:TRP:HA	2.36	0.60
1:A:83:ASN:ND2	1:A:85:GLY:H	2.00	0.60
1:B:442:GLN:HA	1:B:442:GLN:HE21	1.66	0.59
1:A:379:ASP:HB3	1:A:381:GLN:OE1	2.03	0.58
1:A:260:ARG:HG2	1:A:277:THR:HG21	1.86	0.57
1:A:438:MET:HG3	1:A:468:PRO:HB2	1.88	0.56
1:B:282:GLN:HA	1:B:282:GLN:HE21	1.69	0.56
1:A:325:MET:HA	1:A:325:MET:HE2	1.88	0.55
1:B:110:LEU:HG	1:B:113:ILE:HG13	1.89	0.54
1:A:317:PRO:HD2	5:A:924:HOH:O	2.07	0.54
1:B:215:GLN:HE21	1:B:219:GLN:NE2	2.05	0.53
1:B:308:GLN:HE21	1:B:496:GLN:HE22	1.54	0.53
1:A:389:VAL:O	1:A:393:MET:HG3	2.09	0.53
1:A:459:VAL:HG22	1:A:469:VAL:HG12	1.90	0.53
1:B:289:TRP:O	1:B:291:PRO:HD3	2.08	0.52
2:A:901:HEM:HBC2	2:A:901:HEM:HMC2	1.90	0.52
1:A:139:GLU:HG2	5:A:1136:HOH:O	2.08	0.52
1:B:134:LEU:O	1:B:138:ILE:HG12	2.10	0.52
1:A:81:ILE:HD11	1:A:475:LEU:HD13	1.91	0.52
1:A:438:MET:O	1:A:442:GLN:HG2	2.09	0.52
1:A:246:ASP:OD1	1:A:249:HIS:HD2	1.93	0.52
1:B:190:ASN:O	1:B:192:PRO:HD3	2.11	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:83:ASN:HD22	1:B:83:ASN:C	2.13	0.51
2:A:901:HEM:HBA2	4:A:903:AS3:H92	1.92	0.51
1:A:241:PHE:HB3	1:A:242:PRO:CD	2.41	0.51
1:A:134:LEU:O	1:A:138:ILE:HG12	2.11	0.51
1:B:442:GLN:CA	1:B:442:GLN:HE21	2.24	0.51
1:B:330:TYR:HD2	1:B:332:TRP:NE1	2.06	0.51
1:A:190:ASN:O	1:A:192:PRO:HD3	2.11	0.50
1:B:465:SER:O	1:B:471:HIS:HE1	1.95	0.50
1:A:327:HIS:ND1	1:A:328:PRO:HD2	2.27	0.49
1:B:81:ILE:HD11	1:B:475:LEU:HD13	1.95	0.49
1:A:264:TYR:CE1	1:A:293:TYR:HA	2.46	0.49
1:A:393:MET:CE	1:A:411:GLU:HG3	2.42	0.49
1:A:348:ASN:HB2	5:A:1004:HOH:O	2.12	0.49
1:A:324:THR:HG23	5:A:1132:HOH:O	2.12	0.48
1:A:188:TRP:CZ3	1:A:200:TRP:HA	2.48	0.48
1:A:88:GLU:HB3	5:A:1135:HOH:O	2.14	0.47
1:A:239:THR:O	1:A:361:CYS:HA	2.14	0.47
1:B:282:GLN:HA	1:B:282:GLN:NE2	2.30	0.47
1:A:249:HIS:HB3	1:A:306:ASP:OD1	2.14	0.46
1:A:326:GLU:HG2	1:A:334:GLN:CG	2.45	0.46
1:A:332:TRP:HB2	5:A:947:HOH:O	2.16	0.45
1:A:324:THR:H	1:A:423:GLN:HE22	1.63	0.45
1:B:399:THR:O	1:B:402:SER:HB3	2.16	0.45
1:B:298:VAL:HG21	1:B:320:VAL:HG11	1.99	0.45
1:B:326:GLU:HG2	1:B:334:GLN:HG2	1.99	0.45
1:A:78:TYR:HE1	1:A:80:ARG:HG3	1.82	0.44
1:A:86:SER:OG	1:A:88:GLU:HG2	2.18	0.44
1:A:289:TRP:O	1:A:291:PRO:HD3	2.17	0.44
2:B:901:HEM:C1C	4:B:903:AS3:H71	2.52	0.44
1:B:139:GLU:HG2	5:B:1137:HOH:O	2.18	0.44
1:B:342:ALA:CB	1:B:425:VAL:HG11	2.46	0.43
1:A:330:TYR:HB3	1:A:332:TRP:CE2	2.53	0.43
1:B:438:MET:O	1:B:442:GLN:HG2	2.19	0.43
1:B:119:LEU:HD21	5:B:1021:HOH:O	2.18	0.43
1:B:137:ALA:O	1:B:141:ILE:HG12	2.19	0.43
1:B:252:ARG:NH2	1:B:489:PRO:HD3	2.33	0.43
1:B:407:ARG:HD2	5:B:1123:HOH:O	2.19	0.43
1:B:239:THR:O	1:B:361:CYS:HA	2.18	0.43
1:B:385:ILE:O	1:B:389:VAL:HG23	2.19	0.42
1:B:81:ILE:HD11	1:B:475:LEU:CD1	2.49	0.42
1:A:125:ASP:OD1	1:A:126:LYS:HG3	2.19	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:252:ARG:HH21	1:B:489:PRO:HD3	1.85	0.42
1:B:134:LEU:HB3	1:B:135:PRO:HD3	2.01	0.42
1:A:241:PHE:HB3	1:A:242:PRO:HD2	2.01	0.42
1:B:188:TRP:CZ3	1:B:200:TRP:HA	2.53	0.42
1:A:301:LEU:HB3	1:A:303:LEU:HD21	2.02	0.42
1:B:326:GLU:HG2	1:B:334:GLN:CG	2.50	0.42
1:B:393:MET:CE	1:B:411:GLU:HG3	2.50	0.42
1:A:330:TYR:CD1	1:A:330:TYR:N	2.88	0.41
1:B:304:GLN:HG3	1:B:308:GLN:O	2.19	0.41
1:B:83:ASN:HD22	1:B:85:GLY:N	2.11	0.41
1:A:83:ASN:C	1:A:83:ASN:HD22	2.23	0.41
1:A:445:TYR:CE2	1:A:450:GLY:HA2	2.55	0.41
1:B:291:PRO:HB2	1:B:293:TYR:CE2	2.56	0.41
1:B:209:ARG:O	1:B:242:PRO:HG3	2.20	0.41
1:A:283:LEU:O	1:A:287:LEU:HG	2.21	0.41
1:A:323:VAL:HG22	1:A:425:VAL:HG21	2.02	0.41
1:A:80:ARG:NH1	1:A:89:ILE:HG21	2.36	0.41
1:B:332:TRP:CE3	1:B:392:ARG:HD3	2.54	0.41
1:B:453:ALA:HB1	1:B:458:LEU:CD1	2.51	0.41
1:B:453:ALA:HB1	1:B:458:LEU:HD12	2.03	0.41
1:A:83:ASN:HB2	1:A:472:GLN:OE1	2.21	0.40
1:B:83:ASN:HD22	1:B:84:TRP:N	2.19	0.40
1:B:459:VAL:HG22	1:B:469:VAL:HG13	2.02	0.40
2:A:901:HEM:C1C	4:A:903:AS3:H71	2.57	0.40
1:A:265:GLN:HG3	1:A:265:GLN:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/433 (94%)	378 (92%)	28 (7%)	3 (1%)	26	38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	409/433 (94%)	381 (93%)	27 (7%)	1 (0%)	52	69
All	All	818/866 (94%)	759 (93%)	55 (7%)	4 (0%)	34	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	ASP
1	A	293	TYR
1	B	268	ASP
1	A	464	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/381 (94%)	350 (98%)	9 (2%)	55	76
1	B	359/381 (94%)	350 (98%)	9 (2%)	55	76
All	All	718/762 (94%)	700 (98%)	18 (2%)	55	76

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	99	THR
1	A	114	MET
1	A	223	ARG
1	A	260	ARG
1	A	392	ARG
1	A	406	ASP
1	A	409	VAL
1	A	475	LEU
1	B	80	ARG
1	B	83	ASN
1	B	223	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	260	ARG
1	B	282	GLN
1	B	441	MET
1	B	442	GLN
1	B	469	VAL
1	B	475	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	91	HIS
1	A	219	GLN
1	A	249	HIS
1	A	257	GLN
1	A	442	GLN
1	A	471	HIS
1	B	83	ASN
1	B	96	HIS
1	B	219	GLN
1	B	282	GLN
1	B	308	GLN
1	B	442	GLN
1	B	471	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	901	1	30,50,50	2.61	7 (23%)	24,82,82	2.58	11 (45%)
3	H4B	A	902	-	13,18,18	2.43	2 (15%)	11,26,26	2.64	6 (54%)
4	AS3	A	903	-	11,11,11	0.55	0	14,14,14	1.78	3 (21%)
2	HEM	B	901	1	30,50,50	2.49	9 (30%)	24,82,82	2.44	8 (33%)
3	H4B	B	902	-	13,18,18	2.46	3 (23%)	11,26,26	2.52	5 (45%)
4	AS3	B	903	-	11,11,11	0.48	0	14,14,14	1.91	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	901	1	-	0/10/54/54	0/0/8/8
3	H4B	A	902	-	1/1/3/5	0/8/17/17	0/2/2/2
4	AS3	A	903	-	-	0/3/3/3	0/1/1/1
2	HEM	B	901	1	-	0/10/54/54	0/0/8/8
3	H4B	B	902	-	1/1/3/5	0/8/17/17	0/2/2/2
4	AS3	B	903	-	-	0/3/3/3	0/1/1/1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	HEM	C2D-C3D	-7.08	1.33	1.54
2	B	901	HEM	C2D-C3D	-6.71	1.34	1.54
2	A	901	HEM	C3B-CAB	-6.35	1.39	1.51
3	A	902	H4B	C6-N5	-6.28	1.33	1.45
3	B	902	H4B	C6-N5	-6.23	1.33	1.45
2	B	901	HEM	C3C-CAC	-6.12	1.39	1.51
2	A	901	HEM	C3C-CAC	-5.85	1.40	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	902	H4B	C7-N8	-5.83	1.38	1.46
3	A	902	H4B	C7-N8	-5.63	1.38	1.46
2	B	901	HEM	C3B-CAB	-5.34	1.41	1.51
2	A	901	HEM	C2C-C1C	-4.45	1.44	1.52
2	B	901	HEM	C3D-C4D	-3.85	1.46	1.51
2	A	901	HEM	C3D-C4D	-3.61	1.46	1.51
2	A	901	HEM	C3B-C4B	-3.52	1.48	1.51
2	A	901	HEM	C2B-C1B	-3.31	1.41	1.51
2	B	901	HEM	C2B-C1B	-3.30	1.41	1.51
2	B	901	HEM	C2C-C1C	-3.00	1.46	1.52
2	B	901	HEM	C2D-C1D	-2.53	1.43	1.51
3	B	902	H4B	C4A-N5	-2.02	1.33	1.38
2	B	901	HEM	CHC-C1C	2.22	1.41	1.36
2	B	901	HEM	CBC-CAC	2.52	1.43	1.29

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	903	AS3	C4-C5-N6	-3.39	119.14	122.96
4	A	903	AS3	C4-C5-N6	-3.26	119.28	122.96
2	A	901	HEM	CMA-C3A-C4A	-2.75	123.82	128.36
2	A	901	HEM	CAA-C2A-C1A	-2.55	124.23	127.01
3	A	902	H4B	N3-C2-N1	-2.49	121.45	125.53
2	B	901	HEM	CBA-CAA-C2A	-2.30	108.41	112.53
3	B	902	H4B	N3-C2-N1	-2.26	121.83	125.53
2	A	901	HEM	CBA-CAA-C2A	-2.06	108.84	112.53
4	A	903	AS3	N8-C1-N6	2.03	120.20	116.50
3	A	902	H4B	C8A-C4A-N5	2.12	121.52	118.85
2	A	901	HEM	C3B-C4B-CHC	2.32	126.43	123.16
3	B	902	H4B	C2-N1-C8A	2.58	120.35	114.54
2	A	901	HEM	CMD-C2D-C3D	2.61	125.91	114.35
3	A	902	H4B	C4-C4A-C8A	2.72	117.03	114.56
2	B	901	HEM	CMD-C2D-C3D	2.74	126.48	114.35
3	B	902	H4B	C4-C4A-C8A	2.77	117.07	114.56
3	A	902	H4B	C2-N1-C8A	2.84	120.93	114.54
3	B	902	H4B	C4-N3-C2	3.13	120.28	115.94
2	B	901	HEM	CAD-C3D-C2D	3.44	123.12	113.22
3	A	902	H4B	C4-N3-C2	3.54	120.85	115.94
2	A	901	HEM	C2D-C3D-C4D	3.55	107.53	101.50
2	B	901	HEM	C2D-C3D-C4D	3.71	107.78	101.50
2	A	901	HEM	C3B-CAB-CBB	3.71	130.15	124.46
2	A	901	HEM	CAD-C3D-C2D	4.10	125.02	113.22

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	HEM	CAD-C3D-C4D	4.23	127.40	112.47
2	B	901	HEM	C3B-C4B-CHC	4.25	129.15	123.16
2	B	901	HEM	CAD-C3D-C4D	4.57	128.58	112.47
4	A	903	AS3	C1-N6-C5	4.76	121.61	118.23
2	A	901	HEM	CMB-C2B-C3B	4.83	128.59	116.53
2	B	901	HEM	CMC-C2C-C3C	5.22	129.56	116.53
2	B	901	HEM	CMB-C2B-C3B	5.26	129.67	116.53
4	B	903	AS3	C1-N6-C5	5.42	122.08	118.23
3	B	902	H4B	C7-C6-N5	5.68	122.24	110.45
3	A	902	H4B	C7-C6-N5	5.72	122.33	110.45
2	A	901	HEM	CMC-C2C-C3C	6.17	131.92	116.53

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	902	H4B	C6
3	A	902	H4B	C6

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	HEM	3	0
4	A	903	AS3	2	0
2	B	901	HEM	1	0
4	B	903	AS3	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	413/433 (95%)	-0.05	19 (4%) 36 37	25, 47, 79, 98	0
1	B	413/433 (95%)	-0.05	19 (4%) 36 37	27, 46, 74, 95	0
All	All	826/866 (95%)	-0.05	38 (4%) 36 37	25, 46, 76, 98	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	108	SER	10.6
1	A	108	SER	7.2
1	B	109	CYS	5.6
1	B	270	THR	4.8
1	A	270	THR	4.4
1	B	497	ASN	4.1
1	B	494	ILE	4.1
1	A	497	ASN	4.0
1	A	109	CYS	3.9
1	B	269	GLY	3.8
1	B	267	PRO	3.7
1	B	149	LYS	3.6
1	A	267	PRO	3.2
1	A	268	ASP	3.1
1	A	153	ILE	3.1
1	A	149	LYS	3.0
1	A	494	ILE	2.9
1	B	330	TYR	2.7
1	B	496	GLN	2.6
1	A	269	GLY	2.6
1	A	290	LYS	2.6
1	B	153	ILE	2.5
1	B	268	ASP	2.5
1	A	125	ASP	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	398	HIS	2.4
1	B	491	LYS	2.4
1	A	412	ILE	2.3
1	A	295	ARG	2.3
1	A	331	GLU	2.3
1	B	110	LEU	2.3
1	B	331	GLU	2.3
1	B	148	PHE	2.2
1	B	96	HIS	2.2
1	B	422	LYS	2.1
1	B	295	ARG	2.1
1	A	96	HIS	2.1
1	A	99	THR	2.0
1	A	154	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	AS3	B	903	11/11	0.98	0.12	0.18	24,26,29,30	0
2	HEM	A	901	43/43	0.98	0.11	-0.60	19,25,33,35	0
4	AS3	A	903	11/11	0.98	0.10	-0.76	21,26,32,33	0
2	HEM	B	901	43/43	0.98	0.09	-0.93	21,26,33,35	0
3	H4B	B	902	17/17	0.96	0.09	-1.06	28,32,36,37	0
3	H4B	A	902	17/17	0.95	0.09	-1.44	30,33,36,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.