



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:47 AM GMT

PDB ID : 2EBA  
Title : Crystal structure of the putative glutaryl-CoA dehydrogenase from thermus thermophilus  
Authors : Kumarevel, T.S.; Karthe, P.; Kuramitsu, S.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2007-02-07  
Resolution : 2.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*




The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	 61% 32% 5% •
1	C	385	 54% 39% 6% •
1	D	385	 55% 37% 6% •
1	E	385	 54% 39% 5% •
1	F	385	 56% 36% 6% •

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	385	<div><div></div><div>58%32%8%<div><div></div><div></div></div></div></div>
1	H	385	<div><div></div><div>60%34%5%<div><div></div><div></div></div></div></div>
1	I	385	<div><div></div><div>59%34%5%<div><div></div><div></div></div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative glutaryl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			
1	C	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			
1	D	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			
1	E	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			
1	F	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			
1	G	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			
1	H	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			
1	I	380	Total	C	N	O	S	0	0	0
			2986	1915	517	541	13			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	G	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	H	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	I	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total	O	0	0
			87	87		
3	C	77	Total	O	0	0
			77	77		
3	D	71	Total	O	0	0
			71	71		
3	E	71	Total	O	0	0
			71	71		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	55	Total 55	O 55	0	0
3	G	83	Total 83	O 83	0	0
3	H	84	Total 84	O 84	0	0
3	I	72	Total 72	O 72	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Putative glutaryl-CoA dehydrogenase




#### • Molecule 1: Putative glutaryl-CoA dehydrogenase




#### • Molecule 1: Putative glutaryl-CoA dehydrogenase

Chain D:  55% 37% 6%




• Molecule 1: Putative glutaryl-CoA dehydrogenase

Chain E:  54% 39% 5%




• Molecule 1: Putative glutaryl-CoA dehydrogenase


Chain F:  56% 36% 6%









• Molecule 1: Putative glutaryl-CoA dehydrogenase




• Molecule 1: Putative glutaryl-CoA dehydrogenase



• Molecule 1: Putative glutaryl-CoA dehydrogenase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.24Å 86.24Å 157.42Å 75.92° 81.25° 64.24°	Depositor
Resolution (Å)	19.96 – 2.21 40.19 – 2.21	Depositor EDS
% Data completeness (in resolution range)	72.8 (19.96-2.21) 68.8 (40.19-2.21)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.76 (at 2.22Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.260 , 0.275 0.264 , 0.278	Depositor DCC
$R_{free}$ test set	850 reflections (0.68%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.6	Xtriage
Anisotropy	0.612	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 20.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 131981 reflections	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	24912	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	1/3053 (0.0%)	0.72	3/4131 (0.1%)
1	C	0.56	0/3053	0.79	2/4131 (0.0%)
1	D	0.51	0/3053	0.77	3/4131 (0.1%)
1	E	0.52	0/3053	0.77	5/4131 (0.1%)
1	F	0.46	0/3053	0.70	1/4131 (0.0%)
1	G	0.57	3/3053 (0.1%)	0.72	4/4131 (0.1%)
1	H	0.44	0/3053	0.69	2/4131 (0.0%)
1	I	0.40	0/3053	0.64	1/4131 (0.0%)
All	All	0.50	4/24424 (0.0%)	0.73	21/33048 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	155	TRP	CB-CG	-14.56	1.24	1.50
1	G	155	TRP	CD1-NE1	-8.99	1.22	1.38
1	A	274	GLY	CA-C	-5.21	1.43	1.51
1	G	155	TRP	CE3-CZ3	-5.13	1.29	1.38

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	275	GLU	N-CA-C	-9.85	84.41	111.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	274	GLY	N-CA-C	8.46	134.26	113.10
1	D	31	LEU	CA-CB-CG	-7.81	97.34	115.30
1	H	61	LEU	CB-CG-CD1	-6.98	99.13	111.00
1	E	153	ASP	C-N-CA	-6.94	104.35	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	154	THR	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2986	0	3010	162	0
1	C	2986	0	3010	213	1
1	D	2986	0	3010	204	0
1	E	2986	0	3010	206	0
1	F	2986	0	3010	195	0
1	G	2986	0	3010	198	1
1	H	2986	0	3010	199	0
1	I	2986	0	3010	207	1
2	A	53	0	31	4	0
2	C	53	0	31	5	0
2	D	53	0	31	5	0
2	E	53	0	31	3	0
2	F	53	0	31	2	0
2	G	53	0	31	1	0
2	H	53	0	31	1	0
2	I	53	0	31	4	0
3	A	87	0	0	3	0
3	C	77	0	0	4	0
3	D	71	0	0	8	0
3	E	71	0	0	10	1
3	F	55	0	0	4	0
3	G	83	0	0	11	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	84	0	0	14	0
3	I	72	0	0	8	0
All	All	24912	0	24328	1423	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 1423 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:253:MET:CE	1:C:294:HIS:HA	1.60	1.31
1:A:253:MET:CE	1:A:294:HIS:HA	1.72	1.20
1:F:253:MET:CE	1:F:294:HIS:HA	1.74	1.15
1:F:203:ARG:CD	1:G:35:ARG:HH21	1.59	1.15
1:A:19:GLN:NE2	1:A:20:LYS:HE2	1.62	1.15

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:63:PRO:O	1:G:24:ARG:NH1[1_645]	2.10	0.10
1:I:64:GLU:OE1	3:E:4008:HOH:O[1_556]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/385 (97%)	359 (96%)	14 (4%)	1 (0%)	46	50
1	C	374/385 (97%)	351 (94%)	19 (5%)	4 (1%)	17	14
1	D	374/385 (97%)	356 (95%)	16 (4%)	2 (0%)	34	34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	374/385 (97%)	355 (95%)	19 (5%)	0	100	100
1	F	374/385 (97%)	352 (94%)	21 (6%)	1 (0%)	46	50
1	G	374/385 (97%)	357 (96%)	13 (4%)	4 (1%)	17	14
1	H	374/385 (97%)	360 (96%)	14 (4%)	0	100	100
1	I	374/385 (97%)	360 (96%)	12 (3%)	2 (0%)	34	34
All	All	2992/3080 (97%)	2850 (95%)	128 (4%)	14 (0%)	34	34

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	188	LEU
1	C	275	GLU
1	D	275	GLU
1	D	276	PRO
1	G	275	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/310 (99%)	280 (91%)	28 (9%)	12	10
1	C	308/310 (99%)	282 (92%)	26 (8%)	14	12
1	D	308/310 (99%)	283 (92%)	25 (8%)	15	13
1	E	308/310 (99%)	277 (90%)	31 (10%)	9	8
1	F	308/310 (99%)	278 (90%)	30 (10%)	10	8
1	G	308/310 (99%)	274 (89%)	34 (11%)	8	6
1	H	308/310 (99%)	281 (91%)	27 (9%)	12	11
1	I	308/310 (99%)	286 (93%)	22 (7%)	18	18
All	All	2464/2480 (99%)	2241 (91%)	223 (9%)	12	10

5 of 223 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	234	LEU
1	F	193	THR
1	I	78	ILE
1	E	271	SER
1	F	19	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	284	GLN
1	F	281	GLN
1	I	284	GLN
1	E	325	ASN
1	E	368	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	1001	-	48,58,58	1.97	10 (20%)	54,89,89	2.97	17 (31%)
2	FAD	C	2001	-	48,58,58	1.92	9 (18%)	54,89,89	2.93	18 (33%)
2	FAD	D	3001	-	48,58,58	1.83	10 (20%)	54,89,89	2.82	19 (35%)
2	FAD	E	4001	-	48,58,58	1.94	9 (18%)	54,89,89	2.98	18 (33%)
2	FAD	F	5001	-	48,58,58	1.92	7 (14%)	54,89,89	2.95	19 (35%)
2	FAD	G	6001	-	48,58,58	1.93	10 (20%)	54,89,89	2.93	18 (33%)
2	FAD	H	7001	-	48,58,58	1.90	8 (16%)	54,89,89	2.88	19 (35%)
2	FAD	I	8001	-	48,58,58	1.84	10 (20%)	54,89,89	2.84	17 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1001	-	-	0/30/50/50	0/6/6/6
2	FAD	C	2001	-	-	0/30/50/50	0/6/6/6
2	FAD	D	3001	-	-	0/30/50/50	0/6/6/6
2	FAD	E	4001	-	-	0/30/50/50	0/6/6/6
2	FAD	F	5001	-	-	0/30/50/50	0/6/6/6
2	FAD	G	6001	-	-	0/30/50/50	0/6/6/6
2	FAD	H	7001	-	-	0/30/50/50	0/6/6/6
2	FAD	I	8001	-	-	0/30/50/50	0/6/6/6

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	FAD	O4B-C1B	-7.57	1.31	1.41
2	G	6001	FAD	O4B-C1B	-7.16	1.32	1.41
2	E	4001	FAD	O4B-C1B	-7.13	1.32	1.41
2	C	2001	FAD	O4B-C1B	-6.96	1.32	1.41
2	F	5001	FAD	O4B-C1B	-6.91	1.32	1.41

The worst 5 of 145 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	4001	FAD	N3A-C2A-N1A	-10.93	120.52	128.89
2	A	1001	FAD	N3A-C2A-N1A	-10.61	120.77	128.89
2	F	5001	FAD	N3A-C2A-N1A	-10.54	120.83	128.89
2	C	2001	FAD	N3A-C2A-N1A	-10.43	120.91	128.89

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	7001	FAD	N3A-C2A-N1A	-10.36	120.96	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	FAD	4	0
2	C	2001	FAD	5	0
2	D	3001	FAD	5	0
2	E	4001	FAD	3	0
2	F	5001	FAD	2	0
2	G	6001	FAD	1	0
2	H	7001	FAD	1	0
2	I	8001	FAD	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	380/385 (98%)	-0.79	0 100 100	7, 18, 31, 37	0
1	C	380/385 (98%)	-0.68	0 100 100	8, 19, 31, 37	0
1	D	380/385 (98%)	-0.72	0 100 100	7, 23, 33, 39	0
1	E	380/385 (98%)	-0.71	0 100 100	8, 23, 33, 39	0
1	F	380/385 (98%)	-0.61	1 (0%) 94 94	10, 24, 33, 40	0
1	G	380/385 (98%)	-0.82	0 100 100	8, 20, 31, 37	0
1	H	380/385 (98%)	-0.79	0 100 100	8, 19, 30, 37	0
1	I	380/385 (98%)	-0.79	1 (0%) 94 94	9, 22, 32, 40	0
All	All	3040/3080 (98%)	-0.74	2 (0%) 95 95	7, 21, 32, 40	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	232	LEU	2.7
1	F	195	GLY	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	FAD	H	7001	53/53	0.98	0.10	0.44	19,22,27,30	0
2	FAD	G	6001	53/53	0.98	0.09	0.41	16,23,26,27	0
2	FAD	A	1001	53/53	0.97	0.10	0.23	15,20,22,23	0
2	FAD	I	8001	53/53	0.97	0.09	0.18	19,24,28,30	0
2	FAD	C	2001	53/53	0.97	0.09	0.18	17,21,24,25	0
2	FAD	D	3001	53/53	0.97	0.09	0.03	18,23,28,31	0
2	FAD	F	5001	53/53	0.97	0.09	-0.06	18,24,29,30	0
2	FAD	E	4001	53/53	0.97	0.08	-0.56	19,26,30,32	0

## 6.5 Other polymers

There are no such residues in this entry.