



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:18 PM GMT

PDB ID : 4EC4
Title : XIAP-BIR3 in complex with a potent divalent Smac mimetic
Authors : Mastrangelo, E.; Cossu, F.; Bolognesi, M.; Milani, M.
Deposited on : 2012-03-26
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	 70% 16% • 11%
1	B	122	 57% 27% • 13%
1	C	122	 66% 18% • 14%
1	D	122	 53% 29% • 14%
1	E	122	 58% 27% • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	122	<div><div></div><div>61%23%•14%</div></div>
1	G	122	<div>%<div><div></div><div>50%32%6%12%</div></div></div>
1	J	122	<div><div></div><div>59%25%•11%</div></div>
1	K	122	<div><div></div><div>63%20%•12%</div></div>
1	L	122	<div>%<div><div></div><div>54%31%•12%</div></div></div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9357 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Baculoviral IAP repeat-containing protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	0	0
			890	566	154	165	5			
1	B	106	Total	C	N	O	S	0	0	0
			865	553	148	159	5			
1	C	105	Total	C	N	O	S	0	0	0
			854	547	144	158	5			
1	D	105	Total	C	N	O	S	0	1	0
			862	552	147	158	5			
1	E	108	Total	C	N	O	S	0	0	0
			879	560	150	164	5			
1	J	108	Total	C	N	O	S	0	0	0
			879	560	150	164	5			
1	F	105	Total	C	N	O	S	0	0	0
			858	548	147	158	5			
1	G	107	Total	C	N	O	S	0	0	0
			870	556	149	160	5			
1	K	107	Total	C	N	O	S	0	0	0
			872	555	149	163	5			
1	L	107	Total	C	N	O	S	0	0	0
			870	556	149	160	5			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	357	HIS	-	EXPRESSION TAG	UNP P98170
A	358	HIS	-	EXPRESSION TAG	UNP P98170
A	359	HIS	-	EXPRESSION TAG	UNP P98170
A	360	HIS	-	EXPRESSION TAG	UNP P98170
A	361	HIS	-	EXPRESSION TAG	UNP P98170
A	362	HIS	-	EXPRESSION TAG	UNP P98170
B	357	HIS	-	EXPRESSION TAG	UNP P98170
B	358	HIS	-	EXPRESSION TAG	UNP P98170
B	359	HIS	-	EXPRESSION TAG	UNP P98170

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	360	HIS	-	EXPRESSION TAG	UNP P98170
B	361	HIS	-	EXPRESSION TAG	UNP P98170
B	362	HIS	-	EXPRESSION TAG	UNP P98170
C	357	HIS	-	EXPRESSION TAG	UNP P98170
C	358	HIS	-	EXPRESSION TAG	UNP P98170
C	359	HIS	-	EXPRESSION TAG	UNP P98170
C	360	HIS	-	EXPRESSION TAG	UNP P98170
C	361	HIS	-	EXPRESSION TAG	UNP P98170
C	362	HIS	-	EXPRESSION TAG	UNP P98170
D	357	HIS	-	EXPRESSION TAG	UNP P98170
D	358	HIS	-	EXPRESSION TAG	UNP P98170
D	359	HIS	-	EXPRESSION TAG	UNP P98170
D	360	HIS	-	EXPRESSION TAG	UNP P98170
D	361	HIS	-	EXPRESSION TAG	UNP P98170
D	362	HIS	-	EXPRESSION TAG	UNP P98170
E	357	HIS	-	EXPRESSION TAG	UNP P98170
E	358	HIS	-	EXPRESSION TAG	UNP P98170
E	359	HIS	-	EXPRESSION TAG	UNP P98170
E	360	HIS	-	EXPRESSION TAG	UNP P98170
E	361	HIS	-	EXPRESSION TAG	UNP P98170
E	362	HIS	-	EXPRESSION TAG	UNP P98170
J	357	HIS	-	EXPRESSION TAG	UNP P98170
J	358	HIS	-	EXPRESSION TAG	UNP P98170
J	359	HIS	-	EXPRESSION TAG	UNP P98170
J	360	HIS	-	EXPRESSION TAG	UNP P98170
J	361	HIS	-	EXPRESSION TAG	UNP P98170
J	362	HIS	-	EXPRESSION TAG	UNP P98170
F	357	HIS	-	EXPRESSION TAG	UNP P98170
F	358	HIS	-	EXPRESSION TAG	UNP P98170
F	359	HIS	-	EXPRESSION TAG	UNP P98170
F	360	HIS	-	EXPRESSION TAG	UNP P98170
F	361	HIS	-	EXPRESSION TAG	UNP P98170
F	362	HIS	-	EXPRESSION TAG	UNP P98170
G	357	HIS	-	EXPRESSION TAG	UNP P98170
G	358	HIS	-	EXPRESSION TAG	UNP P98170
G	359	HIS	-	EXPRESSION TAG	UNP P98170
G	360	HIS	-	EXPRESSION TAG	UNP P98170
G	361	HIS	-	EXPRESSION TAG	UNP P98170
G	362	HIS	-	EXPRESSION TAG	UNP P98170
K	357	HIS	-	EXPRESSION TAG	UNP P98170
K	358	HIS	-	EXPRESSION TAG	UNP P98170
K	359	HIS	-	EXPRESSION TAG	UNP P98170

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
K	360	HIS	-	EXPRESSION TAG	UNP P98170
K	361	HIS	-	EXPRESSION TAG	UNP P98170
K	362	HIS	-	EXPRESSION TAG	UNP P98170
L	357	HIS	-	EXPRESSION TAG	UNP P98170
L	358	HIS	-	EXPRESSION TAG	UNP P98170
L	359	HIS	-	EXPRESSION TAG	UNP P98170
L	360	HIS	-	EXPRESSION TAG	UNP P98170
L	361	HIS	-	EXPRESSION TAG	UNP P98170
L	362	HIS	-	EXPRESSION TAG	UNP P98170

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Zn 1 1	0	0
2	J	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	K	1	Total Zn 1 1	0	0
2	E	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	L	1	Total Zn 1 1	0	0
2	F	1	Total Zn 1 1	0	0

- Molecule 3 is (3S,6S,7S,9AS,3'S,6'S,7'S,9A'S)-N,N'-(BENZENE-1,4-DIYLBIS{BUTANE-4,1-DIYL-1H-1,2,3-TRIAZOLE-1,4-DIYL[(S)-PHENYLMETHANEDIYL]})BIS[7-(HYDROXYMETHYL)-6-{[(2S)-2-(METHYLAMINO)BUTANOYL]AMINO}-5-OXOOCTAHYDRO-1H-PYRROLO[1,2-A]AZEPINE-3-CARBOXAMIDE] (three-letter code: 0O6) (formula: C₆₄H₈₈N₁₄O₈).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			86	64	14	8		
3	B	1	Total	C	N	O	0	0
			86	64	14	8		
3	D	1	Total	C	N	O	0	0
			86	64	14	8		
3	E	1	Total	C	N	O	0	0
			86	64	14	8		
3	J	1	Total	C	N	O	0	0
			86	64	14	8		

- Molecule 4 is 3,6,9,12,15,18-HEXAOSAICOSANE-1,20-DIOL (three-letter code: P33) (formula: C₁₄H₃₀O₈).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			22	14	8		
4	D	1	Total	C	O	0	0
			22	14	8		
4	J	1	Total	C	O	0	0
			22	14	8		

- Molecule 5 is NONAETHYLENE GLYCOL (three-letter code: 2PE) (formula: C₁₈H₃₈O₁₀).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			28	18	10		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	G	1	Total	C	O	0	0
			28	18	10		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	16	Total	O	0	0
			16	16		
6	B	6	Total	O	0	0
			6	6		
6	C	9	Total	O	0	0
			9	9		
6	D	15	Total	O	0	0
			15	15		
6	E	10	Total	O	0	0
			10	10		
6	J	8	Total	O	0	0
			8	8		
6	F	11	Total	O	0	0
			11	11		
6	G	8	Total	O	0	0
			8	8		
6	K	5	Total	O	0	0
			5	5		
6	L	8	Total	O	0	0
			8	8		

Chain E: 




- Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain J: 




- Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain F: 




- Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain G: 




- Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain K: 



- Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain L: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.42Å 130.49Å 215.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.41 – 3.30 51.41 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (51.41-3.30) 99.8 (51.41-3.30)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.210 , 0.276 0.209 , 0.274	Depositor DCC
R_{free} test set	1446 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	40.5	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtriage
Outliers	2 of 28566 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	9357	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0O6, P33, ZN, 2PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/918	0.61	0/1244
1	B	0.45	0/893	0.59	0/1211
1	C	0.44	0/882	0.56	0/1197
1	D	0.45	0/893	0.56	0/1211
1	E	0.44	0/907	0.61	1/1230 (0.1%)
1	F	0.44	0/886	0.59	0/1201
1	G	0.46	0/898	0.56	0/1218
1	J	0.46	0/907	0.57	0/1230
1	K	0.41	0/900	0.57	0/1220
1	L	0.47	0/898	0.59	0/1218
All	All	0.45	0/8982	0.58	1/12180 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	352	LEU	CA-CB-CG	5.59	128.15	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	890	0	829	12	0
1	B	865	0	808	35	0
1	C	854	0	795	15	0
1	D	862	0	808	37	0
1	E	879	0	816	21	0
1	F	858	0	799	30	0
1	G	870	0	809	28	0
1	J	879	0	816	20	0
1	K	872	0	808	21	0
1	L	870	0	810	23	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	A	86	0	88	4	0
3	B	86	0	88	7	0
3	D	86	0	88	10	0
3	E	86	0	88	11	0
3	J	86	0	88	1	0
4	A	22	0	30	0	0
4	D	22	0	30	1	0
4	J	22	0	30	1	0
5	E	28	0	38	0	0
5	G	28	0	38	2	0
6	A	16	0	0	0	0
6	B	6	0	0	1	0
6	C	9	0	0	0	0
6	D	15	0	0	0	0
6	E	10	0	0	1	0
6	F	11	0	0	0	0
6	G	8	0	0	0	0
6	J	8	0	0	0	0
6	K	5	0	0	0	0
6	L	8	0	0	0	0
All	All	9357	0	8704	245	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 245 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:286[B]:ARG:HG3	1:D:286[B]:ARG:HH11	0.97	1.12
1:G:297:LYS:HG2	1:G:308:THR:HG23	1.34	1.08
1:F:248:ARG:HB3	1:F:249:ASN:HB3	1.37	1.06
1:F:248:ARG:HB3	1:F:249:ASN:CB	1.89	1.02
1:F:248:ARG:HB3	1:F:249:ASN:CA	1.91	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/122 (88%)	98 (92%)	9 (8%)	0	100	100
1	B	104/122 (85%)	100 (96%)	4 (4%)	0	100	100
1	C	103/122 (84%)	98 (95%)	4 (4%)	1 (1%)	19	58
1	D	104/122 (85%)	93 (89%)	11 (11%)	0	100	100
1	E	106/122 (87%)	95 (90%)	10 (9%)	1 (1%)	21	60
1	F	103/122 (84%)	95 (92%)	8 (8%)	0	100	100
1	G	105/122 (86%)	98 (93%)	5 (5%)	2 (2%)	10	45
1	J	106/122 (87%)	102 (96%)	4 (4%)	0	100	100
1	K	105/122 (86%)	96 (91%)	8 (8%)	1 (1%)	19	58
1	L	105/122 (86%)	97 (92%)	6 (6%)	2 (2%)	10	45
All	All	1048/1220 (86%)	972 (93%)	69 (7%)	7 (1%)	26	66

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	312	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	K	322	LYS
1	L	312	PRO
1	G	313	SER
1	L	275	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	94/106 (89%)	84 (89%)	10 (11%)	8	33
1	B	91/106 (86%)	82 (90%)	9 (10%)	10	37
1	C	90/106 (85%)	84 (93%)	6 (7%)	20	58
1	D	91/106 (86%)	82 (90%)	9 (10%)	10	37
1	E	93/106 (88%)	84 (90%)	9 (10%)	10	38
1	F	90/106 (85%)	77 (86%)	13 (14%)	4	19
1	G	91/106 (86%)	74 (81%)	17 (19%)	2	9
1	J	93/106 (88%)	80 (86%)	13 (14%)	4	20
1	K	92/106 (87%)	85 (92%)	7 (8%)	16	51
1	L	91/106 (86%)	79 (87%)	12 (13%)	5	22
All	All	916/1060 (86%)	811 (88%)	105 (12%)	7	29

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	279	VAL
1	F	253	SER
1	L	270	PHE
1	J	283	GLN
1	J	337	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	249	ASN
1	G	259	ASN
1	L	283	GLN
1	F	333	GLN
1	G	283	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 20 ligands modelled in this entry, 10 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	0O6	A	402	-	86,94,94	1.56	7 (8%)	94,130,130	1.36	12 (12%)
4	P33	A	403	-	21,21,21	0.51	0	20,20,20	0.33	0
3	0O6	B	402	-	86,94,94	1.55	8 (9%)	94,130,130	1.43	14 (14%)
3	0O6	D	402	-	86,94,94	1.53	7 (8%)	94,130,130	1.41	13 (13%)
4	P33	D	403	-	21,21,21	0.63	0	20,20,20	0.30	0
3	0O6	E	402	-	86,94,94	1.51	8 (9%)	94,130,130	1.75	17 (18%)
5	2PE	E	403	-	27,27,27	0.76	0	26,26,26	0.34	0
5	2PE	G	402	-	27,27,27	0.69	0	26,26,26	0.47	0
3	0O6	J	401	-	86,94,94	1.53	6 (6%)	94,130,130	1.41	10 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	P33	J	403	-	21,21,21	0.55	0	20,20,20	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	0O6	A	402	-	-	0/66/134/134	0/9/9/9
4	P33	A	403	-	-	0/19/19/19	0/0/0/0
3	0O6	B	402	-	-	0/66/134/134	0/9/9/9
3	0O6	D	402	-	-	0/66/134/134	0/9/9/9
4	P33	D	403	-	-	0/19/19/19	0/0/0/0
3	0O6	E	402	-	-	0/66/134/134	0/9/9/9
5	2PE	E	403	-	-	0/25/25/25	0/0/0/0
5	2PE	G	402	-	-	0/25/25/25	0/0/0/0
3	0O6	J	401	-	-	0/66/134/134	0/9/9/9
4	P33	J	403	-	-	0/19/19/19	0/0/0/0

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	401	0O6	NBX-NBZ	-7.16	1.23	1.34
3	B	402	0O6	NBW-NBY	-7.07	1.23	1.34
3	A	402	0O6	NBW-NBY	-7.06	1.23	1.34
3	A	402	0O6	NBX-NBZ	-7.02	1.24	1.34
3	D	402	0O6	NBX-NBZ	-6.99	1.24	1.34

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	0O6	CB-CA-C	-3.64	103.25	110.44
3	J	401	0O6	OAI-CCK-NDE	-3.57	117.23	121.67
3	E	402	0O6	CCJ-CCZ-NDF	-3.21	106.24	111.84
3	E	402	0O6	CCQ-CDA-NCC	-3.17	103.66	110.15
3	B	402	0O6	OAJ-CCL-NDF	-3.06	117.87	121.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 37 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	0O6	4	0
3	B	402	0O6	7	0
3	D	402	0O6	10	0
4	D	403	P33	1	0
3	E	402	0O6	11	0
5	G	402	2PE	2	0
3	J	401	0O6	1	0
4	J	403	P33	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	109/122 (89%)	-0.16	0 100 100	8, 16, 54, 75	0
1	B	106/122 (86%)	0.12	1 (0%) 85 82	22, 35, 52, 59	0
1	C	105/122 (86%)	-0.07	1 (0%) 84 80	19, 36, 56, 63	0
1	D	105/122 (86%)	-0.29	0 100 100	10, 20, 33, 59	0
1	E	108/122 (88%)	-0.30	0 100 100	12, 23, 40, 67	0
1	F	105/122 (86%)	-0.28	0 100 100	12, 28, 48, 59	0
1	G	107/122 (87%)	-0.09	1 (0%) 85 82	14, 31, 49, 80	0
1	J	108/122 (88%)	-0.26	0 100 100	9, 19, 38, 57	0
1	K	107/122 (87%)	0.18	0 100 100	19, 35, 56, 64	0
1	L	107/122 (87%)	0.27	1 (0%) 85 82	16, 37, 71, 79	0
All	All	1067/1220 (87%)	-0.09	4 (0%) 93 92	8, 27, 55, 80	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	292	LEU	2.3
1	B	270	PHE	2.2
1	C	276	ILE	2.1
1	G	260	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	P33	D	403	22/22	0.85	0.24	1.15	24,29,45,47	0
3	0O6	J	401	86/86	0.96	0.28	0.99	17,25,32,39	0
3	0O6	D	402	86/86	0.96	0.28	0.86	12,18,34,40	0
4	P33	J	403	22/22	0.91	0.20	0.52	24,24,28,29	0
4	P33	A	403	22/22	0.93	0.19	0.51	24,24,30,32	0
5	2PE	E	403	28/28	0.91	0.21	0.50	24,24,28,29	0
3	0O6	E	402	86/86	0.95	0.25	0.33	15,21,35,40	0
3	0O6	B	402	86/86	0.95	0.24	0.21	23,29,38,44	0
3	0O6	A	402	86/86	0.97	0.21	-0.03	11,15,33,36	0
5	2PE	G	402	28/28	0.94	0.17	-0.32	24,24,27,29	0
2	ZN	G	401	1/1	1.00	0.08	-1.64	16,16,16,16	0
2	ZN	C	401	1/1	0.99	0.12	-1.76	23,23,23,23	0
2	ZN	J	402	1/1	1.00	0.13	-1.77	9,9,9,9	0
2	ZN	F	401	1/1	0.99	0.10	-1.84	14,14,14,14	0
2	ZN	D	401	1/1	0.99	0.09	-2.27	11,11,11,11	0
2	ZN	L	401	1/1	0.99	0.05	-2.27	21,21,21,21	0
2	ZN	E	401	1/1	1.00	0.11	-2.40	13,13,13,13	0
2	ZN	B	401	1/1	0.96	0.09	-3.60	25,25,25,25	0
2	ZN	K	401	1/1	1.00	0.06	-6.39	22,22,22,22	0
2	ZN	A	401	1/1	0.99	0.08	-	8,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.